

**Ministry of Higher Education  
Higher Future Institute of Engineering and  
Technology in Mansoura  
Department of Mathematics and  
Basic Sciences**




---

# **Mathematics (III)**

**Dr. Htem Abd Elaziz Elagamy**

**Mathematics and statistics**



---

**Mathematics plays an important role in our practical and scientific life and its applications that have changed many interpretations of natural phenomena. In this course, we will study mathematics 1 and its engineering applications, and its importance lies in linking it to working life.**

**Dr. Htem Abd Elaziz Elagamy**

**Mathematics and statistics**

# Basic Results and Concepts

---

## I. GENERAL INFORMATION

### 1. Greek Letters Used

$\alpha$ alpha	$\theta$ theta	$\kappa$ kappa	$\tau$ tau
$\beta$ beta	$\phi$ phi	$\mu$ mu	$\chi$ chi
$\gamma$ gamma	$\psi$ psi	$\nu$ nu	$\omega$ omega
$\delta$ delta	$\xi$ xi	$\pi$ pi	$\Gamma$ cap. gamma
$\varepsilon$ epsilon	$\eta$ eta	$\rho$ rho	$\Delta$ cap. delta
$i$ iota	$\zeta$ zeta	$\sigma$ sigma	$\Sigma$ cap. sigma
	$\lambda$ lambda		

### 2. Some Notations

$\in$ belongs to	$\cup$ union	$\notin$ doesnot belong to
$\cap$ intersection	$\Rightarrow$ implies	$/$ such that
$\Leftrightarrow$ implies and implied by		

### 3. Unit Prefixes Used

Multiples and Submultiples	Prefixes	Symbols
$10^3$	kilo	k
$10^2$	hecto	h
10	deca	da
$10^{-1}$	deci*	d
$10^{-2}$	centi*	c
$10^{-3}$	milli	m
$10^{-6}$	micro	$\mu$

\* The prefixes 'deci' and 'centi' are only used with the metre, e.g., Centimeter is a recognized unit of length but Centigram is not a recognized unit of mass.

### 4. Useful Data

$e = 2.7183$	$1/e = 0.3679$	$\log_e 2 = 0.6931$	$\log_e 3 = 1.0986$
$\pi = 3.1416$	$1/\pi = 0.3183$	$\log_e 10 = 2.3026$	$\log_{10} e = 0.4343$
$\sqrt{2} = 1.4142$	$\sqrt{3} = 1.732$	$1 \text{ rad.} = 57^{\circ}17'45''$	$1^{\circ} = 0.0174 \text{ rad.}$

## 5. Systems of Units

Quantity	F.P.S. System	C.G.S. System	M.K.S. System
Length	foot (ft)	centimetre (cm)	metre (m)
Mass	pound (lb)	gram (gm)	kilogram (kg)
Time	second (sec)	second (sec)	second (sec)
Force	lb. wt.	dyne	newton (nt)

## 6. Conversion Factors

1 ft. = 30.48 cm = 0.3048 m	1m = 100 cm = 3.2804 ft.
1 ft <sup>2</sup> = 0.0929 m <sup>2</sup>	1 acre = 4840 yd <sup>2</sup> = 4046.77 m <sup>2</sup>
1ft <sup>3</sup> = 0.0283 m <sup>3</sup>	1 m <sup>3</sup> = 35.32 ft <sup>3</sup>
1 m/sec = 3.2804 ft/sec.	1 mile /h = 1.609 km/h.

## II. ALGEBRA

1. Quadratic Equation :  $ax^2 + bx + c = 0$  has roots

$$\alpha = \frac{-b + \sqrt{(b^2 - 4ac)}}{2a}, \quad \beta = \frac{-b - \sqrt{(b^2 - 4ac)}}{2a}$$

$$\alpha + \beta = -\frac{b}{a}, \quad \alpha\beta = \frac{c}{a}.$$

Roots are equal if  $b^2 - 4ac = 0$

Roots are real and distinct if  $b^2 - 4ac > 0$

Roots are imaginary if  $b^2 - 4ac < 0$

### 2. Progressions

(i) Numbers  $a, a + d, a + 2d, \dots$  are said to be in Arithmetic Progression (A.P.)

Its  $n$ th term  $T_n = a + (n - 1)d$  and sum  $S_n = \frac{n}{2} (2a + (n - 1)d)$

(ii) Numbers  $a, ar, ar^2, \dots$  are said to be in Geometric Progression (G.P.)

Its  $n$ th term  $T_n = ar^{n-1}$  and sum  $S_n = \frac{a(1 - r^n)}{1 - r}$ ,  $S_\infty = \frac{a}{1 - r}$  ( $r < 1$ )

(iii) Numbers  $1/a, 1/(a + d), 1/(a + 2d), \dots$  are said to be in Harmonic Progression (H.P.) (i.e., a sequence is said to be in H.P. if its reciprocals are in A.P. Its  $n$ th term  $T_n = 1/(a + (n - 1)d)$ .)

(iv) If  $a$  and  $b$  be two numbers then their

Arithmetic mean =  $\frac{1}{2} (a + b)$ , Geometric mean =  $\sqrt{ab}$ , Harmonic mean =  $2ab/(a + b)$

(v) Natural numbers are  $1, 2, 3, \dots, n$ .

$$\Sigma n = \frac{n(n + 1)}{2}, \quad \Sigma n^2 = \frac{n(n + 1)(2n + 1)}{6}, \quad \Sigma n^3 = \left\{ \frac{n(n + 1)}{2} \right\}^2$$

(vi) Stirling's approximation. When  $n$  is large  $n! \sim \sqrt{2\pi n} \cdot n^n e^{-n}$ .

### 3. Permutations and Combinations

$${}^n P_r = \frac{n!}{(n-r)!}; \quad {}^n C_r = \frac{n!}{r!(n-r)!} = \frac{{}^n P_r}{r!}$$

$$n_{C_{n-r}} = n_{C_r}, \quad n_{C_0} = 1 = n_{C_n}$$

### 4. Binomial Theorem

(i) When  $n$  is a positive integer

$$(1+x)^n = 1 + {}^n C_1 x + {}^n C_2 x^2 + {}^n C_3 x^3 + \dots + {}^n C_n x^n.$$

(ii) When  $n$  is a negative integer or a fraction

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1.2} x^2 + \frac{n(n-1)(n-2)}{1.2.3} x^3 + \dots \infty.$$

### 5. Indices

(i)  $a^m \cdot a^n = a^{m+n}$

(ii)  $(a^m)^n = a^{mn}$

(iii)  $a^{-n} = 1/a^n$

(iv)  $n \sqrt[n]{a}$  (i.e.,  $n$ th root of  $a$ ) =  $a^{1/n}$ .

### 6. Logarithms

(i) Natural logarithm  $\log x$  has base  $e$  and is inverse of  $e^x$ .

Common logarithm  $\log_{10} x = M \log x$  where  $M = \log_{10} e = 0.4343$ .

(ii)  $\log_a 1 = 0$ ;  $\log_a 0 = -\infty$  ( $a > 1$ );  $\log_a a = 1$ .


(iii)  $\log(mn) = \log m + \log n$ ;  $\log(m/n) = \log m - \log n$ ;  $\log(m^n) = n \log m$ .

### III. GEOMETRY

1. Coordinates of a point : Cartesian  $(x, y)$  and polar  $(r, \theta)$ .

Then  $x = r \cos \theta$ ,  $y = r \sin \theta$

or  $r = \sqrt{(x^2 + y^2)}$ ,  $\theta = \tan^{-1} \left( \frac{y}{x} \right)$ .



Distance between two points

$$(x_1, y_1) \text{ and } (x_2, y_2) = \sqrt{[(x_2 - x_1)^2 + (y_2 - y_1)^2]}$$

Points of division of the line joining  $(x_1, y_1)$  and  $(x_2, y_2)$  in the ratio  $m_1 : m_2$  is

$$\left( \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \right)$$

In a triangle having vertices  $(x_1, y_1)$ ,  $(x_2, y_2)$  and  $(x_3, y_3)$

$$(i) \text{ area} = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

(ii) Centroid (point of intersection of medians) is

$$\left( \frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3} \right)$$

(iii) Incentre (point of intersection of the internal bisectors of the angles) is

$$\left( \frac{ax_1 + bx_2 + cx_3}{a + b + c}, \frac{ay_1 + by_2 + cy_3}{a + b + c} \right)$$

where  $a, b, c$  are the lengths of the sides of the triangle.

(iv) Circumcentre is the point of intersection of the right bisectors of the sides of the triangle.

(v) Orthocentre is the point of intersection of the perpendiculars drawn from the vertices to the opposite sides of the triangle.

## 2. Straight Line

(i) Slope of the line joining the points  $(x_1, y_1)$  and  $(x_2, y_2) = \frac{y_2 - y_1}{x_2 - x_1}$

Slope of the line  $ax + by + c = 0$  is  $-\frac{a}{b}$  i.e.,  $-\frac{\text{coeff, of } x}{\text{coeff, of } y}$

(ii) Equation of a line:

(a) having slope  $m$  and cutting an intercept  $c$  on  $y$ -axis is  $y = mx + c$ .

(b) cutting intercepts  $a$  and  $b$  from the axes is  $\frac{x}{a} + \frac{y}{b} = 1$ .

(c) passing through  $(x_1, y_1)$  and having slope  $m$  is  $y - y_1 = m(x - x_1)$

(d) Passing through  $(x_1, y_1)$  and making an  $\angle\theta$  with the  $x$ -axis is

$$\frac{x - x_1}{\cos \theta} = \frac{y - y_1}{\sin \theta} = r$$

(e) through the point of intersection of the lines  $a_1x + b_1y + c_1 = 0$  and  $a_2x + b_2y + c_2 = 0$  is  $a_1x + b_1y + c_1 + k(a_2x + b_2y + c_2) = 0$

(iii) Angle between two lines having slopes  $m_1$  and  $m_2$  is  $\tan^{-1} \frac{m_1 - m_2}{1 - m_1 m_2}$

Two lines are parallel if

$$m_1 = m_2$$

Two lines are perpendicular if

$$m_1 m_2 = -1$$

Any line parallel to the line


$$ax + by + c = 0 \text{ is } ax + by + k = 0$$

Any line perpendicular to

$$ax + by + c = 0 \text{ is } bx - ay + k = 0$$

(iv) Length of the perpendicular from  $(x_1, y_1)$  of the line  $ax + by + c = 0$  is

$$\frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}}$$



### 3. Circle

(i) Equation of the circle having centre  $(h, k)$  and radius  $r$  is

$$(x - h)^2 + (y - k)^2 = r^2$$

(ii) Equation  $x^2 + y^2 + 2gx + 2fy + c = 0$  represents a circle having centre  $(-g, -f)$

and radius  $= \sqrt{g^2 + f^2 - c}$ .

(iii) Equation of the tangent at the point  $(x_1, y_1)$  to the circle  $x^2 + y^2 = a^2$  is  $xx_1 + yy_1 = a^2$ .

(iv) Condition for the line  $y = mx + c$  to touch the circle

$$x^2 + y^2 = a^2 \text{ is } c = a \sqrt{1 + m^2}.$$

(v) Length of the tangent from the point  $(x_1, y_1)$  to the circle

$$x^2 + y^2 + 2gx + 2fy + c = 0 \text{ is } \sqrt{(x_1^2 - y_1^2 + 2gx_1 + 2fy_1 + c)}.$$


### 4. Parabola

(i) Standard equation of the parabola is  $y^2 = 4ax$ .

Its parametric equations are  $x = at^2, y = 2at$ .

Latus - rectum  $LL' = 4a$ , Focus is  $S(a, 0)$

Directrix  $ZM$  is  $x + a = 0$ .




- (ii) Focal distance of any point  $P(x_1, y_1)$  on the parabola  $y^2 = 4ax$  is  $SP = x_1 + a$
- (iii) Equation of the tangent at  $(x_1, y_1)$  to the parabola  $y^2 = 4ax$  is  $yy_1 = 2a(x + x_1)$
- (iv) Condition for the line  $y = mx + c$  to touch the parabola  $y^2 = 4ax$  is  $c = a/m$ .
- (v) Equation of the normal to the parabola  $y^2 = 4ax$  in terms of its slope  $m$  is  $y = mx - 2am - am^3$ .

**5. Ellipse**

(i) Standard equation of the ellipse is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$





Its parametric equations are

$$x = a \cos \theta, \quad y = b \sin \theta.$$

$$\text{Eccentricity } e = \sqrt{1 - b^2 / a^2}.$$

$$\text{Latus - rectum } LSL' = 2b^2/a.$$

Foci S (-ae, 0) and S' (ae, 0)

Directrices ZM (x = -a/e) and Z'M' (x = a/e.)

(ii) Sum of the focal distances of any point on the ellipse is equal to the major axis i.e.,

$$SP + S'P = 2a.$$

(iii) Equation of the tangent at the point (x<sub>1</sub>, y<sub>1</sub>) to the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ is } \frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1.$$

(iv) Condition for the line y = mx + c to touch the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ is } c = \sqrt{a^2 m^2 + b^2}.$$

## 6. Hyperbola


(i) Standard equation of the hyperbola is

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1.$$

Its parametric equations are

$$x = a \sec \theta, \quad y = b \tan \theta.$$

$$\text{Eccentricity } e = \sqrt{1 + b^2 / a^2},$$



$$\text{Latus - rectum } LSL' = 2b^2/a.$$

Directrices ZM (x = a/e) and Z'M' (x = -a/e.)

(ii) Equation of the tangent at the point (x<sub>1</sub>, y<sub>1</sub>) to the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \text{ is } \frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1.$$

(iii) Condition for the line  $y = mx + c$  to touch the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \text{ is } c = \sqrt{(a^2m^2 - b^2)}$$

(iv) Asymptotes of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are  $\frac{x}{a} + \frac{y}{b} = 0$  and  $\frac{x}{a} - \frac{y}{b} = 0$ .

(v) Equation of the rectangular hyperbola with asymptotes as axes is  $xy = c^2$ . Its parametric equations are  $x = ct, y = c/t$ .

### 7. Nature of the a Conic

The equation  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  represents

(i) a pair of lines, if  $\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} (= \Delta) = 0$

(ii) a circle, if  $a = b, h = 0, \Delta \neq 0$

(iii) a parabola, if  $ab - h^2 = 0, c \Delta \neq 0$

(iv) an ellipse, if  $ab - h^2 > 0, \Delta \neq 0$

(v) a hyperbola, if  $ab - h^2 < 0, \Delta \neq 0$

and a rectangular hyperbola if in addition,  $a + b = 0$ .

### 8. Volumes and Surface Areas

Solid	Volume	Curved Surface Area	Total Surface Area
Cube (side a)	$a^3$	$4a^2$	$6a^2$
Cuboid (length l, breadth b, height h)	$lbh$	$2(l + b)h$	$2(lb + bh + hl)$
Sphere (radius r)	$\frac{4}{3} \pi r^3$	—	$4\pi r^2$
Cylinder (base radius r, height h)	$\pi r^2 h$	$2\pi r h$	$2\pi r (r + h)$
Cone	$\frac{1}{3} \pi r^2 h$	$\pi r l$	$\pi r (r + l)$

where slant height  $l$  is given by  $l = \sqrt{(r^2 + h^2)}$ .

#### IV. TRIGONOMETRY

1.

$\theta^\circ = 0$	0	30	45	60	90	180	270	360
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1	0	-1	0
$\cos \theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0	-1	0	1
$\tan \theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	$\infty$	0	$-\infty$	0

2. Any t-ratio of  $(n \cdot 90^\circ \pm \theta) = \pm$  same ratio of  $\theta$ , when  $n$  is even.

$= \pm$  co - ratio of  $\theta$ , when  $n$  is odd.

The sign + or - is to be decided from the quadrant in which  $n \cdot 90^\circ \pm \theta$  lies.

e.g.,  $\sin 570^\circ = \sin (6 \times 90^\circ + 30^\circ) = -\sin 30^\circ = -\frac{1}{2}$ ;

$\tan 315^\circ = \tan (3 \times 90^\circ + 45^\circ) = -\cot 45^\circ = -1$ .

3.  $\sin (A \pm B) = \sin A \cos B \pm \cos A \sin B$

$\cos (A \mp B) = \cos A \cos B \pm \sin A \sin B$

$\sin 2A = 2\sin A \cos A = 2 \tan A / (1 + \tan^2 A)$

$\cos 2A = \cos^2 A - \sin^2 A = 1 - 2 \sin^2 A = 2 \cos^2 A - 1 = \frac{1 - \tan^2 A}{1 + \tan^2 A}$ .

4.  $\tan (A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$ ;  $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$ .

5.  $\sin A \cos B = \frac{1}{2} [\sin (A + B) + \sin (A - B)]$

$\cos A \sin B = \frac{1}{2} [\sin (A + B) - \sin (A - B)]$

$\cos A \cos B = \frac{1}{2} [\cos (A + B) + \cos (A - B)]$

$\sin A \sin B = \frac{1}{2} [\cos (A - B) - \cos (A + B)]$ .

6.  $\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$

$\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$

$\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}$

$\cos C - \cos D = -2 \sin \frac{C+D}{2} \sin \frac{C-D}{2}$

7.  $a \sin x + b \cos x = r \sin (x + \theta)$

$$a \cos x + b \sin x = r \cos (x - \theta)$$

$$\text{where } a = r \cos \theta, b = r \sin \theta \text{ so that } r = \sqrt{a^2 + b^2}, \theta = \tan^{-1} \left( \frac{b}{a} \right)$$

8. In any  $\Delta ABC$ :

(i)  $a/\sin A = b/\sin B = c/\sin C$  (sine formula)

(ii)  $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$  . (cosine formula)

(iii)  $a = b \cos C + c \cos B$  (Projection formula)

(iv) Area of  $\Delta ABC = \frac{1}{2} bc \sin A = \sqrt{s(s-a)(s-b)(s-c)}$  where  $s = \frac{1}{2}(a+b+c)$ .

9. Series

(i) Exponential Series:  $e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \infty$

(ii)  $\sin x, \cos x, \sin hx, \cos hx$  series

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \infty,$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \infty$$

$$\sin hx = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots \infty, \quad \cos hx = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots \infty$$

(iii) Log series

$$\log (1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots \infty, \quad \log (1-x) = - \left( x + \frac{x^2}{2} + \frac{x^3}{3} + \dots \infty \right)$$

(iv) Gregory series

$$\tan^{-1} x = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots \infty, \quad \tan h^{-1} x = \frac{1}{2} \log \frac{1+x}{1-x} = x + \frac{x^3}{3} + \frac{x^5}{5} + \dots \infty.$$

10. (i) Complex number :  $z = x + iy = r (\cos \theta + i \sin \theta) = re^{i\theta}$

(ii) Euler's theorem:  $\cos \theta + i \sin \theta = e^{i\theta}$

(iii) Demoivre's theorem:  $(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$ .

11. (i) Hyperbolic functions:  $\sin hx = \frac{e^x - e^{-x}}{2}; \cos hx = \frac{e^x + e^{-x}}{2};$

$$\tan hx = \frac{\sin hx}{\cosh x}; \cot hx = \frac{\cos hx}{\sinh x}; \sec hx = \frac{1}{\cos hx}; \operatorname{cosec} hx = \frac{1}{\sin hx}$$

(ii) Relations between hyperbolic and trigonometric functions:

$$\sin ix = i \sin hx; \cos hx = \cos ix; \tan ix = i \tan hx.$$

(iii) Inverse hyperbolic functions;

$$\sin h^{-1}x = \log[x + \sqrt{x^2 + 1}]; \cosh^{-1}x = \log[x + \sqrt{x^2 - 1}]; \tan h^{-1}x = \frac{1}{2} \log \frac{1+x}{1-x}.$$

## V. CALCULUS

### 1. Standard limits:

$$(i) \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1},$$

$$(ii) \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

$n$  any rational number

$$(iii) \lim_{x \rightarrow 0} (1+x)^{1/x} = e$$

$$(iv) \lim_{x \rightarrow \infty} x^{1/x} = 1$$

$$(v) \lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \log_e a.$$

### 2. Differentiation

$$(i) \frac{d}{dx} (uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

$$\frac{d}{dx} \left( \frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

$$\frac{du}{dx} = \frac{du}{dy} \cdot \frac{dy}{dx} \text{ (chain Rule)}$$

$$\frac{d}{dx} (ax + b)^n = n(ax + b)^{n-1} \cdot a$$

$$(ii) \frac{d}{dx} (e^x) = e^x$$

$$\frac{d}{dx} (a^x) = a^x \log_e a$$

$$\frac{d}{dx} (\log_e x) = 1/x$$

$$\frac{d}{dx} (\log_a x) = \frac{1}{x \log a}.$$

$$(iii) \frac{d}{dx} (\sin x) = \cos x$$

$$\frac{d}{dx} (\cos x) = -\sin x$$

$$\frac{d}{dx} (\tan x) = \sec^2 x$$

$$\frac{d}{dx} (\cot x) = -\operatorname{cosec}^2 x$$

$$\frac{d}{dx} (\sec x) = \sec x \tan x$$

$$\frac{d}{dx} (\operatorname{cosec} x) = -\operatorname{cosec} x \cot x.$$

$$(iv) \frac{d}{dx} (\sin^{-1}x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} (\cos^{-1}x) = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} (\tan^{-1}x) = \frac{1}{1+x^2}$$

$$\frac{d}{dx} (\cot^{-1}x) = \frac{-1}{1+x^2}$$

$$\frac{d}{dx} (\sec^{-1}x) = \frac{1}{x\sqrt{x^2-1}}$$

$$\frac{d}{dx} (\operatorname{cosec}^{-1}x) = \frac{-1}{x\sqrt{x^2-1}}.$$

$$(v) \frac{d}{dx} (\sin h x) = \cos h x$$

$$\frac{d}{dx} (\cos h x) = \sin h x$$

$$\frac{d}{dx} (\tan h x) = \operatorname{sech}^2 x$$

$$\frac{d}{dx} (\cot h x) = -\operatorname{cosec} h^2 x.$$

$$(vi) D^n (ax + b)^m = m(m-1)(m-2) \dots (m-n+1) (ax + b)^{m-n} \cdot a^n$$

$$D^n \log(ax + b) = (-1)^{n-1} (n-1)! a^n / (ax + b)^n$$

$$D^n (e^{mx}) = m^n e^{mx}$$

$$D^n (a^{mx}) = m^n (\log a)^n \cdot a^{mx}$$

$$D^n \left[ \frac{\sin(ax + b)}{\cos(bx + c)} \right] = (a^2 + b^2)^{n/2} e^{ax} \left[ \frac{\sin(bx + c + n \tan^{-1} b/a)}{\cos(bx + c + n \tan^{-1} b/a)} \right].$$

(vii) Leibnitz theorem:  $(uv)_n$

$$= u_n + {}^n C_1 u_{n-1} v_1 + {}^n C_2 u_{n-2} v_2 + \dots + {}^n C_r u_{n-r} v_r + \dots + {}^n C_n v_n.$$

### 3. Integration

$$(i) \int x^n dx = \frac{x^{n+1}}{n+1} \quad (n \neq -1)$$

$$\int \frac{1}{x} dx = \log_e x$$

$$\int e^x dx = e^x$$

$$\int a^x dx = a^x / \log_e a$$

$$(ii) \int \sin x dx = -\cos x$$

$$\int \cos x dx = \sin x$$

$$\int \tan x dx = -\log \cos x$$

$$\int \cot x dx = \log \sin x$$

$$\int \sec x dx = \log(\sec x + \tan x) = \log \tan \left( \frac{\pi}{4} + \frac{x}{2} \right)$$

$$\int \operatorname{cosec} x dx = \log(\operatorname{cosec} x - \cot x) = \log \tan \left( \frac{x}{2} \right)$$

$$\int \sec^2 x dx = \tan x$$

$$\int \operatorname{cosec}^2 x dx = -\cot x.$$

$$(iii) \int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a}$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a}$$

$$\int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \log \frac{a+x}{a-x}$$

$$\int \frac{dx}{\sqrt{a^2 + x^2}} = \sinh^{-1} \frac{x}{a}$$

$$\int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \log \frac{x-a}{a+x}$$

$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \cosh^{-1} \frac{x}{a}.$$

$$(iv) \int \sqrt{a^2 - x^2} dx = \frac{x \sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a}$$

$$\int \sqrt{a^2 + x^2} dx = \frac{x \sqrt{a^2 + x^2}}{2} + \frac{a^2}{2} \sinh^{-1} \frac{x}{a} = \frac{x}{2} \sqrt{a^2 + x^2} + \frac{a^2}{2} \log \frac{x + \sqrt{a^2 + x^2}}{a}$$

$$\int \sqrt{x^2 - a^2} dx = \frac{x \sqrt{x^2 - a^2}}{2} + \frac{a^2}{2} \cosh^{-1} \frac{x}{a} = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log \frac{x + \sqrt{x^2 - a^2}}{a}$$

$$(v) \int e^{ax} \sin bx dx = \frac{e^{ax}}{a^2 + b^2} (a \sin bx - b \cos bx)$$

$$\int e^{ax} \cos bx \, dx = \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx)$$

$$(vi) \int \sin h x \, dx = \frac{\cos h x}{h}$$

$$\int \cos h x \, dx = \frac{\sin h x}{h}$$

$$\int \tan h x \, dx = \frac{\log \cos h x}{h}$$

$$\int \cot h x \, dx = \frac{\log \sin h x}{h}$$

$$\int \sec h^2 x \, dx = \frac{\tan h x}{h}$$

$$\int \operatorname{cosech}^2 x \, dx = -\frac{\cot h x}{h}$$

$$(vii) \int_0^{\pi/2} \sin^n x \, dx = \int_0^{\pi/2} \cos^n x \, dx$$

$$= \frac{(n-1)(n-3)(n-5)\dots}{n(n-2)(n-4)\dots} \times \left(\frac{\pi}{2}\right), \text{ only if } n \text{ is even}$$

$$\int_0^{\pi/2} \sin^m x \cos^n x \, dx = \frac{(m-1)(m-3)\dots \times (n-1)(n-3)\dots}{(m+n)(m+n-2)(m+n-4)\dots}$$

$$\times \left(\frac{\pi}{2}\right), \text{ only if both } m \text{ and } n \text{ are even}$$

$$(viii) \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx$$

$$\int_{-a}^a f(x) \, dx = 2 \int_0^a f(x) \, dx, \text{ if } f(x) \text{ is an even function.}$$

$$= 0, \text{ if } f(x) \text{ is an odd function.}$$

$$\int_0^{2a} f(x) \, dx = 2 \int_0^a f(x) \, dx, \text{ if } f(2a-x) = f(x)$$

$$= 0, \text{ if } f(2a-x) = -f(x).$$

## VI. Coordinate systems

	Polar coordinates (r, θ)	Cylindrical coordinates (ρ, φ, z)	Spherical polar coordinates (r, θ, φ)
Coordinate transformations	x = r cos θ y = r sin θ	x = ρ cos φ y = ρ sin φ z = z	x = r sin θ cos φ y = r sin θ sin φ z = r cos θ
Jacobian	$\frac{\partial(x, y)}{\partial(r, \theta)} = r$	$\frac{\partial(x, y, z)}{\partial(\rho, \phi, z)} = \rho$	$\frac{\partial(x, y, z)}{\partial(r, \theta, \phi)} = r^2 \sin \theta$
(Arc - length) <sup>2</sup>	(ds) <sup>2</sup> = (dr) <sup>2</sup> + r <sup>2</sup> (dθ) <sup>2</sup> dx dy = rdθ dr	(ds) <sup>2</sup> = (dρ) <sup>2</sup> + ρ <sup>2</sup> (dφ) <sup>2</sup> + (dz) <sup>2</sup>	(ds) <sup>2</sup> = (dr) <sup>2</sup> + r <sup>2</sup> (dθ) <sup>2</sup> + (r sin θ) <sup>2</sup> (dφ) <sup>2</sup>
Volume- element		dV = ρ dρ dφ dz	dV = r <sup>2</sup> sin θ dr dθ dφ

# Contents

Chapters	Page No.
Basic Results and Concepts	
<b>UNIT - I : Differential Equations</b>	
Chapter 1 : Basic Concepts of Differential Equations	3
Chapter 2 : Differential Equations of First Order and First Degree	9
Chapter 3 : Linear Differential Equations with Constant Coefficients and Applications	39
Chapter 4 : Equations Reducible To Linear Equations with Constant Coefficients	81
<b>Unit II : Series Solutions and Special Functions</b>	
Chapter 5 : Series Solutions and Special Functions	153
<b>Unit III : Laplace Transforms</b>	
Chapter 6 : Laplace Transforms	217
<b>Unit IV : Fourier Series and Partial Differential Equations</b>	
Chapter 7 : Fourier Series	291
Chapter 8 : Partial Differential Equations	331
<b>Unit V : Applications of Partial Differential Equations</b>	
Chapter 9 : Applications of Partial Differential Equations	401



---

**UNIT - I**  
**Differential Equations**

---

# Chapter 1

## Basic Concepts of Differential Equations

---

### INTRODUCTION

Differential equations are of fundamental importance in engineering mathematics because many physical laws and relations appear mathematically in the form of such equations. The mathematical formulation of many problems in science, engineering, Economics, sociology, physiology, Biology, Finance and management, give rise to differential equations. For example, the problem of motion of a satellite, the flow of current in an electric circuit, the growth of a population, the changes in price of commodities, decay of radioactive substance, cooling of a body etc. lead to differential equations. Each of the above problems are characterised by some laws which involve the rate of change of one or more quantities, with respect to the other quantities. The laws characterising these problems when expressed mathematically, become equations involving derivatives and such equations are called differential equations.

### DEFINITION

Any relation between known functions and an unknown function is called a differential equation if it involves the differential coefficient (or coefficients) of the unknown function.

It is usual to denote the unknown function by  $y$ . Finding the unknown function is called solving or integrating the differential equation. The solution or integral of the differential equation is also called its primitive, because the differential equation can be regarded as a relation derived from it.

Equations such as

$$(i) \quad (x^2 - y^2) \frac{dy}{dx} = xy \quad (1)$$

$$(ii) \quad \rho = \frac{\left\{ 1 + \left( \frac{dy}{dx} \right)^2 \right\}^{3/2}}{\frac{d^2y}{dx^2}} \quad (2)$$

$$(iii) \quad x^3 \frac{d^3y}{dx^3} + 2x^2 \frac{d^2y}{dx^2} + 3x \frac{dy}{dx} - 3y = x^2 + x \quad (3)$$

$$(iv) \quad x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = xyz \quad (4)$$

$$(v) \quad \frac{\partial^2 u}{\partial x^2} = K \frac{\partial^2 u}{\partial y^2} \quad (5)$$

$$(vi) \quad y = 2x \frac{dy}{dx} + y^2 \left( \frac{dy}{dx} \right)^3 \quad (6)$$

Which involve differential coefficients are called the differential equations.

Differential equations which involve only one independent variable are called ordinary differential equations. Equations (i) (ii), (iii) and (vi) are of this type.

Differential equations which involve two or more independent variables are called partial differential equations. Equations (iv) and (v) are of this type

**The order of a differential Equation.** The order of a differential equation is the order of the highest derivative involving in the equation.

**The Degree of a differential Equation.** The degree of a differential equation is the degree of the highest order derivative involving in the equation, When the equation is free from radicals and fractional powers.

**For example-** The differential equations

$$\frac{dy}{dx} + xy = a \quad (1)$$

$$\frac{d^2y}{dx^2} + x \frac{dy}{dx} = 2 \quad (2)$$

$$\left( \frac{d^3y}{dx^3} \right)^4 - 6x^2 \left( \frac{dy}{dx} \right)^2 + e^x = \sin xy \quad (3)$$

(U.P.T.U. 2009)

$$\frac{d^2y}{dx^2} = \left[ 1 + \left( \frac{dy}{dx} \right)^5 \right]^{-1/3} \quad (4)$$

### Basic Concepts of Differential Equations

The equation (1) is of the first order and first degree. Equation (2) is of second order and first degree. Equation (3) is of third order and fourth degree. Equation (4) is of second order and third degree

#### Formation of A Differential Equation

**Example 1.** Find the differential equation of the family of circles of radius  $r$  whose centre lies on the  $x$  axis. (I.A.S. 1993, 95, 96)

**Solution.** The equation of the circle with radius  $r$  and centre on  $x$  axis is

$$(x - a)^2 + y^2 = r^2 \quad (1)$$

Differentiating (1) with respect to  $x$ , we get

$$2(x - a) + 2y \frac{dy}{dx} = 0$$

Eliminating 'a' between (1) and (2), we get

$$\left(-y \frac{dy}{dx}\right)^2 + y^2 = r^2$$


or 
$$y^2 \left[ \left(\frac{dy}{dx}\right)^2 + 1 \right] = r^2$$

Which is the required differential equation.

**Example 2.** Find the differential equation of the family of parabolas with foci at the origin and axis along the  $x$ -axis. (I.A.S. 1994)

**Solution.** The equation of the parabolas with foci at the origin and axis along the  $x$  axis is given by

$$\sqrt{x^2 + y^2} = \frac{x + 2a}{\sqrt{1^2}}$$



or  $x^2 + y^2 = x^2 + 4ax + 4a^2$   
or  $y^2 = 4a(x + a)$  (1)

Differentiating with respect to x, we get

$$2y \frac{dy}{dx} = 4a$$
$$\Rightarrow y \frac{dy}{dx} = 2a$$
 (2)

Eliminating a between (1) and (2), we get

$$y^2 = 2y \frac{dy}{dx} \left[ x + \frac{1}{2} y \frac{dy}{dx} \right]$$

or  $y \left( \frac{dy}{dx} \right)^2 + 2x \frac{dy}{dx} - y = 0$

which is the required differential equation.

**Example 3.** Determine the differential equation whose set of independent solution is  $\{e^x, xe^x, x^2 e^x\}$

(U.P.T.U. 2002)

**Solution.** Here we have

$$y = C_1 e^x + C_2 x e^x + C_3 x^2 e^x$$
 (1)

Differentiating both sides of (1) w.r.t "x" we get

$$y' = C_1 e^x + C_2 x e^x + C_2 e^x + C_3 x^2 e^x + C_3 2x e^x$$
$$\Rightarrow y' = y + C_2 e^x + 2 C_3 x e^x$$
$$\Rightarrow y' - y = C_2 e^x + 2 C_3 x e^x$$
 (2)

Again differentiating both sides, we get

$$y'' - y' = C_2 e^x + 2 C_3 x e^x + 2 C_3 e^x$$
$$\Rightarrow y'' - y' = y' - y + 2 C_3 e^x \quad \text{using (2)}$$
$$\Rightarrow y'' - 2y' + y = 2C_3 e^x$$
 (3)

Again differentiating (3), we get

$$y''' - 2y'' + y' = 2 C_3 e^x$$
$$\Rightarrow y''' - 2y'' + y' = y'' - 2y' + y \quad \text{using (3)}$$

Basic Concepts of Differential Equations

$$\Rightarrow y''' - 3y'' + 3y' - y = 0$$

$$\text{or } \frac{d^3y}{dx^3} - 3 \frac{d^2y}{dx^2} + 3 \frac{dy}{dx} - y = 0$$

is the required differential equation

### EXERCISE

1. Form the differential equation of simple harmonic motion given by  $x = A \cos (nt + \alpha)$

$$\text{Ans. } \frac{d^2x}{dt^2} + n^2x = 0$$

2. Obtains the differential equation of all circles of radius  $a$  and centre  $(h, k)$  and hence prove that the radius of curvature of a circle at any point is constant.

$$\text{Ans. } \left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^3 = a^2 \left( \frac{d^2y}{dx^2} \right)^2$$

3. Show that  $v = \frac{A}{r} + B$  is a solution of  $\frac{d^2v}{dr^2} + \frac{2}{r} \frac{dv}{dr} = 0$

4. Show that  $Ax^2 + By^2 = 1$  is the solution of  $x \left\{ y \frac{d^2y}{dx^2} + \left( \frac{dy}{dx} \right)^2 \right\} = y \frac{dy}{dx}$

5. By eliminating the constants  $a$  and  $b$  obtain differential equation of which  $xy = ae^x + be^{-x} + x^2$  is a constant.

### Objective Type of Questions

Each question possesses four alternative answers, but only one answer is correct tick mark the correct one.

1. Degree and order of the differential equation  $\sqrt{2 \left( \frac{dy}{dx} \right)^3 + 4} = \left( \frac{d^2y}{dx^2} \right)^{3/2}$  are respectively.

- (a) order 2, degree 3                      .(b)            order 1, degree 3  
(c) order 3, degree 2                      (d)            order 3, degree 1

Ans. (a)

2. The degree of the differential equation  $\left[ y + x \left( \frac{d^2y}{dx^2} \right)^2 \right]^{1/4} = \frac{d^3y}{dx^3}$  is given by

- (a) 2                      (b) 3                      (c) 4                      (d) 1

(R.A.S. 1993)

Ans. (c)

3. The order of the differential equation  $\left[ 1 + \left( \frac{d^3y}{dx^3} \right)^2 \right]^{4/3} = \frac{d^2y}{dx^2}$  is given by

- (a) 1                      (b) 2                      (c) 3                      (d) 4

(R.A.S. 1993)

Ans. (c)

4. If  $x = A \cos (mt - \alpha)$  then the differential equation satisfying this relation is

(a)  $\frac{dx}{dt} = 1 - x^2$                       (b)  $\frac{d^2x}{dt^2} = -\alpha^2 x$

(c)  $\frac{d^2x}{dt^2} = -m^2 x$                       (d)  $\frac{dx}{dt} = -m^2 x$

(I.A.S. 1993)

5. The equation  $y \frac{dy}{dx} = x$  represents a family of

- (a) Circles                      (b) hyperbola  
(c) parabolas                      (d) ellipses

(U.P.P.C.S. 1995)

Ans. (b)

## Chapter 2

# Differential Equations of First Order and First Degree

---

### INTRODUCTION

An equation of the form  $F\left(x, y, \frac{dy}{dx}\right) = 0$  in which  $x$  is the independent variable and  $\frac{dy}{dx}$  appears with first degree is called a first order and first degree differential equation. It can also be written in the form  $\frac{dy}{dx} = f(x, y)$  or in the form  $Mdx + Ndy = 0$ , where  $M$  and  $N$  are functions of  $x$  and  $y$ . Generally, it is difficult to solve the first order differential equations and in some cases they may not possess any solution. There are certain standard types of first order, first degree equations. In this chapter we shall discuss the methods of solving them.

### VARIABLES SEPARABLE

If the equation is of the form  $f_1(x) dx = f_2(y) dy$  then, its solution, by integration is  $\int f_1(x) dx = \int f_2(y) dy + C$

where  $C$  is an arbitrary constant.

**Example 1.** Solve  $\frac{dy}{dx} = e^{x-y} + x^2 e^{-y}$

**Solution.** The given equation is

$$\frac{dy}{dx} = e^{x-y} + x^2 e^{-y}$$

or  $e^y dy = (e^x + x^2) dx$

Integrating,  $e^y = e^x + \frac{x^3}{3} + c$ , where  $C$  is an arbitrary constant is the required solution.

**Example 2.** Solve  $\left(y - x \frac{dy}{dx}\right) = a \left(y^2 + \frac{dy}{dx}\right)$



**Solution.** The given equation is

$$y - x \frac{dy}{dx} = ay^2 + a \frac{dy}{dx}$$

or  $(a + x) \frac{dy}{dx} = (y - ay^2)$

or  $\frac{dy}{y - ay^2} = \frac{dx}{a + x}$

or  $\frac{dy}{y - ay^2} = \frac{dx}{a + x}$

or  $\frac{dy}{y(1 - ay)} = \frac{dx}{x + a}$

or  $\left[ \frac{a}{1 - ay} + \frac{1}{y} \right] dy = \frac{dx}{x + a}$ , resolving into partial fractions.

Integrating,  $[-\log(1 - ay) + \log y] = \log(x + a) + \log C$

where C is an arbitrary constant

or  $\log \left( \frac{y}{1 - ay} \right) = \log \{C(x + a)\}$

or  $\frac{y}{1 - ay} = C(x + a)$

or  $y = C(x + a)(1 - ay)$  is the required solution.

**Example 2.** Solve  $(x + y)^2 \frac{dy}{dx} = a^2$

**Solution.** Let  $x + y = v$

then from (1) on differentiating with respect to x, we have

$$1 + \frac{dy}{dx} = \frac{dv}{dx}$$

or  $\frac{dy}{dx} = \left( \frac{dv}{dx} - 1 \right)$

Substituting these values from (1) and (2) in the given equation, we get

Differential Equations of First Order and First Degree

$$v^2 \left[ \frac{dv}{dx} - 1 \right] = a^2$$

or  $v^2 \frac{dv}{dx} = a^2 + v^2$

or  $\frac{v^2}{a^2 + v^2} dv = dx$

or  $\frac{a^2 + v^2 - a^2}{a^2 + v^2} dv = dx$

or  $\left[ 1 - \frac{a}{a^2 + v^2} \right] dv = dx$

Integrating,

$$v - a^2 \frac{1}{a} \tan^{-1} \left( \frac{v}{a} \right) = x + c$$

where c is an arbitrary constant

or  $v - a \tan^{-1} (v/a) = x + c$

or  $(x + y) - a \tan \{(x + y)/a\} = x + c$ , from (1)

or  $y - a \tan^{-1} \{(x + y)/a\} = C$

**Method of Solving Homogeneous Differential Equation**

It is a differential equation of the form

$$\frac{dy}{dx} = \frac{\phi(x, y)}{\psi(x, y)} \tag{1}$$

where  $\phi(x, y)$  and  $\psi(x, y)$  are homogeneous functions of x and y at the same degree, n (say)

Such equation can be solved by putting  $y = vx$

where  $\frac{dy}{dx} = v + x \frac{dv}{dx}$  (2)

Now the given differential equation is

$$\frac{dy}{dx} = \frac{\phi(x, y)}{\psi(x, y)} = \frac{x^n \phi(y/x)}{x^n \psi(y/x)}$$

or 
$$\frac{dy}{dx} = \frac{\phi(y/x)}{\psi(y/x)} = f\left(\frac{y}{x}\right), \text{ say} \quad (3)$$

Substituting the values of  $y$  and  $\frac{dy}{dx}$  from, (1) and (2) in (3),

we have 
$$v + x \frac{dv}{dx} = f(v)$$

or 
$$x \frac{dv}{dx} = f(v) - v$$

or 
$$\frac{dv}{f(v) - v} = \frac{dx}{x}$$

The variables has now been separated and its solution is

$$\int \frac{dv}{f(v) - v} = \log x + c \text{ where } c \text{ is an arbitrary constant.}$$

After integration  $v$  should be replaced by  $(y/x)$  to get the required solution.

**Example 4.** Solve  $x dy - y dx = \sqrt{(x^2 + y^2)} dx$

**Solution.** The given equation can be rewritten as

$$x \frac{dy}{dx} - y = \sqrt{(x^2 + y^2)}$$

or 
$$x \left( v + x \frac{dv}{dx} \right) - vx = \sqrt{(x^2 + v^2 x^2)}, \text{ putting } y = vx$$

or 
$$x \frac{dv}{dx} = \sqrt{(1 + v^2)}$$

or 
$$\frac{dv}{\sqrt{(1 + v^2)}} = \frac{dx}{x}$$

Integrating,  $\log \left\{ v + \sqrt{(v^2 + 1)} \right\} = \log x + \log c$ , where  $c$  is an arbitrary constant

or 
$$\left\{ v + \sqrt{(v^2 + 1)} \right\} = Cx$$

or 
$$\left[ y + \sqrt{(y^2 + x^2)} \right] / x = Cx \quad \because v = y/x$$

Differential Equations of First Order and First Degree

or 
$$y + \sqrt{(y^2 + x^2)} = Cx^2$$

**Example 5.** Solve  $(1 + e^{x/y}) dx + e^{x/y} [1 - (x/y)] dy = 0$  (U.P.P.C.S. 1999)

**Solution.** The given equation may be rewritten as

$$e^{x/y} \left( 1 - \frac{x}{y} \right) + (1 + e^{x/y}) \frac{dx}{dy} = 0$$

or 
$$e^v (1 - v) + (1 + e^v) \left( v + y \frac{dv}{dy} \right) = 0$$

putting  $x = vy$  or  $\frac{dx}{dy} = v + y \frac{dv}{dy}$

or 
$$e^v - ve^v + v + ve^v + (1 + e^v) y \frac{dv}{dy} = 0$$

or 
$$(v + e^v) + (1 + e^v) y \frac{dv}{dy} = 0$$

or 
$$\frac{(1 + e^v)}{v + e^v} dv + \frac{dy}{y} = 0$$

Integrating,  $\log (v + e^v) + \log y = \log C$ , when C is an arbitrary constant

or 
$$\log \{(v + e^v) y\} = \log C$$

or 
$$(v + e^v) y = C$$

or 
$$\left[ \frac{x}{y} + e^{x/y} \right] y = C, \text{ putting } v = x/y$$

or 
$$(x + y e^{x/y}) = C$$

**Equations Reducible to Homogeneous Form.**

Consider the equation 
$$\frac{dy}{dx} = \frac{ax + by + C}{a'x + b'y + C'} \quad (1)$$

where  $a:b \neq a':b'$

If C and C' are both zero, the equation is homogenous and can be solved by the method of homogeneous equation. If C and C' are not both zero, we change the variables so that constant terms are no longer present, by the substitutions  $x = X + h$  and  $y = Y + k$  Where h and k are constants yet to be chosen. (2)

From (2)  $dx = dX$  and  $dy = dY$

and so (1) reduces to  $\frac{dY}{dX} = \frac{a(X+h) + b(Y+k) + C}{a'(X+h) + b'(Y+k) + C'}$

or 
$$\frac{dY}{dX} = \frac{aX + bY + (ah + bk + c)}{a'X + b'Y + (a'h + b'k + C')} \quad (3)$$

Now, choose  $h$  and  $k$  such that

$$ah + bk + c = 0 \text{ and } a'h + b'k + c' = 0 \quad (4)$$

Then (3) reduces to  $\frac{dY}{dX} = \frac{aX + bY}{a'X + b'Y}$ , which is homogeneous and can be solved by the substitution  $Y = vX$ . Replacing  $X$  and  $Y$  in the solution so obtained by  $x-h$  and  $y-k$  [from (3)] respectively

We can get the required solution is terms of  $x$  and  $y$ , the original variables.

#### A Special Case When $a:b = a':b'$

In this case we cannot solve the equations given by (4) above and the differential equation is of the form

$$\frac{dy}{dx} = \frac{ax + by + C}{kax + kby + C'} \quad (5)$$

In this case the differential equation is solve by putting

$$v = ax + by \quad (6)$$

Differentiating both sides of (6) with respect to  $x$ , we get

$$\frac{dv}{dx} = a + b \frac{dy}{dx} \text{ or } \frac{dy}{dx} = \frac{1}{b} \left( \frac{dv}{dx} - a \right)$$

$\therefore$  The equation (5) reduces to

$$\frac{1}{b} \left( \frac{dv}{dx} - a \right) = \frac{v+C}{kv + C'} \text{, or } \frac{dv}{dx} = a + \frac{b(v+C)}{kv+C'}$$

The variables, are now separable, and we can determine  $v$  in terms of  $x$ . Replacing  $v$  by  $ax + by$  in this solution, we can obtain the final solution.

**Example 6.** Solve  $(2x + y - 3) dy = (x + 2y - 3) dx$

**Solution.** The given differential equation is  $\frac{dy}{dx} = \frac{x + 2y - 3}{2x + y - 3} \quad (1)$

Differential Equations of First Order and First Degree

putting  $x = X + h$  and  $y = Y + k$  (2)

the equation (1) reduce to  $\frac{dY}{dX} = \frac{(X + h) + 2(Y + k) - 3}{2(X + h) + (Y + k) - 3}$

or  $\frac{dY}{dX} = \frac{X + 2Y + (h + 2K - 3)}{2X + Y + (2h + K - 3)}$  (3)

Choose  $h$  and  $k$  such that

$$h + 2k - 3 = 0 \text{ and } 2h + k - 3 = 0 \quad (4)$$

Solving the equations (4) we have  $h = 1 = k$

$\therefore$  From (2)  $x = X + 1$  and  $y = Y + 1$

or  $X = x - 1$  and  $Y = y - 1$  (5)

Also (3) reduce to  $\frac{dY}{dX} = \frac{X + 2Y}{2X + Y}$

putting  $Y = vX$ ,  $v + X \frac{dv}{dX} = \frac{X + 2vX}{2X + vX}$

or  $v + X \frac{dv}{dX} = \frac{1 + 2v}{2 + v}$

or  $X \frac{dv}{dX} = \frac{1 + 2v}{2 + v} - v = \frac{1 + 2v - 2v - v^2}{2 + v} = \frac{1 - v^2}{2 + v}$

or  $\frac{2 + v}{1 - v^2} dv = \frac{dX}{X}$

or  $\left[ \frac{1}{2} \frac{1}{1 + v} + \frac{3}{2} \frac{1}{1 - v} \right] dv = \frac{dX}{X}$

Integrating,  $\frac{1}{2} \log(1 + v) - \frac{3}{2} \log(1 - v) = \log X + \log C$ , where  $C$  is an arbitrary constant

or  $\log \left[ \frac{1 + v}{(1 - v)^3} \right] = 2 \log(CX) = \log(CX)^2$

or  $\frac{1 + v}{(1 - v)^3} = (CX)^2$

or 
$$\frac{1 + Y/k}{(1 - Y/k)^3} = C^2 X^2$$

or 
$$\frac{X + Y}{(X - Y)^3} = C^2$$

or 
$$\frac{(x - 1) + (y - 1)}{\{(x - 1) - (y - 1)\}^3} = C^2 \text{ from (5)}$$

or 
$$\frac{x + y - 2}{(x - y)^3} = C^2$$

**Example 7.** Solve  $(2x + 2y + 3) dy - (x + y + 1) dx = 0$

**Solution.** The given equation is 
$$\frac{dy}{dx} = \frac{x + y + 1}{2x + 2y + 3} \quad (1)$$

put  $x + y + 1 = v$

• Differentiating

$$1 + \frac{dy}{dx} = \frac{dv}{dx} \text{ or } \frac{dy}{dx} = \frac{dv}{dx} - 1$$

Therefore (1) reduce to 
$$\frac{dv}{dx} - 1 = \frac{v}{2v + 1}$$

or 
$$\frac{dv}{dx} = \frac{3v + 1}{2v + 1}$$

or 
$$\frac{2v + 1}{3v + 1} dv = dx$$

or 
$$\left( \frac{2}{3} + \frac{1}{3(3v + 1)} \right) dv = dx$$

Integrating,  $\frac{2}{3} v + \frac{1}{9} \log(3v + 1) = x + c$ , where C is an arbitrary constant

or 
$$6v + \log(3v + 1) = 9x + C_1, \text{ where } C_1 = 9C$$

or 
$$6(x + y + 1) + \log(3x + 3y + 4) = 9x + C_1$$

or 
$$6y - 3x + \log(3x + 3y + 4) = C_2 \text{ where } C_2 = C_1 + 6$$

Differential Equations of First Order and First Degree

**Linear Differential Equations**

A differential equation of the form  $\frac{dy}{dx} + Py = Q$

where P and Q are constants or functions of x alone (and not of y) is called a linear differential equation of the first order in y

its integrating factor =  $e^{\int P dx}$

Multiplying both sides of (1) by this integrating factor (I.F.) and then integrating we get

$y \cdot e^{\int P dx} = C + \int Q \cdot e^{\int P dx} dx$ , where C is an arbitrary constant, is the complete solution of (1)

**Example 8.** Solve  $\frac{dy}{dx} + 2y \tan x = \sin x$ , given that  $y = 0$  when  $x = \pi/3$ .

(U.P.P.C.S. 2003)

**Solution.** Here  $P = 2 \tan x$  and  $Q = \sin x$

$$\begin{aligned} \therefore \text{Integrating factor} &= e^{\int P dx} = e^{\int 2 \tan x dx} \\ &= e^{2 \log \sec x} \\ &= e^{\log (\sec x)^2} = \sec^2 x \end{aligned}$$

Multiplying the given equation by  $\sec^2 x$ , we get

$$\sec^2 x \left( \frac{dy}{dx} + 2y \tan x \right) = \sin x \sec^2 x$$

$$\text{or } \frac{d}{dx} (y \sec^2 x) = \sec x \tan x$$

Integrating both sides with respect to x, we get

$$y \sec^2 x = C + \int \sec x \tan x dx, \text{ where } C \text{ is an arbitrary constant}$$

$$\text{or } y \sec^2 x = C + \sec x \tag{1}$$

it is given that when  $x = \pi/3, y = 0$

$$\therefore \text{from (1) } 0 \times \sec^2 \frac{\pi}{3} = C + \sec \frac{\pi}{3}$$



or  $0 = C + 2 \quad \therefore \sec \frac{\pi}{3} = 2$

$\therefore$  from (1) the required solution is

$$y \sec^2 x = -2 + \sec x$$

or  $y = -2 \cos^2 x + \cos x$

**Example 9.** Solve  $(1 + y^2) dx + (x - e^{\tan^{-1}y}) dy = 0$

(Bihar P.C.S. 2002, I.A.S. 2006)

**Solution.** The given equation can be written as

$$\frac{dx}{dy} + \frac{x}{1 + y^2} = \frac{e^{\tan^{-1}y}}{1 + y^2}$$

Therefore the integrating factor  $= e^{\int \frac{1}{1 + y^2} dy}$   
 $= e^{\tan^{-1}y}$

Multiplying both sides of (1) by the integrating factor and integrating, we have

$$x \cdot e^{\tan^{-1}y} = C + \int \frac{e^{\tan^{-1}y}}{1 + y^2} \times e^{\tan^{-1}y} dy$$

where C is an arbitrary constant

or  $x e^{\tan^{-1}y} = C + \int e^{2t} dt$ , where  $t = \tan^{-1}y$   
 $= C + \frac{1}{2} e^{2t}$

or  $x \cdot e^{\tan^{-1}y} = C + \frac{1}{2} e^{2 \tan^{-1}y}$

**Example 10.** Solve  $\frac{dy}{dx} + y \cos x = \frac{1}{2} \sin 2x$  (I.A.S. 2004)

**Solution.** Here  $P = \cos x$  and  $Q = \frac{1}{2} \sin 2x = \sin x \cos x$

$\therefore$  Integrating factor  $= e^{\int P dx} = e^{\int \cos x dx} = e^{\sin x}$

Multiplying the given equation by the integrating factor  $e^{\sin x}$  and integrating with respect to x, we get

Differential Equations of First Order and First Degree

$y \cdot e^{\sin x} = C + \int e^{\sin x} \sin x \cos x \, dx$ , where  $C$  is an arbitrary constant.

or 
$$y \cdot e^{\sin x} = C + \int e^t t \, dt, \quad \text{where } t = \sin x$$
$$= C + t \cdot e^t - e^t$$
$$= C + e^{\sin x} (\sin x - 1)$$

or 
$$y \cdot e^{\sin x} = C + e^{\sin x} (\sin x - 1)$$

**Equations Reducible to the Linear Form.**

The equation

$$\frac{dy}{dx} + Py = Qy^n \tag{1}$$

Where  $P$  and  $Q$  are constants or functions of  $x$  alone and  $n$  is a constant other than zero or unity is called the extended form of linear equation or Bernoulli's Equation.

This type of equation can be reduced to the linear form on dividing by  $y^n$  and putting  $\frac{1}{y^{n-1}}$  equal to  $v$

Dividing (1) by  $y^n$ , we get 
$$\frac{1}{y^n} \frac{dy}{dx} + P \cdot \frac{1}{y^{n-1}} = Q \tag{2}$$

Put  $\frac{1}{y^{n-1}} = v$  or  $y^{-n+1} = v$

Differentiating both sides with respect to  $x$ , get

$$(-n+1) y^{-n} \frac{dy}{dx} = \frac{dv}{dx}$$

or 
$$\frac{1}{y^n} \frac{dy}{dx} = \frac{1}{1-n} \frac{dv}{dx}$$

Making these substitutions in (2), we have

$$\frac{1}{(1-n)} \frac{dv}{dx} + Pv = Q$$

or 
$$\frac{dv}{dx} + P(1-n)v = Q(1-n)$$

which is linear in  $v$  and can be solve by method of linear differential equation.

**Example 11.** Solve  $\frac{dy}{dx} + x \sin 2y = x^3 \cos^2 y$  (U.P.P.C.S. 1994)

**Solution.** Dividing both sides of the given equation by  $\cos^2 y$  we get

$$\sec^2 y \frac{dy}{dx} + x (2 \tan y) = x^3$$

putting  $\tan y = v$  or  $\sec^2 y \frac{dy}{dx} = \frac{dv}{dx}$

the above equation reduces to  $\frac{dv}{dx} + 2xv = x^3$  (1)

which is a linear equation. whose integrating factor

$$= e^{\int 2x dx} = e^{x^2}$$

Multiplying both sides of (1) by the integrating factor and integrating, we have

$$v \cdot e^{x^2} = C + \int x^3 e^{x^2} dx, \text{ where } C \text{ is an arbitrary constant}$$

or 
$$v \cdot e^{x^2} = C + \frac{1}{2} \int x^2 e^{x^2} 2x dx$$

$$= C + \frac{1}{2} \int t \cdot e^t dt, \text{ where } t = x^2$$

$$= C + \frac{1}{2} (t e^t - \int e^t dt)$$

or 
$$e^{x^2} \tan y = C + \frac{1}{2} e^t (t - 1) \quad \because v = \tan y$$

or 
$$e^{x^2} \tan y = C + \frac{1}{2} e^{x^2} (x^2 - 1)$$

or 
$$2 \tan y = 2 C e^{-x^2} + (x^2 - 1)$$

Differential Equations of First Order and First Degree

**Example 2.** Solve  $\frac{dz}{dx} + \frac{z}{x} \log z = \frac{z}{x^2} (\log z)^2$

(I.A.S. 2001, U.P.P.C.S., 1999)

**Solution.** Dividing both sides of the given equation by  $z (\log z)^2$ , we get

$$\frac{1}{z (\log z)^2} \frac{dz}{dx} + \frac{1}{(\log z)} \frac{1}{x} = \frac{1}{x^2}$$

putting  $\frac{1}{\log z} = v$  or  $-\frac{1}{z (\log z)^2} \frac{dz}{dx} = \frac{dv}{dx}$ , the above equation reduces to

$$-\frac{dv}{dx} + \frac{v}{x} = \frac{1}{x^2}$$

or  $\frac{dv}{dx} - \frac{1}{x} v = -\frac{1}{x^2}$ , which is linear equation in  $v$  and whose integrating factor

$$= e^{-\int \frac{1}{x} dx} = e^{-\log x} = e^{\log x^{-1}}$$

$$= \frac{1}{x}$$

Proceeding in the usual way, the solution of above equation is

$$v \frac{1}{x} = C - \int \frac{1}{x^2} \frac{1}{x} dx$$

or  $\frac{1}{x (\log z)} = C + \frac{1}{2x^2}$

**Example 13.** Solve  $y \sin 2x dx - (1 + y^2 + \cos^2 x) dy = 0$

(I.A.S. 1996)

**Solution.** The given equation can be rewritten as

$$y \sin 2x \frac{dx}{dy} - \cos^2 x = 1 + y^2$$

or  $\sin 2x \frac{dx}{dy} - \frac{1}{y} \cos^2 x = \frac{1 + y^2}{y}$  (1)

putting  $-\cos^2 x = v$  or  $-2 \cos x (-\sin x) dx = dv$

or  $\sin 2x dx = dv$

the equation (1) reduce to

$$\frac{dv}{dy} + \frac{1}{y} v = \frac{1 + y^2}{y} \quad (2)$$

Which is linear in v with y as independent variable

Its integrating factor =  $e^{\int \frac{1}{y} dy} = e^{\log y} = y$

Multiplying both sides of (2) by y and integrating, we get

$$vy = C + \int (1 + y^2) dy, \text{ where } C \text{ is constant of integration}$$

or  $-y \cos^2 x = C + y + \frac{1}{3} y^3 \quad \therefore v = -\cos^2 x$

or  $y \cos^2 x + C + y + \frac{1}{3} y^3 = 0$

**Example 14.** Solve  $x (dy/dx) + y = y^2 \log x$

(U.P.C.S. 1995)

**Solution.** The given equation can be written as

$$\frac{1}{y^2} \frac{dy}{dx} + \frac{1}{x} \frac{1}{y} = \frac{1}{x} \log x \quad (1)$$

putting  $-\frac{1}{y} = v$  or  $\frac{1}{y^2} \frac{dy}{dx} = \frac{dv}{dx}$  in (1), we get

$$\frac{dv}{dx} - \frac{1}{x} v = \frac{1}{x} \log x \quad (2)$$

which is in the standand form of the linear equation and integrating factor

$$e^{-\int \frac{1}{x} dx} = e^{-\log x} = \frac{1}{x}$$

Multiplying both sides of (2) by the integrating factor and integrating, we get

$$v \cdot \frac{1}{x} = C + \int \frac{1}{x^2} \log x dx \text{ where } C \text{ is an arbitrary constant}$$

or  $\frac{v}{x} = C + \int t e^{-t} dt, \text{ putting } t = \log x$

Differential Equations of First Order and First Degree

$$\begin{aligned} \text{or} \quad \frac{v}{x} &= C + \left[ -t e^{-t} + \int e^{-t} dt \right] \\ &= C - (t+1) e^{-t} = C - (1 + \log x) e^{-\log x} \end{aligned}$$

$$\text{or} \quad -\frac{1}{xy} = C - (1 + \log x) \left( \frac{1}{x} \right) \quad \therefore v = -\frac{1}{y}$$

$$\text{or} \quad 1 = (1 + \log x) y - C xy$$

**Example 15.** Solve  $x (dy/dx) + y \log y = x ye^x$

(I.A.S. 2003, M.P.P.C.S. 1996)

**Solution.** Dividing both sides of the given equation by  $y$ , we get

$$\frac{x}{y} \frac{dy}{dx} + (\log y) = x e^x$$

$$\text{or} \quad \frac{1}{y} \frac{dy}{dx} + \frac{1}{x} (\log y) = e^x$$

putting  $v = \log y$

$$\text{or} \quad \frac{dv}{dx} = \frac{1}{y} \frac{dy}{dx}, \text{ the above equation reduce to}$$

$$\frac{dv}{dx} + \frac{1}{x} v = e^x \quad (1)$$

which is a linear equation in  $v$

$$\text{Its integrating factor} = e^{\int \frac{1}{x} dx} = e^{\log x} = x$$

Multiplying both sides of (1) by the integrating factor  $x$  and integrating, we have

$$v \cdot x = C + \int x e^x dx, \text{ where } C \text{ is an arbitrary constant}$$

$$\text{or} \quad v \cdot x = C + x e^x - \int 1 \cdot e^x dx$$

$$\text{or} \quad (\log y) x = C + x e^x - e^x$$

### Exact Differential Equations

A differential equation which can be obtained by direct differentiation of some function of  $x$  and  $y$  is called exact differential equation, consider the equation

$$Mdx + Ndy = 0 \text{ is exact}$$

$$\text{if } \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

where M and N are the functions of x and y.

Solution of a exact differential equation is

$$\int M dx + \int N dy = C$$

Regarding y as a constant                      only those terms of N not containing x

**Example 16.** Solve

$$\left\{ y \left( 1 + \frac{1}{x} \right) + \cos y \right\} dx + (x + \log x - x \sin y) dy = 0$$

(I.A.S. 1993)

**Solution.** Here  $M = y \left( 1 + \frac{1}{x} \right) \cos y$  and  $N = x + \log x - x \sin y$

$$\therefore \frac{\partial M}{\partial y} = \left( 1 + \frac{1}{x} \right) - \sin y \text{ and } \frac{\partial N}{\partial x} = 1 + \frac{1}{x} - \sin y$$

$$\Rightarrow \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}, \text{ hence the given equation is exact}$$

$$\text{Regarding } y \text{ as constant, } \int M dx = \int \left\{ y \left( 1 + \frac{1}{x} \right) + \cos y \right\} dx$$

$$= y \int \left( 1 + \frac{1}{x} \right) dx + \cos y \int dx$$

$$= y (x + \log x) + (\cos y) x \tag{1}$$

Also no new term is obtained by integrating N with respect to y.

$\therefore$  From (1), the required solution is

$$y (x + \log x) + x \cos x = C, \text{ where } C \text{ is an arbitrary constant}$$

**Example 17.** Verify that the equation

$$(x^4 - 2xy^2 + y^4) dx - (2x^2y - 4xy^3 + \sin y) dy = 0 \text{ is exact and solve it}$$

(U.P.P.C.S. 1996)

$$\text{Solution. Here } M = x^4 - 2xy^2 + y^4, \quad N = -2x^2y + 4xy^3 - \sin y$$

Differential Equations of First Order and First Degree

$$\therefore \frac{\partial M}{\partial y} = -4xy + 4y^3 \text{ and } \frac{\partial N}{\partial x} = -4xy + 4y^3$$

$$\therefore \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}, \text{ hence the given equation is exact}$$

Regarding y as constant,  $\int M dx = \int (x^4 - 2xy^2 + y^4) dx$

$$= \frac{x^5}{5} - x^2y^2 + y^4x \tag{1}$$

Integrating N with respect to y, we get

$$\int (-2x^2y + 4xy^3 - \sin y) dy = -x^2y^2 + xy^4 + \cos y$$

omitting from this the term  $-x^2y^2$  and  $xy^4$  which are already occurring in (1) we get  $\cos y$

$\therefore$  From (1) and (2) the required solution is

$$\frac{x^5}{5} - x^2y^2 + y^2x + \cos y = C$$

where C is an arbitrary constant

**Rules for finding integrating factor**

A differential equation of the type  $Mdx + Ndy = 0$  which is not exact can be made exact by multiplying the equation by some function of x and y, which is called an integrating factor.

A few methods (without proof) are given below for finding the integrating factor in certain cases.

**Method I. Integrating factor found by inspection.**

In the case of some differential equation the integrating factor can be found by inspection. A few exact differentials are given below which would help students (if they commit these to memory) in finding the integrating factors.

$$(a) \quad d\left(\frac{y}{x}\right) = \frac{x dy - y dx}{x^2}$$

$$(b) \quad d\left(\frac{x}{y}\right) = \frac{y dx - x dy}{y^2}$$

$$(c) \quad d(xy) = x dy + y dx$$

$$(d) \quad d\left(\frac{x^2}{y}\right) = \frac{2yx dx - x^2 dy}{y^2}$$



$$(e) \quad d\left(\frac{y^2}{x}\right) = \frac{2xy \, dy - y^2 \, dx}{x^2} \qquad (f) \quad d\left(\frac{y^2}{x^2}\right) = \frac{2x^2y \, dy - 2xy^2 \, dx}{x^4}$$

$$(g) \quad d\left(\tan^{-1} \frac{x}{y}\right) = \frac{y \, dx - x \, dy}{x^2 + y^2} \qquad (h) \quad d\left(\tan^{-1} \frac{y}{x}\right) = \frac{x \, dy - y \, dx}{x^2 + y^2}$$

$$(i) \quad d\left(\frac{1}{2} \log(x^2 + y^2)\right) = \frac{x \, dx + y \, dy}{x^2 + y^2} \qquad (j) \quad d\left(-\frac{1}{xy}\right) = \frac{x \, dy + y \, dx}{x^2y^2}$$

$$(k) \quad d \log\left(\frac{x}{y}\right) = \frac{y \, dx - x \, dy}{xy} \qquad (l) \quad d \log\left(\frac{y}{x}\right) = \frac{x \, dy - y \, dx}{xy}$$

$$(m) \quad d\left(\frac{e^x}{y}\right) = \frac{ye^x \, dx - e^x \, dy}{y^2}$$

**Example 18.** Solve  $(1 + xy) y \, dx + (1 - xy) x \, dy = 0$

(I.A.S. 1992, M.P.P.C.S. 1974)

**Solution.** The given equation can be written as

$$(y \, dx + x \, dy) + (xy^2 \, dx - x^2y \, dy) = 0$$

or 
$$d(yx) + xy^2 \, dx - x^2y \, dy = 0$$

Dividing both sides of this equation by  $x^2y^2$ , we get

$$\frac{d(yx)}{x^2y^2} + \frac{1}{x} \, dx - \frac{1}{y} \, dy = 0$$

or 
$$\frac{1}{z^2} \, dz + \frac{1}{x} \, dx - \frac{1}{y} \, dy = 0, \quad \text{where } z = xy$$

Integrating,  $-\frac{1}{z} + \log x - \log y = C$ , where  $C$  is an arbitrary constant

or 
$$-\frac{1}{xy} + \log x - \log y = C, \quad \text{putting } z = xy$$

or 
$$\log(x/y) = C + \frac{1}{xy}$$

**Example 19.** Solve  $(xy^2 + 2x^2y^3) \, dx + (x^2y - x^3y^2) \, dy = 0$

(U.P.P.C.S. 1993)

**Solution.** The given equation can be rewritten as

Differential Equations of First Order and First Degree

$$xy^2 (1 + 2xy) dx + x^2y (1 - xy) dy = 0$$

or  $y (1 + 2xy) dx + x (1 - xy) dy = 0$

or  $(y dx + x dy) + 2xy^2 dx - x^2y dy = 0$

or  $d(xy) + 2xy^2 dx - x^2y dy = 0$

Dividing both sides of this equation by  $x^2y^2$ , we get

$$\frac{d(xy)}{x^2 y^2} + \frac{2}{x} dx - \frac{1}{y} dy = 0$$

or  $\left(\frac{1}{z^2}\right) dz + (z/x) dx - (1/y) dy = 0$ , where  $z = xy$

Integrating,  $-\frac{1}{z} + 2 \log x - \log y = C$ , where  $C$  is an arbitrary constant

or  $-\left(\frac{1}{xy}\right) + \log\left(\frac{x^2}{y}\right) = C$ , putting  $z = xy$

or  $\log\left(\frac{x^2}{y}\right) = C + \left(\frac{1}{xy}\right)$

**Example 20.** Solve  $y \sin 2x dx = (1 + y^2 + \cos^2 x) dy$

(I.A.S. 1996)

**Solution.** The given equation can be written as

$$2y \sin x \cos x dx - \cos^2 x dy = (1 + y^2) dy$$

or  $-d[y \cos^2 x] = (1 + y^2) dy$

Integrating,  $-y \cos^2 x = y + \frac{y^3}{3} + C$ , where  $C$  is an arbitrary constant.

**Method II.** In the differential equation  $Mdx + Ndy = 0$ , If  $M = y f_1(xy)$  and  $N = x f_2(xy)$ , then  $\frac{1}{Mx - Ny}$  is an integrating factor, Provided  $Mx - Ny \neq 0$

**Example 21.** Solve

$$(xy \sin xy + \cos xy) y dx + (xy \sin xy - \cos xy) x dy = 0$$

(Bihar P.C.S. 2007, U.P.P.C.S. 1990, 94)

**Solution.** Integrating factor

$$= \frac{1}{(xy \sin xy + \cos xy) xy - (xy \sin xy - \cos xy) xy}$$
$$= \frac{1}{2yx \cos xy}$$

Multiplying both sides of the given equation by this integrating factor, we get

$$\frac{1}{2} \left( \tan xy + \frac{1}{xy} \right) y dx + \frac{1}{2} \left( \tan xy - \frac{1}{xy} \right) x dy = 0$$

or  $(\tan xy) (y dx + x dy) + \frac{1}{x} dx - \frac{1}{y} dy = 0$

or  $(\tan xy) d(xy) + \frac{1}{x} dx - \frac{1}{y} dy = 0$

or  $\tan z dz + \frac{1}{x} dx - \frac{1}{y} dy = 0$ , where  $z = xy$

Integrating term by term, we get

$\log (\sec z) + \log x - \log y = \log C$ , where  $C$  is an arbitrary constant

or  $\log \left\{ \frac{x \sec z}{y} \right\} = \log C$

or  $\frac{x}{y} (\sec z) = C$

or  $x \sec (xy) = Cy$

**Method III.** In the differential equation  $Mdx + Ndy = 0$  if  $\frac{1}{N} \left( \frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right)$  is a

function of  $x$  alone, say  $f(x)$ , then the integrating factor is  $e^{\int f(x) dx}$

**Example 22.** solve  $(x^2 + y^2 + 1) dx - 2xy dy = 0$

(U.P.P.C.S. 1988, 82)

**Solution.** Here  $M = x^2 + y^2 + 1$  and  $N = -2xy$

$$\therefore \frac{\partial M}{\partial y} = 2y \quad \text{and} \quad \frac{\partial M}{\partial x} = -2y$$

Differential Equations of First Order and First Degree

Therefore,  $\frac{1}{N} \left( \frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) = \frac{1}{-2xy} (2y + 2y) = -\frac{2}{x}$ , which is a function of  $x$  alone. Hence method III is applicable

Here  $f(x) = -2/x$

$$\begin{aligned} \therefore \text{Integrating factor} &= e^{\int f(x) dx} = e^{-\int \frac{2}{x} dx} \\ &= e^{-2 \log x} = 1/x^2 \end{aligned}$$

Multiplying the given equation by this integrating factor  $1/x^2$ , we get

$$\frac{1}{x^2} (x^2 + y^2 + 1) dx - \frac{1}{x^2} (2xy) dy = 0$$

or 
$$\left[ 1 + \frac{y^2}{x^2} + \frac{1}{x^2} \right] dx - 2 \frac{y}{x} dy = 0$$

or 
$$\left[ 1 + \frac{1}{x^2} \right] dx + \left[ \frac{y^2}{x^2} dx - 2 \frac{y}{x} dy \right]$$
 which is an exact

or 
$$\left[ 1 + \frac{1}{x^2} \right] dx + d \left( -\frac{y^2}{x} \right) = 0$$
 using method I

Integrating term by term, we get

$$x - \frac{1}{x} + \left( -\frac{y^2}{x} \right) = C, \text{ where } C \text{ is constant of integration}$$

or  $x^2 - 1 - y^2 = Cx$  is the required solution.

**Method IV.** In the equation  $Mdx + Ndy = 0$

If  $\frac{1}{M} \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right)$  is a function of  $y$  alone, say  $f(y)$ ,

then the integrating factor is  $e^{\int f(y) dy}$

**Example 23.** Solve  $(xy^3 + y) dx + 2(x^2y^2 + x + y^4) dy = 0$

**Solution.** Here  $M = xy^3 + y$  and  $N = 2x^2y^2 + 2x + 2y^4$

$$\therefore \frac{\partial M}{\partial y} = 2xy^2 + 1, \text{ and } \frac{\partial N}{\partial x} = 4xy^2 + 2$$

$$\begin{aligned}\therefore \frac{1}{M} \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) &= \frac{1}{(xy^3 + y)} \left\{ (4xy^2 + 2) - (3xy^2 + 1) \right\} \\ &= \frac{1}{y(xy^2 + 1)} (xy^2 + 1) = \frac{1}{y}, \text{ which is a function of } y \text{ alone}\end{aligned}$$

and equal to  $f(y)$  say.

$$\begin{aligned}\text{Then integrating factor} &= e^{\int f(y) dy} = e^{\int \frac{1}{y} dy} \\ &= e^{\log y} = y\end{aligned}$$

Multiplying the given equation by integrating factor  $y$  we get

$$(xy^4 + y^2) dx + (2x^2 y^3 + 2xy + 2y^5) dy = 0$$

which is an exact differential equation and solving by the method of exact, we have

$$3x^2y^4 + 6xy^2 + 2y^6 = 6 C \text{ is the required solution.}$$

**Example 24.** Solve  $(3x^2 y^4 + 2xy) dx + (2x^3 y^3 - x^2) dy = 0$

(U.P.P.C.S. 2001)

**Solution.** Here  $M = 3x^2 y^4 + 2xy$ ,  $N = 2x^3 y^3 - x^2$

$$\text{and } \frac{\partial M}{\partial y} = 12x^2 y^3 + 2x, \quad \frac{\partial N}{\partial x} = 6x^2 y^3 - 2x$$

As  $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$ , so the equation is not exact in this form. Thus, we have to find the integrating factor by trial. In the present case, we see that

$$\frac{1}{M} \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) = \frac{6x^2 y^3 - 2x - 12x^2 y^3 - 2x}{3x^2 y^4 + 2xy} = -\frac{2}{y}, \text{ is the function of } y \text{ alone}$$

$$\text{The integrating factor is } e^{-\int \frac{2}{y} dy} = \frac{1}{y^2}$$

Thus, the differential equation becomes

$$(3x^2 y^2 + 2x/y) dx + (2x^3 y - x^2/y^2) dy = 0$$

$$\text{Which is an exact, as } \frac{\partial M}{\partial y} = 6x^2 y - 2x/y^2 = \frac{\partial N}{\partial x}$$

Differential Equations of First Order and First Degree

Its solution is

$$\int (3x^2 y^2 + 2x/y) dx = 0$$

or  $x^3 y^2 + x^2/y = C$

Where C is an arbitrary constant.

**Method V.** If the equation  $Mdx + N dy = 0$  is homogeneous then  $\frac{1}{Mx + Ny}$  is an

integrating factor, Provided  $Mx + Ny \neq 0$

**Example 25.** Solve  $x^2y dx - (x^3 + y^3) dy = 0$

**Solution.** Here  $M = x^2y$  and  $N = -x^3 - y^3$

$$\therefore Mx + Ny = x^3y - x^3y - y^4 = -y^4 \neq 0$$

$$\therefore \text{Integrating factor} = \frac{1}{Mx + Ny} = -\frac{1}{y^4} \neq 0$$

Multiplying the given equation by this integrating factor  $-1/y^4$ , we get

$$-\frac{x^2}{y^3} dx + \left( \frac{x^3}{y^4} + \frac{1}{y} \right) dy = 0$$

In this form of the equation,  $M = \frac{-x^2}{y^3}$  and  $N = \frac{x^3}{y^4} + \frac{1}{y}$

$$\therefore \frac{\partial M}{\partial y} = \frac{3x^2}{y^4} \text{ and } \frac{\partial N}{\partial x} = \frac{3x^2}{y^4}$$

$$\Rightarrow \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Hence this equation is an exact

Solving, we get

$$x^3 = 3y^3 (\log y - C)$$

**Method VI.** If the equation be of the form  $x^a y^b (my dx + nx dy) + x^c y^d (py dx + qx dy) = 0$

where a, b, c, d, m, n, p and q are constants, then the integrating factor is  $x^h y^k$ , where h and k can be obtained by applying the condition that after multiplication by  $x^h y^k$  the equation is exact.

**Example 26.** Solve  $(y^2 + 2x^2y) dx + (2x^3 - xy) dy = 0$

or  $y(y dx - x dy) + 2x^2(y dx + x dy) = 0$  (U.P.P.C.S. 2000)

**Solution.** The given equation can be rewritten as

$$y(y + 2x^2) dx + x(2x^2 - y) dy = 0$$

which is of the form as given in method VI

Let  $x^h y^k$  be an integrating factor

Multiplying the given equation by  $x^h y^k$ , we get  $(x^h y^{k+2} + 2x^{h+2} y^{k+1}) dx + (2x^{h+3} y^k - x^{h+1} y^{k+1}) dy = 0$

Here  $M = x^h y^{k+2} + 2x^{h+2} y^{k+1}$  and  $N = 2x^{h+3} y^k - x^{h+1} y^{k+1}$

$$\therefore \frac{\partial M}{\partial y} = (k+2)x^h y^{k+1} + 2(k+1)x^{h+2} y^k \quad (1)$$

and 
$$\frac{\partial N}{\partial x} = 2(h+3)x^{h+2} y^k - (h+1)x^h y^{k+1} \quad (2)$$

If the equation (A) be exact we must have  $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$

Or  $(k+2)x^h y^{k+1} + 2(k+1)x^{h+2} y^k = -(h+1)x^h y^{k+1} + 2(h+3)x^{h+2} y^k$  from (1) and (2)

Equating coefficients of  $x^h y^{k+1}$  and  $x^{h+2} y^k$  on both sides we get  $k+2 = -(h+1)$  and  $2(k+1) = 2(h+3)$

hence solving we get

$$h = -\frac{5}{2}, k = -\frac{1}{2}$$

$\therefore$  The integrating factor =  $x^h y^k = x^{-5/2} y^{-1/2}$

Multiplying the given by  $x^{-5/2} y^{-1/2}$ , we get

$$(x^{-5/2} y^{3/2} + 2x^{-1/2} y^{1/2}) dx + (2x^{1/2} y^{-1/2} - x^{-3/2} y^{1/2}) dy = 0$$

In this form, we have  $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$ , the equation is exact

$\therefore$  Regarding  $y$  as constant

$$\int M dx = \int (x^{-5/2} y^{3/2} + 2x^{-1/2} y^{1/2}) dx$$

Differential Equations of First Order and First Degree

$$= \frac{2}{3} x^{-3/2} y^{3/2} + 4x^{1/2} y^{1/2}$$

Also no new term is obtained by integrating N with respect to y, Hence the required solution is

$$- \frac{2}{3} x^{-3/2} y^{3/2} + 4x^{1/2} y^{1/2} = C$$

Where C is an arbitrary constant

### EXERCISE

Solve the following differential equations

1.  $(x^2 - 2x + 2y^2) dx + 2xy (1 + \log x^2) dy = 0$

(I.A.S. 1991)

Ans.  $x^2 - 4x + 4y^2 \log x + 2y^2 = C$

2.  $x^2 \frac{dy}{dx} = x^2 + xy + y^2$

(U.P.P.C.S. 1998)

Ans.  $x = C e^{\tan^{-1}\left(\frac{y}{x}\right)}$

3.  $(3y + 2x + 4) dx - (4x + 6y + 5) dy = 0$

(U.P.P.C.S. 2001)

Ans.  $14(2x + 3y) - 9 \log(14x + 21y + 22) = 49x + C$

4.  $xy - \frac{dy}{dx} = y^3 e^{-x^2}$

(I.A.S. 1998)

Ans.  $e^x = y^2(x + C)$

5. Solve the initial value problem  $\frac{dy}{dx} = \frac{x}{x^2y + y^3}$ ,  $y(0) = 0$

(I.A.S. 1997)

Ans.  $(x^2 + y^2 + 2) = 2e^{y^2/2}$

6. Show that the equation  $(4x + 3y + 1) dx + (3x + 2y + 1) dy = 0$  represents a family of hyperbolas having as asymptotes the line  $x + y = 0$ ,  $2x + y + 1 = 0$

(I.A.S. 1998)



7. The equations of motion of a particle are given by  $\frac{dx}{dt} + wy = 0$ ,  
 $\frac{dy}{dt} - wx = 0$ , Find the path of the particle and so that it is a circle.

(U.P.T.U. 2009)

Hint.  $y(t) = C_1 \cos wt + C_2 \sin wt$

$$x(t) = C_2 \cos wt - C_1 \sin wt$$

$$\text{so } x^2 + y^2 = C_1^2 + C_2^2 = R^2$$

### Objective Type of Questions

Choose a correct answer from the four answers given in each of the following questions:

1. The differential equation  $y dx + x dy = 0$  represents a family of

(U.P.P.C.S. 1999)

- (a) Circles (b) Ellipses  
(c) Cycloids (d) Rectangular hyperbolas

Ans. (d)

2. The solution of  $(xy^2 + 1) dx + (x^2y + 1) dy = 0$  is

(U.P.P.C.S. 1999)

- (a)  $x^2y^2 + 2x^2 + 2y^2 = C$  (b)  $x^2y^2 + x^2 + y^2 = C$   
(c)  $x^2y^2 + x + y = C$  (d)  $x^2y^2 + 2x + 2y = C$

Ans. (d)

3. The differential equation for the family of all tangents to the parabola  $y^2 = 2x$

(U.P.P.C.S. 1999)

- (a)  $2x (y')^2 + 1 = 2yy'$  (b)  $2xy + 1 = 2yy'$   
(c)  $2x^2y' + 1 = 2yy'$  (d)  $2 (y')^2 + x = 2yy'$

Ans. (a)

4. The general solution of the differential equation  $(1 + x) y dx + (1 - y) x dy = 0$  is

(U.P.P.C.S. 2000)

- (a)  $xy = C e^{x-y}$  (b)  $x + y = C e^{xy}$   
(c)  $xy = C e^{y-x}$  (d)  $x - y = C e^{xy}$

Ans. (c)

Differential Equations of First Order and First Degree

5. An integrating factor of the differential equation  $(1 + x^2) \frac{dy}{dx} + 2xy = \cos x$  is

(U.P.P.C.S. 2000)

- |                     |                         |
|---------------------|-------------------------|
| (a) $1 + x^2$       | (b) $\frac{1}{1 + x^2}$ |
| (c) $\log(1 + x^2)$ | (d) $-\log(1 + x^2)$    |

Ans. (a)

6. If  $\frac{dy}{dx} = e^{-2y}$ ,  $y = 0$  when  $x = 5$ , then the value of  $x$  for  $y = 3$  is

(U.P.P.C.S. 2000)

- |                   |               |
|-------------------|---------------|
| (a) $(e^6 + 9)/2$ | (b) $e^5$     |
| (c) $\log_e 6$    | (d) $e^6 + 1$ |

Ans. (a)

7. The solution of  $(x - 1) dy = y dx$ ,  $y(0) = -5$  is

- |                    |                     |
|--------------------|---------------------|
| (a) $y = 5(x - 1)$ | (b) $y = -5(x - 1)$ |
| (c) $y = 5(x + 1)$ | (d) $y = -5(x + 1)$ |

Ans. (a)

8. The differential equation  $x dx - y dy = 0$  represents a family of

(U.P.P.C.S. 2001)

- |                            |              |
|----------------------------|--------------|
| (a) Circles                | (b) Ellipse  |
| (c) rectangular hyperbolas | (d) Cycloids |

Ans. (c)

9. The solution of the differential equation  $(x + 2y) dy - (2x - y) dx = 0$  is

(U.P.P.C.S. 2001)

- |                           |                          |
|---------------------------|--------------------------|
| (a) $x^2 + y^2 - 2xy = C$ | (b) $xy + y^2 + x^2 = C$ |
| (c) $x^2 + 4xy + y^2 = C$ | (d) $xy + y^2 - x^2 = C$ |

Ans. (d)

10. In the differential equation  $\frac{x dy}{dx} + my = e^{-x}$ , if the integrating factor is  $\frac{1}{x^2}$ ,

then the value of  $m$  is

- |       |        |
|-------|--------|
| (a) 2 | (b) -2 |
| (c) 1 | (d) -1 |

Ans. (b)

11. The solution of the variable separable equation  $(x^2 + 1)(y^2 - 1) dx + xy dy = 0$  is

(I.A.S. 1988)

- (a)  $y^2 - 1 = x^2 + 1 + C$                       (b)  $\log(y^2 - 1) = C \log(x^2 + 1)$   
(c)  $y^2 = 1 + C \frac{e^{-x^2}}{x^2}$                                   (d) none of these

Ans. (c)

12. The solution of the differential equation  $\frac{dy}{dx} = e^{x-y} + x^2 e^{-y}$  is

(M.P.P.C.S. 1991, R.A.S. 1993, U.P.P.C.S. 1995)

- (a)  $e^y = e^x + \frac{1}{3} x^3 + C$                       (b)  $e^y = e^{-x} + \frac{1}{3} x^3 + C$   
(c)  $e^y = e^x + x^3 + C$                       (d)  $e^{-y} = \frac{1}{3} x^3 + e^x + C$

Ans. (a)

13. If  $\frac{1}{M} \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) = f(y)$  a function of  $y$  alone, then the integrating factor of  $Mdx + Ndy = 0$  is

(M.P.P.C.S. 1991, 93)

- (a)  $e^{-\int f(y) dy}$                                   (b)  $e^{\int f(y) dy}$   
(c)  $f(y) \int e^{f(y)} dy$                       (d)  $\int e^{f(y)} f(y) dy$

Ans. (b)

14. The solution of the differential equation  $(x + y)^2 \frac{dy}{dx} = a^2$  is given by

(I.A.S. 1994)

- (a)  $y + x = a \tan \left( \frac{y - c}{a} \right)$                       (b)  $(y - x) = a \tan (y - c)$   
(c)  $y - x = \tan \left( \frac{y - c}{a} \right)$                       (d)  $a(y - x) = \tan \left( \frac{y - c}{a} \right)$

Ans. (a)

Differential Equations of First Order and First Degree

15.  $Pdx + x \sin y \, dy = 0$  is exact, then P can be

(MP.P.P.C.S. 1994)

(a)  $\sin y + \cos y$                       (b)  $-\sin y$

(c)  $x^2 - \cos y$                       (d)  $\cos y$

Ans. (c)

16. The solution of the differential equation  $(x - y^2) \, dx + 2xy \, dy = 0$  is

(I.A.S. 1993)

(a)  $ye^{y^2/x} = A$                       (b)  $xe^{y^2/x} = A$

(c)  $ye^{x/y^2} = A$                       (d)  $xe^{x/y^2} = A$

17. The solution of the equation  $\frac{dy}{dx} + 2xy = 2xy^2$  is

(I.A.S. 1994)

(a)  $y = \frac{cx}{1 + e^{x^2}}$                       (b)  $y = \frac{1}{1 - ce^x}$

(c)  $y = \frac{1}{1 + ce^{x^2}}$                       (d)  $y = \frac{cx}{1 + e^{x^2}}$

Ans. (c)

18. The homogeneous differential equation

$$M(x, y) \, dx + N(x, y) \, dy = 0$$

can be reduced to a differential equation in which the variables are separable, by the substitution-

(I.A.S. 1996)

(a)  $y = vx$                               (b)  $xy = v$

(c)  $x + y = v$                         (d)  $x - y = v$

Ans. (a)

19. The solution of the differential equation

$$\frac{dy}{dx} + \frac{y}{x} = x^2$$

Under the condition that  $y = 1$ , when  $x = 1$  is

(I.A.S. 1996)

(a)  $4xy = x^3 + 3$                       (b)  $4xy = y^4 + 3$

(c)  $4xy = x^4 + 3$

(d)  $4xy = y^3 + 3$

Ans. (c)

20. The necessary condition to exact the differential equation  $Mdx + Ndy = 0$  will be

(MP.P.C.S. 1993, R.A.S. 1995)

(a)  $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$

(b)  $\frac{\partial M}{\partial x} = \frac{\partial N}{\partial y}$

(c)  $\frac{\partial^2 M}{\partial x^2} = \frac{\partial^2 N}{\partial y^2}$

(d)  $\frac{\partial^2 M}{\partial y^2} = \frac{\partial^2 N}{\partial x^2}$

21. If  $I_1, I_2$  are integrating factors of the equations  $xy' + 2y = 1$  and  $xy' - 2y = 1$  then

(M.P.P.C.S. 1994)

(a)  $I_1 = -I_2$

(b)  $I_1 I_2 = x$

(c)  $I_1 = x^2 I_2$

(d)  $I_1 I_2 = 1$

Ans. (d)

22. The family of conic represented by the solution of the differential equation  $(4x + 3y + 1) dx + (3x + 2y + 1) dy = 0$  is

(U.P.P.C.S. 1994)

(a) Circles

(b) Parabolas

(c) hyperbolas

(d) ellipse

Ans. (c)

# Chapter 3

## Linear Differential Equations with Constant Coefficients and Applications

---

### INTRODUCTION

A differential equation is of the form

$$\frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + a_n y = Q \quad (1)$$

where  $a_1, a_2, \dots, a_n$  are constants and  $Q$  is a function of  $x$  only, is called a linear differential equation of  $n^{\text{th}}$  order. Such equations are most important in the study of electro-mechanical vibrations and other engineering problems.

The operator  $\frac{d}{dx}$  is denoted by  $D$ .

$$\therefore D^n y + a_1 D^{n-1} y + \dots + a_n y = Q$$

or  $f(D) y = Q$

where  $f(D) = D^n + a_1 D^{n-1} + \dots + a_n$

### Solution of the Differential Equation

If the given equation is

$$\frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + a_n y = 0 \quad (1)$$

or  $(D^n + a_1 D^{n-1} + a_2 D^{n-2} + \dots + a_n) y = 0 \quad (2)$

Let  $y = e^{mx}$

$$\Rightarrow D^r y = m^r e^{mx}, \quad 1 \leq r \leq n$$

$\therefore$  Then from equation (2)

$$m^n + a_1 m^{n-1} + a_2 m^{n-2} + \dots + a_n e^{mx} = 0$$

$y = e^{mx}$  is a solution of (1), if

$$m^n + a_1 m^{n-1} + a_2 m^{n-2} + \dots + a_n = 0$$

This equation is called the auxiliary equation.

**Case 1. When Auxiliary Equation has Distinct and real Roots**

Let  $m_1, m_2, \dots, m_n$  are distinct roots of the auxiliary equation, then the general solution of (1) is  $y = C_1 e^{m_1 x} + C_2 e^{m_2 x} + \dots + C_n e^{m_n x}$

where  $C_1, C_2, \dots, C_n$  are arbitrary constants

**Illustration.** Solve the differential equation

$$\frac{d^2 y}{dx^2} + 3 \frac{dy}{dx} - 54 y = 0$$

**Solution.** The given equation is

$$(D^2 + 3D - 54) y = 0$$

Here auxiliary equation is

$$m^2 + 3m - 54 = 0$$

or  $(m + 9)(m - 6) = 0$

$\Rightarrow m = 6, -9$

Hence the general solution of the given differential equation is  $y = C_1 e^{6x} + C_2 e^{-9x}$

**Case II. When Auxiliary Equation has real and some equal roots.**

If the auxiliary equation has two roots equal, say  $m_1 = m_2$  and others are distinct say  $m_3, m_4, \dots, m_n$ . In this Case the general solution of the equation (1) is

$$y = (C_1 + C_2 x) e^{m_1 x} + C_3 e^{m_3 x} + \dots + C_n e^{m_n x}$$

where  $C_1, C_2, C_3, \dots, C_n$  are arbitrary constants

**Illustration.** Solve the differential equation

$$(D^4 - D^3 - 9D^2 - 11D - 4) y = 0$$

**Solution.** The auxiliary equation of the give equation is

$$m^4 - m^3 - 9m^2 - 11m - 4 = 0$$

or  $(m + 1)^3 (m - 4) = 0$

$\Rightarrow m = -1, -1, -1, 4$

Hence, the required solution is

$$y = (C_1 + C_2 x + C_3 x^2) e^{-x} + C_4 e^{4x}$$

or 
$$y = (C_1 x^2 + C_2 x + C_3) e^{-x} + C_4 e^{4x}$$

**Case III. When the auxiliary equation has imaginary roots**

If there are one pair of imaginary roots say  $m_1 = \alpha + i\beta$ ,  $m_2 = \alpha - i\beta$  i.e.  $\alpha \pm i\beta$  say then the required solution is

$e^{x \text{ (Real part)}} [C_1 \{\cos (\text{imaginary part}) x\} + C_2 \{\sin (\text{imaginary part}) x\}]$

i.e. 
$$e^{\alpha x} [C_1 \cos \beta x + C_2 \sin \beta x]$$

or 
$$y = C_1 e^{\alpha x} \cos (\beta x + C_2)$$

**Illustration.** Solve  $(D^2 - 2D + 5) y = 0$

**Solution.** Here the auxiliary equation is

$$m^2 - 2m + 5 = 0$$

or 
$$m = \frac{1}{2} [2 \pm \sqrt{(4 - 20)}] = 1 \pm 2i$$

$\therefore$  The required solution is

$$y = e^x (C_1 \cos 2x + C_2 \sin 2x)$$

**Particular Integral (P.I.)**

when the equation is

$$D^n + a_1 D^{n-1} + \dots + a_n y = Q$$

or 
$$f(D) y = Q$$

The general solution of  $f(D) y = Q$  is equal to the sum of the general solution of  $f(D) y = 0$  called complementary function (C.F.) and any particular integral of the equation  $f(D) y = Q$

$\therefore$  General solution = C.F. + P.I.

A particular integral of the differential equation

$$f(D) y = Q \text{ is given by } \frac{1}{f(D)} Q$$

**Methods of finding Particular integral**

**(A)**

**Case I.** P.I., when  $Q$  is of the form of  $e^{ax}$ , where  $a$  is any constant and  $f(a) \neq 0$

we know that  $D(e^{ax}) = a e^{ax}$



$$D^2 (e^{ax}) = a^2 e^{ax}$$

$$D^3 (e^{ax}) = a^3 e^{ax}$$

In general  $D^n (e^{ax}) = a^n e^{ax}$

$$\therefore f(D) (e^{ax}) = f(a) e^{ax}$$

or  $\frac{1}{f(D)} f(D) e^{ax} = \frac{1}{f(D)} f(a) e^{ax}$

or  $e^{ax} = f(a) \frac{1}{f(D)} e^{ax} \because f(a) \text{ is constant}$

or  $\frac{1}{f(a)} e^{ax} = \frac{1}{f(D)} e^{ax}$

Hence P.I. =  $\frac{1}{f(D)} e^{ax} = \frac{1}{f(a)} e^{ax}$ , if  $f(a) \neq 0$

**Case II.** P.I., when Q is of the form of  $e^{ax}$ , and  $f(a) = 0$

Then  $\frac{1}{f(D)} (e^{ax}) = e^{ax} \frac{1}{f(D+a)} 1$ , which shows that if  $e^{ax}$  is brought to the left from the right of  $\frac{1}{f(D)}$ , then D should be replaced by  $(D + a)$

### Another method for Exceptional Case

If  $f(a) = 0$ , then

$$\begin{aligned} \text{P.I.} &= \frac{1}{f(D)} e^{ax} = x \frac{1}{f'(D)} e^{ax} \\ &= x \frac{e^{ax}}{f'(a)}, \text{ if } f'(a) \neq 0 \end{aligned}$$

If  $f'(a) = 0$ , then

$$\begin{aligned} \text{P.I.} &= \frac{1}{f(D)} e^{ax} \\ &= x^2 \frac{e^{ax}}{f''(a)}, \text{ if } f''(a) \neq 0 \end{aligned}$$

**Example 1.** Solve  $(D^2 - 2D + 5) y = e^{-x}$

Linear Differential Equations with Constant Coefficients and Applications

**Solution.** Here auxiliary equation is  $m^2 - 2m + 5 = 0$ , whose roots are

$$m = -1 \pm 2i$$

$\therefore$  C.F. =  $e^{-x} [C_1 \cos 2x + C_2 \sin 2x]$ , where  $C_1$  and  $C_2$  are arbitrary constants

and

$$\begin{aligned} \text{P.I.} &= \frac{1}{D^2 - 2D + 5} e^{-x} \\ &= \frac{1}{(-1)^2 - 2(-1) + 5} e^{-x} \quad \because \text{here } a = -1 \\ &= \frac{1}{8} e^{-x} \end{aligned}$$

$\therefore$  The required solution is  $y = \text{C.F.} + \text{P.I.}$

$$\text{i.e. } y = e^{-x} (C_1 \cos 2x + C_2 \sin 2x) + \frac{1}{8} e^{-x}$$

**Example 2.** Solve  $(D - 1)^2 (D^2 + 1)^2 y = e^x$

**Solution.** Here the auxiliary equation is

$$(m - 1)^2 (m^2 + 1)^2 = 0$$

or

$$m = 1, 1, \pm i, \pm i$$

$$\therefore \text{C.F.} = (C_1 + C_2 x) e^x + (C_3 x + C_4) \cos x + (C_5 x + C_6) \sin x$$

where c's are arbitrary constant

and

$$\begin{aligned} \text{P.I.} &= \frac{1}{(D - 1)^2 (D^2 + 1)^2} e^x \\ &= \frac{1}{(D - 1)^2} \frac{1}{(1^2 + 1)^2} e^x \\ &= \frac{1}{(D - 1)^2} \frac{1}{(2)^2} e^x \\ &= \frac{1}{(D - 1)^2} \frac{1}{4} e^x \\ &= e^x \frac{1}{(D + 1 - 1)^2} \frac{1}{4} \end{aligned}$$

$$= e^x \frac{1}{D^2} \frac{1}{4} = \frac{1}{4} e^x \frac{1}{D^2} (1) = \frac{1}{4} e^x \frac{x^2}{2} = \frac{1}{8} x^2 e^x$$

∴ The required solution is  $y = C. F + P.I.$

or 
$$y = (C_1 x + C_2) e^x + (C_3 x + C_4) \cos x + (C_5 x + C_6) \sin x + \frac{1}{8} x^2 e^x$$

**Example 3.** Solve  $(D + 2) (D - 1)^3 y = e^x$

**Solution.** Here the auxiliary equation is

$$(m + 2) (m - 1)^3 = 0$$

or  $m = -2$  and  $m = 1$  (thrice)

Therefore C. F =  $C_1 e^{-2x} + (C_2 x^2 + C_3 x + C_4) e^x$ , where  $C_1, C_2$  and  $C_3$  are constants and

$$\begin{aligned} P.I. &= \frac{1}{(D + 2) (D - 1)^3} e^x \\ &= \frac{1}{(1 + 2) (D - 1)^3} e^x \\ &= \frac{1}{3} \frac{1}{(D - 1)^3} e^x = \frac{1}{3} e^x \frac{1}{\{(D + 1) - 1\}^3} 1 \\ &= \frac{1}{3} e^x \frac{1}{D^3} 1 \\ &= \frac{1}{3} e^x \left( \frac{1}{6} x^3 \right) = \frac{1}{18} x^3 e^x \end{aligned}$$

∴ the required solution is  $y = C.F + P.I$

or 
$$y = C_1 e^{-2x} + (C_2 x^2 + C_3 x + C_4) e^x + \frac{1}{18} x^3 e^x$$

**(B) (i) P.I. when Q is of the form  $\sin ax$  or  $\cos ax$  and  $f(-a^2) \neq 0$**

$$\frac{1}{f(D^2)} \sin ax = \frac{1}{f(-a^2)} \sin ax, \text{ if } f(-a^2) \neq 0$$

$$\& \frac{1}{f(D^2)} \cos ax = \frac{1}{f(-a^2)} \cos ax, \text{ if } f(-a^2) \neq 0$$

**(ii) P.I. when Q is of the form  $\sin ax$  or  $\cos ax$  and  $f(-a^2) = 0$**

Linear Differential Equations with Constant Coefficients and Applications

Let  $f(D) = D^2 + a^2$  and  $Q = \sin ax$

Then

$$\begin{aligned} \text{P.I.} &= \frac{1}{D^2 + a^2} \sin ax \\ &= \frac{1}{D^2 + a^2} (\text{Imaginary part of } e^{iax}) \\ &= \text{Imaginary part of } \frac{1}{D^2 + a^2} e^{iax} \\ &= \text{I.P. of } e^{iax} \frac{1}{\{(D + ia)^2 + a^2\}} 1 \\ &= \text{I.P. of } e^{iax} \frac{1}{(D^2 + 2iaD + i^2a^2 + a^2)} 1 \\ &= \text{I.P. of } e^{iax} \frac{1}{2iaD \left[ 1 + \frac{D}{2ia} \right]} 1 \\ &= \text{I.P. of } \frac{e^{iax}}{2ia} \frac{1}{D} \left( 1 - \frac{D}{2ia} + \dots \right) 1 \\ &= \text{I.P. of } \frac{e^{iax}}{2ia} \frac{1}{D} 1 = \text{I.P. of } \frac{e^{iax}}{2ia} x \\ &= \text{I.P. of } \frac{e^{iax} xi}{2i^2 a} \\ &= \text{I.P. of } \frac{1}{2} \left( -\frac{x}{a} \right) i (\cos ax + i \sin ax) \\ &= \text{I.P. of } \frac{1}{2} \left( -\frac{x}{a} \right) (i \cos ax - \sin ax) \\ &= -\frac{1}{2} \frac{x}{a} \cos ax \end{aligned}$$

$$\therefore \frac{1}{D^2 + a^2} \sin ax = -\frac{x}{2a} \cos ax$$

Now if  $f(D) = D^2 + a^2$  and  $Q = \cos ax$

Then P.I. =  $\frac{1}{D^2 + a^2} \cos ax = \frac{1}{D^2 + a^2}$  (Real part of  $e^{iax}$ )

$$= \text{Real part (or R.P.) of } \frac{1}{D^2 + a^2} e^{iax}$$

$$= \text{R.P. of } -\frac{1}{2} \left( \frac{x}{a} \right) (i \cos ax - \sin ax)$$

$$= \frac{x}{2a} \sin ax$$

$$\therefore \frac{1}{D^2 + a^2} \cos ax = \frac{x}{2a} \sin ax$$

**Example 4.** Solve  $(D^2 + D + 1)y = \sin 2x$

**Solution.** Here the auxiliary equation is  $m^2 + m + 1 = 0$

$$\text{which gives } m = -\frac{1}{2} \pm i \left( \frac{1}{2} \sqrt{3} \right)$$

$$\therefore \text{C.F.} = e^{-x/2} \left\{ C_1 \cos \left( \frac{1}{2} x \sqrt{3} \right) + C_2 \sin \left( \frac{1}{2} x \sqrt{3} \right) \right\}$$

where  $C_1$  and  $C_2$  are arbitrary constants

$$\text{and P.I.} = \frac{1}{D^2 + D + 1} \sin 2x$$

$$= \frac{1}{(-2)^2 + D + 1} \sin 2x \text{ replacing } D^2 \text{ by } -2^2$$

$$= \frac{1}{D - 3} \sin 2x$$

$$= \frac{1}{(D - 3)(D + 3)} (D + 3) \sin 2x$$

$$= \frac{1}{D^2 - 9} (D + 3) \sin 2x = \frac{1}{-2^2 - 9} (D + 3) \sin 2x$$

$$= -\frac{1}{13} (D + 3) \sin 2x = \frac{-1}{13} [D(\sin 2x) + 3 \sin 2x]$$

$$= -\frac{1}{13} [2 \cos 2x + 3 \sin 2x] \text{ Since } D \text{ means differentiation with respect to } x$$

$\therefore$  The complete solution is

Linear Differential Equations with Constant Coefficients and Applications

$$y = e^{-x/2} \left[ C_1 \cos \left( \frac{1}{2} x\sqrt{3} \right) + C_2 \sin \left( \frac{1}{2} x\sqrt{3} \right) \right] - \frac{1}{13} (2 \cos 2x + 3 \sin 2x)$$

**Example 5.** Solve  $(D^2 - 5D + 6)y = \sin 3x$

**Solution.** Its auxiliary equation is  $m^2 - 5m + 6 = 0$  which gives

$$m = 2, 3$$

$\therefore$  C.F. =  $C_1 e^{2x} + C_2 e^{3x}$ , where  $C_1$  and  $C_2$  are arbitrary constants

and

$$\begin{aligned} \text{P.I.} &= \frac{1}{D^2 - 5D + 6} \sin 3x \\ &= \frac{1}{-3^2 - 5D + 6} \sin 3x, \text{ replacing } D^2 \text{ by } -3^2 \\ &= \frac{1}{-(5D + 3)} \sin 3x = \frac{-1}{(5D + 3)(5D - 3)} (5D - 3) \sin 3x \\ &= \frac{-1}{(25D^2 - 9)} (5D - 3) \sin 3x \\ &= \frac{-1}{\{25(-3^2) - 9\}} (5D - 3) \sin 3x \\ &= \frac{1}{234} [5D(\sin 3x) - 3\sin 3x] \\ &= \frac{1}{234} [5 \times 3 \cos 3x - 3 \sin 3x] \\ &= \frac{1}{78} (5 \cos 3x - \sin 3x) \end{aligned}$$

Hence the required solution is

$$y = C_1 e^{2x} + C_2 e^{3x} + \frac{1}{78} (5 \cos 3x - \sin 3x)$$

**Example 6.** Solve  $(D^3 + D^2 - D - 1)y = \cos 2x$

**Solution.** Its auxiliary equation is  $m^3 + m^2 - m - 1 = 0$

or  $(m^2 - 1)(m + 1) = 0$

or  $m = 1, -1, -1$

$\therefore$  C.F. =  $C_1 e^x + (C_2 x + C_3) e^{-x}$ , where  $C_1, C_2$  and  $C_3$  are arbitrary constants

and

$$\begin{aligned}
 \text{P.I.} &= \frac{1}{D^3 + D^2 - D - 1} \cos 2x \\
 &= \frac{1}{D(D^2) + D^2 - D - 1} \cos 2x \\
 &= \frac{1}{D(-2^2) + (-2^2) - D - 1} \cos 2x = -\frac{1}{5} \frac{1}{D + 1} \cos 2x \\
 &= -\frac{1}{5} \frac{D - 1}{(D - 1)(D + 1)} \cos 2x \\
 &= -\frac{1}{5} \frac{1}{D^2 - 1} (D - 1) \cos 2x = -\frac{1}{5} \frac{1}{(-2^2 - 1)} [D(\cos 2x) - \cos 2x] \\
 &= \frac{1}{25} (-2 \sin 2x - \cos 2x)
 \end{aligned}$$

Therefore the complete solution is

$$y = \text{C.F.} + \text{P.I.}$$

or

$$y = C_1 e^x + (C_2 x + C_3) e^{-x} - \frac{1}{25} (2 \sin 2x + \cos 2x)$$

**Example 7.** Solve  $(D^2 - 4D + 4) y = \sin 2x$ , given that  $y = 1/8$  and  $Dy = 4$  when  $x = 0$ . Find also the value of  $y$  when  $x = \pi/4$ . (Here  $D \equiv \frac{d}{dx}$ )

**Solution.** Its auxiliary equation is  $m^2 - 4m + 4 = 0$  or  $m = 2, 2$

$\therefore$  C.F. =  $(C_1 x + C_2) e^{2x}$ , where  $C_1$  and  $C_2$  are arbitrary constants

and P.I. =  $\frac{1}{D^2 - 4D + 4} \sin 2x = \frac{1}{(-2^2 - 4D + 4)} \sin 2x$

$$= -\frac{1}{4} \frac{1}{D} \sin 2x = -\frac{1}{4} \int \sin 2x \, dx = \frac{1}{8} \cos 2x$$

$\therefore$  The solution of the given equation is

$$y = \text{C.F.} + \text{P.I.} = (C_1 x + C_2) e^{2x} + \frac{1}{8} \cos 2x \quad (1)$$

$\therefore$   $Dy = C_1 e^{2x} + 2(C_1 x + C_2) e^{2x} - \frac{1}{4} \sin 2x$

Linear Differential Equations with Constant Coefficients and Applications

or  $Dy = (2 C_1 x + 2 C_2 + C_1) e^{2x} - \frac{1}{4} \sin 2x$

According to the problem  $y = \frac{1}{8}$  and  $Dy = 4$  when

$x = 0$ , so from (1) and (2) we get

$$\frac{1}{8} = C_2 e^0 + \frac{1}{8} \cos 0 = C_2 + \frac{1}{8} \text{ or } C_2 = 0$$

And  $4 = (2 C_2 + 4) e^0 = \frac{1}{4} \sin 0 = C_1 \therefore C_2 = 0$

or  $C_1 = 4$

Substituting these values of  $C_1, C_2$  in (1) we get

$$y = 4 x e^{2x} + \frac{1}{8} \cos 2x$$

when  $x = \pi/4, y = 4 \left( \frac{\pi}{4} \right) e^{\pi/2} + \frac{1}{8} \cos (\pi/2) = \pi e^{\pi/2}$

**Example 8.** Solve  $(D^3 + a^2 D) y = \sin ax$

**Solution.** Here the auxiliary equation is

$$m^3 + a^2 m = 0 \text{ or } m = 0, \pm ai$$

$\therefore$  C.F. =  $C_1 + C_2 \cos ax + C_3 \sin ax$ , where  $C_1, C_2$  and  $C_3$  are arbitrary constants and

$$\text{P.I.} = \frac{1}{D^3 + a^2 D} \sin ax = \frac{1}{D^3 + a^2 D} \text{ (Imaginary part of } e^{iax} \text{)}$$

$$= \text{I.P. of } \frac{1}{D^3 + a^2 D} e^{iax}$$

$$= \text{I.P. of } e^{iax} \frac{1}{(D + ia)^3 + a^2 (D + ia)} 1$$

$$= \text{I.P. of } e^{iax} \frac{1}{D^3 + 3ia D^2 - 2a^2 D} 1$$

$$= \text{I.P. of } e^{iax} \frac{1}{-2a^2 D \left[ 1 - \frac{3i}{2a} D - \frac{1}{2a^2} D^2 \right]}$$



$$\begin{aligned}
 &= \text{I.P. of } \frac{e^{iax}}{-2a^2} \frac{1}{D} \left( 1 - \frac{3i}{2a} D - \frac{1}{2a^2} D^2 \right)^{-1} 1 \\
 &= \text{I.P. of } \frac{e^{iax}}{-2a^2} \frac{1}{D} \left( 1 - \frac{3i}{2a} D - \frac{1}{2a^2} D^2 \right)^{-1} 1 \\
 &= \text{I.P. of } \frac{e^{iax}}{-2a^2} \frac{1}{D} \left( 1 + \frac{3i}{2a} D + \dots \right) (1) \\
 &= \text{I.P. of } \frac{e^{iax}}{-2a^2} \frac{1}{D} (1) = \text{I.P. of } \frac{e^{iax}}{-2a^2} (x) \\
 &= -\left( \frac{1}{2a^2} \right) x \sin ax
 \end{aligned}$$

∴ The required solution is  $y = \text{C.F.} + \text{P.I.}$

or  $y = C_1 + C_2 \cos ax + C_3 \sin ax - \frac{x}{2a^2} \sin ax$

**Example 9.** Solve  $\frac{d^4y}{dx^4} - m^4y = \sin mx$

(I.A.S. 1991)

**Solution.** We have  $\frac{d^4y}{dx^4} - m^4y = \sin mx$

or  $(D^4 - m^4) y = \sin mx, D = \frac{d}{dx}$

Here the auxiliary equation is

$$M^4 - m^4 = 0$$

$$\Rightarrow (M^2 - m^2) (M^2 + m^2) = 0$$

∴  $M = -m, m, \pm mi$

Therefore, C<sub>1</sub> F =  $C_1 e^{mx} + C_2 e^{-mx} + C_3 \cos mx + C_4 \sin mx$

where  $C_1, C_2, C_3$  and  $C_4$  are arbitrary constants

$$\text{P.I.} = \frac{1}{D^4 - m^2} \sin mx = \frac{1}{(D^2 - m^2) (D^2 + m^2)} \sin mx$$

$$\begin{aligned} &= \frac{1}{(D^2 + m^2)} \left\{ \frac{1}{D^2 - m^2} \sin mx \right\} = \frac{1}{D^2 + m^2} \left\{ \frac{1}{-m^2 - m^2} \sin mx \right\} \text{ replacing } D^2 \text{ by } -m^2 \\ &= \frac{1}{D^2 + m^2} \left( -\frac{1}{2m^2} \sin mx \right) = -\frac{1}{2m^2} \left( \frac{1}{D^2 + m^2} \sin mx \right) \\ &= -\frac{1}{2m^2} \left( -\frac{x}{2m} \cos mx \right) = \frac{x}{4m^3} \cos mx \\ \therefore \frac{1}{D^2 + a^2} \sin ax &= -\frac{x}{2a} \cos ax, \text{ if } f(-a^2) = 0 \end{aligned}$$

Hence the required solution is  $y = C.F + P.I$

$$\text{or } y = C_1 e^{mx} + C_2 e^{-mx} + C_3 \cos mx + C_4 \sin mx + \frac{x}{4m^3} \cos mx$$

**Example 10.** Solve  $\frac{d^3y}{dx^3} - 3 \frac{d^2y}{dx^2} + 4 \frac{dy}{dx} - 2y = e^x + \cos x$

(I.A.S. 1999, U.P.T.U. 2001, 2006)

**Solution.** The given can be rewritten as

$$(D^3 - 3D^2 + 4D - 2)y = e^x + \cos x, D \equiv \frac{d}{dx}$$

Here the auxiliary equation is

$$\begin{aligned} &m^3 - 3m^2 + 4m - 2 = 0 \\ \Rightarrow &(m - 1)(m^2 - 2m + 2) = 0 \end{aligned}$$

i.e.  $m = 1, 1 \pm i$

$\therefore$  C.F. =  $C_1 e^x + e^x (C_2 \cos x + C_3 \sin x)$ , where  $C_1, C_2$  and  $C_3$  are arbitrary constants.

$$\& \text{ P.I.} = \frac{1}{D^3 - 3D^2 + 4D - 2} (e^x + \cos x)$$

$$\begin{aligned}
 &= \frac{1}{(D-1)(D^2-2D+2)} e^x + \frac{1}{D^3-3D^2+4D-2} \cos x \\
 &= \frac{1}{(D-1)} \left\{ \frac{1}{1-2+2} e^x \right\} + \frac{1}{(-1)^2 D - 3(-1^2) + 4D - 2} \cos x \\
 &= \frac{1}{D-1} e^x + \frac{1}{3D+1} \cos x \\
 &= e^x \frac{1}{(D+1)-1} + \frac{(3D-1)}{9D^2-1} \cos x \\
 &= e^x \frac{1}{D} + \frac{3D-1}{9(-1^2)-1} \cos x \\
 &= x e^x - \frac{1}{10} (3D-1) \cos x \\
 &= x e^x - \frac{1}{10} (3D \cos x - \cos x) \\
 &= x e^x - \frac{1}{10} (-3 \sin x - \cos x) \\
 &= x e^x + \frac{1}{10} (3 \sin x + \cos x)
 \end{aligned}$$

Hence, the required solution is  $y = C.F + P.I$

$$y = C_1 e^x + e^x (C_2 \cos x + C_3 \sin x) + x e^x + \frac{1}{10} (3 \sin x + \cos x)$$

**(C) To find P.I. when Q is of the form  $x^m$**

In this case  $P.I = \frac{1}{f(D)} x^m$ , where  $m$  is a positive integer

To evaluate this we take common the lowest degree from  $f(D)$ , so that the remaining factor reduces to the form  $[1 + F(D)]$  or  $[1 - F(D)]$ . Now take this factor in the numerator with a negative index and expand it by Binomial theorem in powers of  $D$  upto the term  $D^m$ , (Since other higher derivatives of  $x^m$  will be zero) and operate upon  $x^m$ . The following examples will illustrate the method.

**Example 11.** Solve  $\frac{d^2y}{dx^2} + \frac{dy}{dx} - 6y = x$

**Solution.** The given equation can be written as

$$(D^2 + D - 6) y = x$$

Its auxiliary equation is  $m^2 + m - 6 = 0$  or  $m = 2, -3$

$\therefore$  C.F. =  $C_1 e^{2x} + C_2 e^{-3x}$ , where  $C_1$  and  $C_2$  are arbitrary constants

and 
$$\begin{aligned} \text{P.I.} &= \frac{1}{D^2 + D - 6} x = \frac{1}{-6 \left( 1 - \frac{1}{6} D - \frac{1}{6} D^2 \right)} x \\ &= -\frac{1}{6} \left[ 1 - \frac{1}{6} (D + D^2) \right]^{-1} x \\ &= -\frac{1}{6} \left[ 1 + \frac{1}{6} (D + D^2) + \dots \right] x \\ &= -\frac{1}{6} \left( x + \frac{1}{6} \right) = \frac{-1}{36} (6x + 1) \end{aligned}$$

∴ The required solution is  $y = \text{C.F.} + \text{P.I.}$

$$y = C_1 e^{2x} + C_2 e^{-3x} - \frac{1}{36} (6x + 1)$$

**Example 12.** Solve  $(D^3 - D^2 - 6D) y = x^2 + 1$  Where  $D \equiv \frac{d}{dx}$

**(Bihar P.C.S. 1993)**

**Solution.** Here the auxiliary equation is  $m^3 - m^2 - 6m = 0$

or  $m(m^2 - m - 6) = 0$

or  $m(m - 3)(m + 2) = 0$

or  $m = 0, 3, -2$

∴  $C_1 F = C_1 e^{0x} + C_2 e^{3x} + C_3 e^{-2x}$

or C.F. =  $C_1 + C_2 e^{3x} + C_3 e^{-2x}$ , where  $C_1, C_2$  and  $C_3$  are arbitrary constants.

and 
$$\begin{aligned} \text{P.I.} &= \frac{1}{D^2 - D^2 - 6D} (x^2 + 1) \\ &= \frac{1}{-6D \left( 1 + \frac{1}{6} D - \frac{1}{6} D^2 \right)} (x^2 + 1) \\ &= -\frac{1}{6D} \left[ 1 - \left( \frac{D}{6} - \frac{D^2}{6} \right) + \left( \frac{D}{6} - \frac{D^2}{6} \right)^2 \dots \right] (x^2 + 1) \\ &= -\frac{1}{6D} \left[ 1 + x^2 - \frac{1}{6} D(x^2) + \frac{7}{36} D^2(x^2) \right] \quad \because D(1) = 0 \end{aligned}$$

$$\begin{aligned}
 &= -\frac{1}{6D} \left( 1 + x^2 - \frac{1}{3}x + \frac{7}{18} \right) \\
 &= -\frac{1}{6} \int \left( 1 + x^2 - \frac{x}{3} + \frac{7}{18} \right) dx \qquad \because D \equiv \frac{d}{dx} \\
 &= -\frac{1}{6} \left[ x + \frac{x^3}{3} - \frac{1}{6}x^2 + \frac{7}{18}x \right] = -\frac{1}{6} \left( \frac{25}{18}x + \frac{x^3}{3} - \frac{x^2}{6} \right)
 \end{aligned}$$

∴ The required solution is  $y = \text{C.F.} + \text{P.I.}$

or 
$$y = C_1 + C_2 e^{3x} + C_3 e^{-2x} - \frac{25}{108}x - \frac{x^3}{18} + \frac{x^2}{36}$$

**Example 13.** Solve  $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + 2y = x + e^x \cos x$

(U.P.T.U. 2002)

**Solution.** Given differential equation is

$$(D^2 - 2D + 2)y = x + e^x \cos x$$

Here the auxiliary equation is

$$m^2 - 2m + 2 = 0 \Rightarrow m = 1 \pm i$$

∴ C.F =  $e^x (C_1 \cos x + C_2 \sin x)$ , where  $C_1$  and  $C_2$  are arbitrary constants,

$$\begin{aligned}
 \text{and P.I.} &= \frac{1}{D^2 - 2D + 2} (x + e^x \cos x) \\
 &= \frac{1}{D^2 - 2D + 2} x + \frac{1}{D^2 - 2D + 2} (e^x \cos x) \\
 &= \frac{1}{2 \left[ 1 - D + \frac{D^2}{2} \right]} x + e^x \frac{1}{\left\{ (D+1)^2 - 2(D+1) + 2 \right\}} \cos x \\
 &= \frac{1}{2} \left( 1 - D + \frac{D^2}{2} \right)^{-1} (x) + e^x \frac{1}{D^2 + 1} \cos x \quad \text{Here } f(-a^2) = 0 \\
 &= \frac{1}{2} \left( 1 + D - \frac{D^2}{2} + \dots \right) (x) + e^x \frac{x}{2} \sin x \\
 &= \frac{1}{2} \left\{ x + D(x) - \frac{1}{2} D^2(x) \right\} + \frac{x e^x}{2} \sin x \\
 &= \frac{1}{2} (x + 1) + \frac{x e^x}{2} \sin x
 \end{aligned}$$

∴ The required solutions  $y = \text{C.F.} + \text{P.I.}$

$$\text{or } y = e^x (C_1 \cos x + C_2 \sin x) + \frac{1}{2} (x + 1) + \frac{x e^x}{2} \sin x$$

**Example 15.** Find the complete solution of

$$\frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 2y = x e^{3x} + \sin 2x \quad (\text{U.P.T.U. 2003})$$

**Solution.** The given differential equation is

$$(D^2 - 3D + 2)y = x e^{3x} + \sin 2x$$

Here the auxilliary equation is

$$m^2 - 3m + 2 = 0 \Rightarrow (m - 1)(m - 2) = 0 \Rightarrow m = 1, 2$$

$\therefore$  C.F =  $C_1 e^x + C_2 e^{2x}$ , where  $C_1$  and  $C_2$  being arbitrary constants

$$\begin{aligned} \& \text{ P.I.} &= \frac{1}{D^2 - 3D + 2} (x e^{3x} + \sin 2x) \\ &= \frac{1}{D^2 - 3D + 2} (x e^{3x}) + \frac{1}{D^2 - 3D + 2} (\sin 2x) \\ &= e^{3x} \frac{1}{(D + 3)^2 - 3(D + 3) + 2} (x) + \frac{1}{-2^2 - 3D + 2} (\sin 2x) \\ &= e^{3x} \frac{1}{D^2 + 3D + 2} (x) + \frac{1}{-3D - 2} \sin 2x \\ &= e^{3x} \frac{1}{2} \frac{1}{\left(1 + \frac{3}{2}D + \frac{D^2}{2}\right)} (x) - \frac{1}{(3D + 2)} (\sin 2x) \\ &= \frac{1}{2} e^{3x} \left(1 + \frac{3D}{2} + \frac{D^2}{2}\right)^{-1} (x) - \frac{3D - 2}{9D^2 - 4} (\sin 2x) \\ &= \frac{1}{2} e^{3x} \left[1 - \frac{3D}{2} - \frac{D^2}{2} \dots\dots\dots\right] (x) - \frac{(3D - 2)}{9(-2^2) - 4} (\sin 2x) \text{ replacing } D^2 \text{ by } -2^2 \end{aligned}$$

$$\begin{aligned} &= \frac{1}{2} e^{3x} \left( x - \frac{3}{2} \right) + \frac{1}{40} (3D \sin 2x - 2 \sin 2x) \\ &= \frac{1}{2} e^{3x} \left( x - \frac{3}{2} \right) + \frac{1}{40} (6 \cos 2x - 2 \sin 2x) \\ &= e^{3x} \left( \frac{x}{2} - \frac{3}{4} \right) + \frac{1}{20} (3 \cos 2x - \sin 2x) \end{aligned}$$

Hence the required solution is  $y = C.F + P.I$

$$\text{or } y = C_1 e^x + C_2 e^{2x} + e^{3x} \left( \frac{x}{2} - \frac{3}{4} \right) + \frac{1}{20} (3 \cos 2x - \sin 2x)$$

**Example 16.** A body executes damped forced vibrations given by the equation.

$$\frac{d^2x}{dt^2} + 2k \frac{dx}{dt} + b^2x = e^{-kt} \sin wt$$

Solve the equation for both the cases, when  $w^2 \neq b^2 - k^2$  and  $w^2 = b^2 - k^2$

[U.P.T.U. 2001, 03, 04 (C.O) 2005]

**Solution.** We have  $\frac{d^2x}{dt^2} + 2k \frac{dx}{dt} + b^2x = e^{-kt} \sin wt$

or  $(D^2 + 2kD + b^2)x = e^{-kt} \sin wt$

Here the auxiliary equation is  $m^2 + 2km + b^2 = 0$

$$\Rightarrow m = -k \pm i\sqrt{(b^2 - k^2)}$$

Case 1. When  $w^2 \neq b^2 - k^2$

$$C.F = e^{-kt} [C_1 \cos \sqrt{(b^2 - k^2)} t + C_2 \sin \sqrt{(b^2 - k^2)} t]$$

where  $C_1$  and  $C_2$  are arbitrary constants

$$\& P.I. = \frac{1}{D^2 + 2kD + b^2} (e^{-kt} \sin wt)$$

Linear Differential Equations with Constant Coefficients and Applications

$$\begin{aligned}
 &= e^{-kt} \frac{1}{(D-k)^2 + 2k(D-k) + b^2} (\sin wt) \\
 &= e^{-kt} \frac{1}{D^2 + (b^2 - k^2)} (\sin wt) = e^{-kt} \frac{1}{w^2 + (b^2 - k^2)} \sin wt \\
 &= \frac{e^{-kt} \sin wt}{b^2 - k^2 - w^2}
 \end{aligned}$$

Here, complete solution is  $y = C.F + P.I$

or 
$$y = e^{-kt} \left[ C_1 \cos \sqrt{(b^2 - k^2)} t + C_2 \sin \sqrt{(b^2 - k^2)} t \right] + \frac{e^{-kt} \sin wt}{b^2 - k^2 - w^2}$$

Case 2. When  $w^2 = b^2 - k^2$

C.F. =  $e^{-kt} (C_1 \cos wt + C_2 \sin wt)$

& P.I. = 
$$\frac{1}{D^2 + 2kD + b^2} (e^{-kt} \sin wt)$$

$$\begin{aligned}
 &= e^{-kt} \frac{1}{(D-k)^2 + 2k(D-k) + b^2} \sin wt \\
 &= e^{-kt} \frac{1}{D^2 + b^2 - k^2} \sin wt \text{ Here } f(-a^2) = 0 \\
 &= e^{-kt} \left( -\frac{t}{2w} \cos wt \right) \\
 &= -\frac{t e^{-kt}}{2w} \cos wt
 \end{aligned}$$

Hence, the required solution is  $y = C.F + P.I$

or 
$$y = e^{-kt} (C_1 \cos wt + C_2 \sin wt) - \frac{t}{2w} e^{-kt} \cos wt$$

**Example 17.** Solve  $(D^2 - 2D + 1) y = x e^x \sin x$

(Bihar P.C.S., 1997; 2007, U.P.P.C.S., 2001; L.D.A. 1995, U.P.T.U. 2005)

**Solution.** Here the auxiliary equation is

$m^2 - 2m + 1 = 0 \Rightarrow (m - 1)^2 = 0 \therefore m = 1, 1$

$\therefore$  C.F. =  $(C_1 + C_2 x) e^x$  where  $C_1$  and  $C_2$  are arbitrary constants

and 
$$P.I. = \frac{1}{D^2 - 2D + 1} x e^x \sin x$$



$$\begin{aligned}
 &= e^x \frac{1}{(D+1)^2 - 2(D+1) + 1} (x \sin x) \\
 &= e^x \frac{1}{D^2 + 1 + 2D - 2D - 2 + 1} x \sin x \\
 &= e^x \frac{1}{D^2} (x \sin x) \\
 &= e^x \frac{1}{D} \int x \sin x \, dx = e^x \frac{1}{D} [-x \cos x + \sin x] \\
 &= e^x [-\int x \cos x \, dx + \int \sin x \, dx] \\
 &= e^x [-x \sin x - \cos x - \cos x] \\
 &= -e^x (x \sin x + 2 \cos x)
 \end{aligned}$$

Hence, the required solution is  $y = C.F + P.I$

$$y = (C_1 + C_2 x) e^x - e^x (x \sin x + 2 \cos x)$$

**Example 18.** Find the complete solution of the differential equation

$$\frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + y = x e^x \cos x \quad \text{(U.P.T.U. 2009)}$$

**Solution.** Here the auxiliary equation is

$$m^2 - 2m + 1 = 0 \Rightarrow (m - 1)^2 = 0; \therefore m = 1, 1$$

$$\therefore C.F. = (C_1 + C_2 x) e^x$$

and 
$$P.I. = \frac{1}{D^2 - 2D + 1} x e^x \cos x$$

$$= e^x \frac{1}{D^2} (x \cos x)$$

$$= e^x (-x \cos x + 2 \sin x)$$

Therefore, complete solution is  $y = C.F + P.I$

or 
$$y = (C_1 + C_2 x) e^x + e^x (2 \sin x - x \cos x)$$

**Example 19.** Solve  $(D^4 + 6D^3 + 11D^2 + 6D) y = 20 e^{-2x} \sin x$

[U.P.T.U. (C.O.) 2005]

**Solution.** Here the auxiliary equation is

$$m^4 + 6m^3 + 11m^2 + 6m = 0$$

$$\Rightarrow m (m^3 + 6m^2 + 11m + 6) = 0$$

Linear Differential Equations with Constant Coefficients and Applications

$$\Rightarrow m(m+1)(m+2)(m+3) = 0$$

$$\therefore m = 0, -1, -2, -3$$

$$\therefore \text{C.F.} = C_1 + C_2 e^{-x} + C_3 e^{-2x} + C_4 e^{-3x}$$

$$\begin{aligned} \& \text{ P.I} &= \frac{1}{D^4 + 6D^3 + 11D^2 + 6D} 20 e^{-2x} \sin x \\ &= 20 e^{-2x} \frac{1}{(D-2)^4 + 6(D-2)^3 + 11(D-2)^2 + 6(D-2)} \sin x \\ &= 20 e^{-2x} \frac{1}{(D-2)(D^3 - D)} \sin x \\ &= 20 e^{-2x} \frac{1}{(D-2)\{(D)^2 D - D\}} \sin x \\ &= 20 e^{-2x} \frac{1}{(D-2)\{(-1)^2 D - D\}} \sin x \text{ replacing } D^2 \text{ by } -1^2 \\ &= 20 e^{-2x} \frac{1}{(D-2)(-D-D)} \sin x \\ &= 20 e^{-2x} \frac{1}{4D - 2D^2} \sin x = 20 e^{-2x} \frac{1}{4D - 2(-1)^2} \sin x \\ &= 20 e^{-2x} \frac{1}{4D + 2} \sin x \\ &= 10 e^{-2x} \frac{2D - 1}{4D^2 - 1} \sin x = 10 e^{-2x} \frac{(2 \cos x - \sin x)}{4(-1) - 1} \\ &= 2 e^{-2x} (\sin x - 2 \cos x) \end{aligned}$$

Hence, Complete solution is  $y = \text{C.F.} + \text{P.I}$

$$y = C_1 + C_2 e^{-x} + C_3 e^{-2x} + C_4 e^{-3x} + 2e^{2x} (\sin x - 2\cos x)$$

**Example 20.** Solve  $(D^2 - 4D - 5)y = e^{2x} + 3 \cos(4x + 3)$   $D \equiv \frac{d}{dx}$

(U.P.T.U. 2008)

**Solution.** We have  $(D^2 - 4D - 5)y = e^{2x} + 3 \cos(4x + 3)$

Here the auxiliary equation is

$$m^2 - 4m - 5 = 0 \Rightarrow (m - 5)(m + 1) = 0 \therefore m = 5, -1$$

$$\text{C.F.} = C_1 e^{5x} + C_2 e^{-x}$$

$$\begin{aligned}
 \& \text{ P.I.} &= \frac{1}{D^2 - 4D - 5} \{e^{2x} + 3 \cos (4x + 3)\} \\
 &= \frac{1}{D^2 - 4D - 5} e^{2x} + 3 \frac{1}{D^2 - 4D - 5} \{\cos 4x \cos 3 - \sin 4x \sin 3\} \\
 &= \frac{1}{(2)^2 - 4(2) - 5} e^{2x} + 3 \cos 3 \frac{1}{D^2 - 4D - 5} \cos 4x - 3 \sin 3 \frac{1}{D^2 - 4D - 5} \sin 4x \\
 &= -\frac{1}{9} e^{2x} + 3 \cos 3 \frac{1}{-4^2 - 4D - 5} \cos 4x - 3 \sin 3 \frac{1}{-4^2 - 4D - 5} \sin 4x \\
 &= -\frac{1}{9} e^{2x} + 3 \cos 3 \frac{1}{-16 - 4D - 5} \cos 4x - 3 \sin 3 \frac{1}{-16 - 4D - 5} \sin 4x \\
 &= -\frac{1}{9} e^{2x} + 3 \cos 3 \frac{1}{-(4D + 21)} \cos 4x + 3 \sin 3 \frac{1}{4D + 2} \sin 4x \\
 &= -\frac{1}{9} e^{2x} - 3 \cos 3 \frac{(4D - 21)}{(6D^2 - 44)} \cos 4x + 3 \sin 3 \frac{(4D - 21)}{16D^2 - 441} \sin 4x \\
 &= -\frac{1}{9} e^{2x} - 3 \cos 3 \frac{(4D - 21)}{4(-4^2) - 441} \cos 4x + 3 \sin 3 \frac{4D - 21}{16(-4^2)441} \sin 4x \\
 &= -\frac{1}{9} e^{2x} - 3 \cos 3 \frac{(-16 \sin 4x - 21 \cos 4x)}{-697} + 3 \sin 3 \frac{16 \cos 4x - 21 \sin 4x}{-697} \\
 &= -\frac{1}{9} e^{2x} - \frac{3}{697} \cos 3 (16 \sin 4x + 21 \cos 4x) - \frac{3 \sin 3}{697} (16 \cos 4x - 21 \sin 4x) \\
 &= -\frac{1}{9} e^{2x} - \frac{3}{697} [16 \sin 4x \cos 3 + 21 \cos 4x \cos 3 + 16 \cos 4x \sin 3 - 21 \sin 4x \sin 3] \\
 &= -\frac{1}{9} e^{2x} - \frac{3}{697} [16 \sin (4x + 3) + 21 \cos (4x + 3)]
 \end{aligned}$$

Hence the required solution is  $y = \text{C.F.} + \text{P.I.}$

$$\text{or } y = C_1 e^{5x} + C_2 e^{-x} - \frac{1}{9} e^{2x} - \frac{3}{697} [16 \sin (4x + 3) + 21 \cos (4x + 3)]$$

**Example 21.** Solve  $\frac{d^2y}{dx^2} + 2 \frac{dy}{dx} + 10y + 37 \sin 3x = 0$ , and find the value of  $y$

when  $x = \pi/2$ , if it is given that  $y = 3$  and  $\frac{dy}{dx} = 0$  when  $x = 0$  (I.A.S. 1996)

**Solution.** The given equation can be written as

$$(D^2 + 2D + 10)y = -37 \sin 3x$$

its auxiliary equation is  $m^2 + 2m + 10 = 0$ , which gives

Linear Differential Equations with Constant Coefficients and Applications

$$m = \frac{1}{2} \left[ -2 \pm \sqrt{(4 - 40)} \right] = -1 \pm 3i$$

$\therefore$  C.F. =  $e^{-x} [C_1 \cos 3x + C_2 \sin 3x]$ , where  $C_1$  &  $C_2$  are arbitrary constants,

and

$$\begin{aligned} \text{P.I.} &= \frac{1}{D^2 + 2D + 10} (-37 \sin 3x) \\ &= -37 \frac{1}{D^2 + 2D + 10} \sin 3x \\ &= -37 \frac{1}{-3^2 + 2D + 10} \sin 3x \\ &= -37 \frac{1}{2D + 1} \sin 3x = -37 \frac{1}{(4D^2 - 1)} (2D - 1) \sin 3x \\ &= -37 \frac{1}{\{4(-3^2) - 1\}} (2D - 1) \sin 3x \\ &= (2D - 1) \sin 3x = 6 \cos 3x - \sin 3x \end{aligned}$$

Hence the required solution of the given differential equation is

$$y = e^{-x} (C_1 \cos 3x + C_2 \sin 3x) + 6 \cos 3x - \sin 3x \quad (1)$$

Differentiating both sides of (1) with respect to  $x$ ,

we get

$$\frac{dy}{dx} = e^{-x} (-3 C_1 \sin 3x + 3 C_2 \cos 3x) - e^{-x} (C_1 \cos 3x + C_2 \sin 3x) - 18 \sin 3x - 3 \cos 3x \quad (2)$$

it is given that when  $x = 0$ ,  $y = 3$  and  $\frac{dy}{dx} = 0$

$\therefore$  From (1) and (2) we, have

$$3 = C_1 + 6 \quad (3)$$

and

$$0 = 3 C_2 - C_1 - 3 \quad (4)$$

From (3) and (4) we get  $C_1 = -3$  and  $C_2 = 0$

$\therefore$  From (1), we have

$$y = -3 e^{-x} \cos 3x + 6 \cos 3x - \sin 3x \quad (5)$$

$\therefore$  when  $x = \pi/2$  we have from (5)

$$y = 3e^{-\pi/2} \cos \frac{3\pi}{2} + 6 \cos \frac{3\pi}{2} - \sin \frac{3\pi}{2}$$

$$y = -3e^{-\frac{\pi}{2}} \cdot 0 + 6 \cdot 0 + 1$$

$$\text{or } y = 1$$

**Example 22.** Solve  $(D^2 + 1)^2 = 24 x \cos x$  given the initial conditions  $x = 0, y = 0, Dy = 0, D^2y = 0, D^3y = 12$  **(I.A.S. 2001)**

**Solution.** Here the auxiliary equation is

$$(m^2 + 1)^2 = 0$$

$$\text{or } m = \pm i \text{ (twice)}$$

$$\therefore \text{C.F.} = (C_1 x + C_2) \cos x + (C_3 x + C_4) \sin x$$

$$\text{and P.I.} = \frac{1}{(D^2 + 1)^2} (24 x \cos x)$$

$$= \text{R.P. of } \frac{1}{(D^2 + 1)^2} 24 x e^{ix}$$

$$= \text{R.P. of } 24 e^{ix} \frac{1}{[(D + i)^2 + 1]^2} x$$

$$= \text{R.P. of } 24 e^{ix} \frac{1}{(D^2 + 2iD)^2} x$$

$$= \text{R.P. of } 24 e^{ix} \frac{1}{-4D^2 \left[1 + \frac{D}{2i}\right]^2} x$$

$$= \text{R.P. of } -6 e^{ix} \frac{1}{D^2} \left(1 + iD - \frac{3}{4} D^2 + \dots\right) x$$

$$= \text{R.P. of } -6 e^{ix} \frac{1}{D^2} (x + i)$$

$$= \text{R.P. of } -6 e^{ix} \left(\frac{1}{6} x^3 + \frac{1}{2} i x^2\right)$$

$$= \text{R.P. of } -(\cos x + i \sin x) (x^3 + 3i x^2)$$

$$= -x^2 \cos x + 3x^2 \sin x$$

$\therefore$  The solution of the given differential equation is

Linear Differential Equations with Constant Coefficients and Applications

$$y = (C_1 x + C_2) \cos x + (C_3 x + C_4) \sin x - x^3 \cos x + 3x^2 \sin x$$

or  $y = (C_1 x + C_2 - x^3) \cos x + (C_3 x + C_4 + 3x^2) \sin x$  (1)

$$\therefore Dy = (C_1 - 3x^2) \cos x - (C_1 x + C_2 - x^3) \sin x + (C_3 x + C_4 + 3x^2) \cos x + (C_3 + 6x) \sin x$$

or  $Dy = (C_1 + C_4 + C_3 x) \cos x + (C_3 - C_2 + 6x - C_1 x + x^3) \sin x$  (2)

$$\therefore D^2y = -(C_1 + C_4 + C_3 x) \sin x + C_3 \cos x + (C_3 - C_2 + 6x - C_1 x + x^3) \cos x + (6 - 4 + 3x^2) \sin x$$

or  $D^2y = (6 - 2C_1 - C_4 - C_3 x + 3x^2) \sin x + (2C_3 - C_2 + 6x - C_1 x + x^3) \cos x$  (3)

$$\therefore D^2y = (6 - 2C_1 - C_4 - C_3 x + 3x^2) \cos x + (-C_3 + 6x) \sin x - (2C_3 - C_2 + 6x - C_1 x + x^3) \sin x + (6 - C_1 + 3x^2) \cos x$$

or  $D^3y = (12 - 3C_1 - C_4 - C_3 x + 6x^2) \cos x + (C_2 - 3C_3 - 6x + C_1 x + 6x - x^2) \sin x$  (4)

Since we are given that  $x = 0, y = 0, Dy = 0, D^2y = 0, D^3y = 12$  so from (1), (2), (3) and (4) we get

$$0 = C_2 \quad (5)$$

$$0 = C_1 + C_4 \quad (6)$$

$$0 = 2C_3 - C_2 \quad (7)$$

$$12 = 12 - 3C_1 - C_4 \quad (8)$$

From (5) and (7) we get  $C_2 = 0 = C_3$

From (6) and (8) we get  $C_1 = 0 = C_4$

$\therefore$  From (1) the required solution is

$$y = 3x^2 \sin x - x^3 \cos x$$

(D) P.I when Q is of the form  $x \cdot V$ , where V is any function of x

$$\frac{1}{f(D)} (x \cdot V) = x \cdot \frac{1}{f(D)} V - \frac{f'(D)}{\{f(D)\}^2} V$$

**Example 23.** Solve  $(D^2 - 2D + 1)y = x \sin x$

**Solution.** Here the auxiliary equation is  $m^2 - 2m + 1 = 0$

or  $(m - 1)^2 = 0$  or  $m = 1, 1$

$\therefore$  C.F. =  $(C_1 x + C_2) e^x$ , where  $C_1$  and  $C_2$  are arbitrary constants

and  $P.I = \frac{1}{D^2 - 2D + 1} x \sin x$

$$\begin{aligned}
 &= x \frac{1}{D^2 - 2D + 1} \sin x - \frac{(2D - 2)}{(D^2 - 2D + 1)^2} \sin x \\
 &= x \frac{1}{-1^2 - 2D + 1} \sin x - \frac{(2D - 2)}{(-1^2 - 2D + 1)^2} \sin x \\
 &= -\frac{x}{2} \frac{1}{D} \sin x - \frac{1}{4D^2} (2D - 2) \sin x \\
 &= -\frac{x}{2} \int \sin x \, dx - \frac{1}{2} \frac{1}{D^2} (\cos x - \sin x) \\
 &= \frac{x}{2} \cos x - \frac{1}{2} \frac{1}{D} \int (\cos x - \sin x) \, dx \\
 &= \frac{x}{2} \cos x - \frac{1}{2} \frac{1}{D} (\sin x + \cos x) \\
 &= \frac{x}{2} \cos x - \frac{1}{2} \int (\sin x + \cos x) \, dx \\
 &= \frac{x}{2} \cos x - \frac{1}{2} (-\cos x + \sin x) \\
 &= \frac{1}{2} (x \cos x + \cos x - \sin x)
 \end{aligned}$$

∴ The required solution is  $y = C.F + P.I$

or 
$$y = (C_1 x + C_2) e^x + \frac{1}{2} (x \cos x + \cos x - \sin x)$$

(E) To show that  $\frac{1}{D - a} Q = e^{ax} \int Q e^{-ax} \, dx$ , where  $Q$  is a function of  $x$ .

**Proof** Let  $y = \frac{1}{D - a} Q$

Then  $(D - a)y = Q$ , operating both sides with  $D - a$

or  $\frac{dy}{dx} - ay = Q$ , Which is a linear equation in  $y$  whose integrating factor is  $e^{-ax}$  its

solution is  $ye^{-ax} = \int Q e^{-ax} \, dx$ , neglecting the constant of integration as P.I. is required

or 
$$y = e^{ax} \int Q e^{-ax} \, dx$$

or 
$$\frac{1}{D - a} Q = e^{ax} \int Q e^{-ax} \, dx$$

**Example 24.** Solve  $(D^2 + a^2)y = \sec ax$  (U.P.P.C.S. 1971, 1973, 1977)

**Solution.** The auxiliary equation is  $m^2 + a^2 = 0$  or  $m = \pm ai$

$\therefore$  C.F. =  $C_1 \cos ax + C_2 \sin ax$ , where  $C_1$  and  $C_2$  are arbitrary constants

$$\text{and P.I} = \frac{1}{D^2 + a^2} \sec ax = \frac{1}{2ia} \left[ \frac{1}{D - ia} - \frac{1}{D + ia} \right] \sec ax \quad (1)$$

$$\begin{aligned} \text{Now } \frac{1}{D - ia} \sec ax &= e^{iax} \int \sec ax \cdot e^{-iax} dx \\ &= e^{iax} \int \sec ax \cdot (\cos ax - i \sin ax) dx \\ &= e^{iax} \int (1 - i \tan ax) dx \\ &= e^{iax} \left[ x + \left( \frac{i}{a} \right) \log \cos ax \right] \end{aligned}$$

similarly,

$$\begin{aligned} \frac{1}{D + ia} \sec ax &= e^{-iax} \int \sec ax \cdot e^{iax} dx \\ &= e^{-iax} \int \sec ax (\cos ax + i \sin ax) dx \\ &= e^{-iax} \int (1 + i \tan ax) dx \\ &= e^{-iax} \left[ x - \left( \frac{i}{a} \right) \log \cos ax \right] \end{aligned}$$

$\therefore$  From (1) we, have

$$\begin{aligned} \text{P.I} &= \frac{1}{2ia} \left[ e^{iax} \left\{ x + \frac{i}{a} \log \cos ax \right\} - e^{-iax} \left\{ x - \frac{i}{a} \log \cos ax \right\} \right] \\ &= \left[ x \left( \frac{e^{iax} - e^{-iax}}{2ia} \right) + \frac{i}{a} (\log \cos ax) \cdot \left( \frac{e^{iax} + e^{-iax}}{2ia} \right) \right] \\ &= \left( \frac{x}{a} \right) \sin ax + \frac{1}{a^2} \cos ax \cdot (\log \cos ax) \end{aligned}$$

$\therefore$  The required solution is

$$y = \text{C.F.} + \text{P.I.}$$



or  $y = C_1 \cos ax + C_2 \sin ax + \left(\frac{x}{a}\right) \sin ax + \left(\frac{1}{a^2}\right) \cos ax \cdot \log \cos ax$

**Example 25.**  $(D^2 + a^2)y = \operatorname{cosec} ax$

**Solution.** Here the auxiliary equation is  $m^2 + a^2 = 0$

or  $m = \pm ai$

$\therefore$  C.F. =  $C_1 \cos ax + C_2 \sin ax$  and

$$\text{P.I.} = \frac{1}{D^2 + a^2} \operatorname{cosec} ax = \frac{1}{2ia} \left[ \frac{1}{D - ia} - \frac{1}{D + ia} \right] \operatorname{cosec} ax \quad (1)$$

$$\begin{aligned} \text{Now } \frac{1}{D - ia} \operatorname{cosec} ax &= e^{iax} \int (\operatorname{cosec} ax) e^{-iax} dx \\ &= e^{iax} \int \operatorname{cosec} ax (\cos ax - i \sin ax) dx \\ &= e^{iax} \int (\cot ax - i) dx \\ &= e^{iax} \left[ \frac{1}{a} (\log \sin ax) - ix \right] \end{aligned}$$

$$\begin{aligned} \text{and } \frac{1}{D + ia} \operatorname{cosec} ax &= e^{-iax} \int (\operatorname{cosec} ax) e^{iax} dx \\ &= e^{-iax} \int (\operatorname{cosec} ax) (\cos ax + i \sin ax) dx \\ &= e^{-iax} \int (\cot ax + i) dx \\ &= e^{-iax} \left[ \frac{1}{a} (\log \sin ax) + ix \right] \end{aligned}$$

$\therefore$  From (1) we have

$$\begin{aligned} \text{P.I.} &= \frac{1}{2ia} \left[ e^{iax} \left\{ \frac{1}{a} (\log \sin ax) - ix \right\} - e^{-iax} \left\{ \frac{1}{a} (\log \sin ax) + ix \right\} \right] \\ &= \left[ \frac{1}{a^2} (\log \sin ax) \left( \frac{e^{iax} - e^{-iax}}{2i} \right) - \frac{x}{a} \left( \frac{e^{iax} + e^{-iax}}{2} \right) \right] \\ &= \left( \frac{1}{a^2} \right) \sin ax (\log \sin ax) - \left( \frac{x}{a} \right) (\cos ax) \end{aligned}$$

$\therefore$  The required solution is  $y = \text{C.F.} + \text{P.I.}$

or  $y = C_1 \cos ax + C_2 \sin ax + \left(\frac{1}{a^2}\right) \sin ax \log (\sin ax) - \left(\frac{x}{a}\right) \cos ax$

**Example 26.**  $(D^2 + a^2) y = \tan ax$

**Solution.** Here C.F. =  $C_1 \cos ax + C_2 \sin ax$

and P.I. =  $\frac{1}{D^2 + a^2} \tan ax = \frac{1}{2ia} \left[ \frac{1}{D - ia} - \frac{1}{D + ia} \right] \tan ax$  (1)

Now  $\frac{1}{D - ia} \tan ax = e^{i ax} \int e^{-i ax} \tan ax \, dx$

$$= e^{i ax} \int (\cos ax - i \sin ax) \tan ax \, dx$$

$$= e^{i ax} \int \left[ \sin ax - i \frac{\sin^2 ax}{\cos ax} \right] dx$$

$$= e^{i ax} \int \sin ax \, dx - i e^{i ax} \int \left( \frac{1 - \cos^2 ax}{\cos ax} \right) dx$$

$$= e^{i ax} \{(-\cos ax)/a\} - i e^{i ax} \int (\sec ax - \cos ax) \, dx$$

$$= -\frac{1}{a} \{e^{i ax} \cos ax\} - i e^{i ax} \left[ \frac{1}{a} \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) - \frac{1}{a} \sin ax \right]$$

$$= -\left(\frac{1}{a}\right) e^{i ax} \left[ \cos ax + i \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) - i \sin ax \right]$$

$$= -\frac{1}{a} e^{i ax} \left[ (\cos ax - i \sin ax) + i \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) \right]$$

$$= -\frac{1}{a} e^{i ax} \left[ e^{-i ax} + i \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) \right]$$

$$= -\frac{1}{a} \left[ 1 + i e^{i ax} \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) \right]$$
 (2)

Similarly replacing  $i$  by  $-i$  we get

$$\frac{1}{D + ia} \tan ax = -\left(\frac{1}{a}\right) \left[ 1 - i e^{-i ax} \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) \right]$$
 (3)

$\therefore$  From (1), (2) and (3), we have

$$\begin{aligned}
 \text{P.I} &= \frac{1}{2ia} \left[ -\frac{1}{a} \left\{ 1 + i e^{i ax} \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) \right\} + \frac{1}{a} \left\{ 1 - i e^{-i ax} \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) \right\} \right] \\
 &= \frac{1}{2i a} \left[ -\frac{i}{a} (e^{i ax} + e^{-i ax}) \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) \right] \\
 &= -\frac{1}{a^2} (\cos ax) \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right)
 \end{aligned}$$

∴ The required solution is  $y = \text{C.F} + \text{P.I}$

$$\text{or } y = C_1 \cos ax + C_2 \sin ax - \frac{1}{a^2} \cos ax \log \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right)$$

### Miscellaneous solved Examples

**Example 27.** Solve  $(D^2 - 2D + 1) y = x^2 e^{3x}$

**Solution.** Here the auxiliary equation is

$$m^2 - 2m + 1 = 0$$

$$\text{or } (m - 1)^2 = 0$$

$$\text{or } m = 1, 1$$

$$\therefore \text{C.F.} = (C_1 x + C_2) e^x$$

$$\begin{aligned}
 \text{and P.I.} &= \frac{1}{D^2 - 2D + 1} x^2 e^{3x} = e^{3x} \frac{1}{(D + 3)^2 - 2(D + 3) + 1} x^2 \\
 &= e^{3x} \frac{1}{D^2 + 4D + 4} x^2 = e^{3x} \frac{1}{4 \left( 1 + \frac{D}{2} \right)^2} x^2 \\
 &= \frac{e^{3x}}{4} \left( 1 + \frac{1}{2} D \right)^{-2} x^2 = \frac{e^{3x}}{4} \left[ \left( 1 - D + \frac{3}{4} D^2 + \dots \right) x^2 \right] \\
 &= \frac{1}{4} e^{3x} \left( x^2 - Dx^2 + \frac{3}{4} D^2 x^2 \right) \\
 &= \frac{1}{4} e^{3x} \left( x^2 - 2x + \frac{3}{2} \right) \\
 &= \frac{1}{8} e^{3x} (2x^2 - 4x + 3)
 \end{aligned}$$

∴ The required solution is  $y = \text{C.F} + \text{P.I}$

or  $y = (C_1 x + C_2) e^x + \frac{1}{8} e^{3x} (2x^2 - 4x + 3)$

**Example 28.** Solve  $(D^2 - 1) y = \cos hx$

**Solution.** The given equation can be written as

$$(D^2 - 1) y = \frac{1}{2} (e^x + e^{-x})$$

its auxiliary equation is  $m^2 - 1 = 0$

or  $m = \pm 1$

$\therefore$  C.F. =  $C_1 e^x + C_2 e^{-x}$ , where  $C_1, C_2$  are arbitrary constants

$$\begin{aligned} \text{and P.I} &= \frac{1}{D^2 - 1} \frac{1}{2} (e^x + e^{-x}) \\ &= \frac{1}{2} \frac{1}{(D^2 - 1)} e^x + \frac{1}{2} \frac{1}{D^2 - 1} e^{-x} \\ &= \frac{1}{2} e^x \frac{1}{(D + 1)^2 - 1} 1 + \frac{1}{2} \frac{1}{(-1)^2 - 1} e^{-x} \\ &= \frac{1}{2} e^x \frac{1}{D^2 + 2D} (1) - \frac{1}{4} e^{-x} \\ &= \frac{1}{2} e^x \frac{1}{2D \left(1 + \frac{D}{2}\right)} (1) - \frac{1}{4} e^{-x} \\ &= \frac{1}{4} e^x \frac{1}{D} \left(1 - \frac{D}{2} + \dots\dots\right) 1 - \frac{1}{4} e^{-x} \\ &= \frac{1}{4} e^x \frac{1}{D} (1) - \frac{1}{4} e^{-x} \\ &= \frac{1}{4} e^x \int 1 dx - \frac{1}{4} e^{-x} = \frac{1}{4} e^x x - \frac{1}{4} e^{-x} \end{aligned}$$

$\therefore$  The required solution is  $y = \text{C.F} + \text{P.I}$

or  $y = C_1 e^x + C_2 e^{-x} + \frac{1}{4} (x e^x - e^{-x})$

**Example 29.** Solve  $(D^2 - 4) y = \cos^2 x$

**Solution.** Its auxiliary equation is  $m^2 - 4 = 0$

or  $m = \pm 2$

∴ C.F. =  $C_1 e^{2x} + C_2 e^{-2x}$ , where  $C_1$  and  $C_2$  are arbitrary constants

$$\begin{aligned} \text{and P.I} &= \frac{1}{D^2 - 4} \cos^2 x = \frac{1}{D^2 - 4} \left[ \frac{1}{2} (1 + \cos 2x) \right] \\ &= \frac{1}{D^2 - 4} \frac{1}{2} + \frac{1}{D^2 - 4} \left( \frac{1}{2} \cos 2x \right) \\ &= -\frac{1}{4} \left( 1 - \frac{1}{4} D^2 \right)^{-1} \frac{1}{2} + \frac{1}{2} \frac{1}{-2^2 - 4} \cos 2x \\ &= -\frac{1}{4} \left[ 1 + \frac{1}{4} D^2 + \dots \right] \frac{1}{2} - \frac{1}{16} \cos 2x \\ &= -\frac{1}{4} \left( \frac{1}{2} \right) - \frac{1}{16} \cos 2x \\ &= -\frac{1}{16} (2 + \cos 2x) \end{aligned}$$

∴ The required solution is  $y = \text{C.F} + \text{P.I}$

$$\text{or } y = C_1 e^{2x} + C_2 e^{-2x} - \frac{1}{16} (2 + \cos 2x)$$

**Example 30.** Solve  $(D^2 + 1) y = x^2 \sin 2x$

**Solution.** Here the auxiliary equation is  $m^2 + 1 = 0$

or  $m = \pm i$

∴ C.F =  $C_1 \cos x + C_2 \sin x$ , where  $C_1$  and  $C_2$  are arbitrary constants

$$\text{and P.I} = \frac{1}{D^2 + 1} x^2 \sin 2x = \text{I.P of } \frac{1}{D^2 + 1} x^2 e^{2ix}$$

$$\begin{aligned} &= \text{I.P of } e^{2ix} \frac{1}{\{(D + 2i)^2 + 1\}} x^2 \\ &= \text{I.P of } e^{2ix} \frac{1}{(D^2 + 4iD - 3)} x^2 \\ &= \text{I.P of } e^{2ix} \frac{1}{-3 \left( 1 - \frac{4iD}{3} - \frac{D^2}{3} \right)} x^2 \\ &= \text{I.P. of } \frac{e^{2ix}}{-3} \left[ 1 - \left( \frac{4iD + D^2}{3} \right) \right]^{-1} x^2 \end{aligned}$$

$$\begin{aligned}
 &= \text{I.P of } \frac{e^{2ix}}{-3} \left[ 1 + \left( \frac{4iD + D^2}{3} \right) + \left( \frac{16 i^2 D^2}{9} + \dots \right) + \dots \right] x^2 \\
 &= \text{I.P of } -\frac{1}{3} e^{2ix} \left( 1 + \frac{4}{3} iD - \frac{13}{9} D^2 + \dots \right) x^2 \\
 &= \text{I.P of } -\frac{1}{3} e^{2ix} \left[ x^2 + \frac{4}{3} i (2x) - \frac{13}{9} (2) \right] \\
 &= \text{I.P of } -\frac{1}{3} (\cos 2x + i \sin 2x) \left[ \left( x^2 - \frac{26}{9} \right) + \frac{8}{3} xi \right] \\
 &= -\frac{1}{3} \left( \frac{8}{3} x \right) \cos 2x - \frac{1}{3} \left( x^2 - \frac{26}{9} \right) \sin 2x \\
 &= -\frac{1}{27} [24 x \cos 2x + (9x^2 - 26) \sin 2x]
 \end{aligned}$$

∴ The required solution is

$$y = C_1 \cos x + C_2 \sin x - \frac{1}{27} [24 x \cos 2x + (9x^2 - 26) \sin 2x]$$

### EXERCISE

**Solve the following differential Equations**

1.  $(D^3 + 6D^2 + 11D + 6)y = 0$

Ans.  $y = C_1 e^{-x} + C_2 e^{-2x} + C_3 e^{-3x}$

2.  $\frac{d^2x}{dt^2} - 3 \frac{dx}{dt} + 2x = 0$  given that when  $t = 0$ ,  $x = 0$  and  $\frac{dx}{dt} = 0$

Ans.  $x = 0$

3.  $(D^2 + D + 1)y = e^{-x}$ , where  $D \equiv \frac{d}{dx}$

Ans.  $y = e^{-x/2} \left[ C_1 \cos \left( \frac{1}{2} x\sqrt{3} \right) + C_2 \sin \left( \frac{1}{2} x\sqrt{3} \right) \right] + e^{-x}$

4.  $(D - 1)^2 (D^2 + 1)^2 y = e^x$

Ans.  $y = (C_1x + C_2) e^x + (C_3x + C_4) \cos x + (C_5x + C_6) \sin x + \frac{1}{8} x^2 e^x$

5.  $\frac{d^2y}{dx^2} - 3 \frac{dy}{dx} + 2y = e^x$ ,  $y = 3$  and  $\frac{dy}{dx} = 3$ , when  $x = 0$

Ans.  $y = 2e^x + e^{2x} - x e^x$

6.  $\frac{d^2y}{dx^2} - \frac{dy}{dx} - 2y = \sin 2x$

Ans.  $y = C_1 e^{2x} + C_2 e^{-x} + \frac{1}{20} (\cos 2x - 3 \sin 2x)$

7.  $(D^2 + 16)y = \sin 2x$ , given that  $y = 0$  and  $\frac{dy}{dx} = \frac{5}{6}$  where  $x = 0$

Ans.  $12y = 2 \sin 4x + \sin 2x$

8.  $\frac{d^2y}{dx^2} - 4y = e^x + \sin 2x$

Ans.  $y = C_1 e^{2x} + C_2 e^{-2x} - \frac{1}{3} e^x - \frac{1}{8} \sin 2x$

9.  $(4D^2 + 9)y = \sin x$ , given that  $y = \frac{1}{2}$ ,  $\frac{dy}{dx} = \frac{4}{5}$  when  $x = \pi$

Ans.  $y = \frac{2}{3} \cos \frac{3x}{2} - \frac{1}{2} \sin \frac{3x}{2} + \frac{1}{5} \sin x$

10. Find the integral of the equation  $\frac{d^2x}{dt^2} + 2n \cos x \frac{dx}{dt} + n^2 x = a \cos nt$   
which is such that when  $t = 0$ ,  $x = 0$  and  $\frac{dx}{dt} = 0$

Ans.  $x = e^{-nt \cos \alpha} \left\{ -\frac{a}{n^2 \sin 2\alpha} \sin (n \sin \alpha) t \right\} + \frac{a \sin nt}{2n^2 \cos \alpha}$

11. Solve  $\frac{d^2y}{dx^2} + a^2y = \sin ax$

Ans.  $y = C_1 \cos ax + C_2 \sin ax - \frac{x}{2a} \cos ax$

12.  $(D^2 + a^2)y = \cos ax$

Ans.  $y = C_1 \cos ax + C_2 \sin ax + \frac{1}{2} \left( \frac{x}{a} \right) \sin ax$

13.  $(D^4 - 1)y = \sin x$

Ans.  $y = C_1 e^x + C_2 e^{-x} + C_3 \cos x + C_4 \sin x + \frac{x}{4} \cos x$

Linear Differential Equations with Constant Coefficients and Applications

14.  $(D^4 + D^2 + 1)y = e^{-x/2} \cos\left(\frac{1}{2}x\sqrt{3}\right)$

Ans.  $y = e^{-x/2} \left[ C_1 \cos \frac{1}{2}x\sqrt{3} + C_2 \sin \frac{1}{2}x\sqrt{3} \right] + e^{x/2} \left[ C_3 \cos \frac{x}{2}\sqrt{3} + C_4 \sin \frac{x}{2}\sqrt{3} \right] - 16e^{-x/2} \cos\left(\frac{x}{2}\sqrt{3}\right)$

15.  $(D^2 + 1)y = \sin x \sin 2x$

Ans.  $y = C_1 \cos x + C_2 \sin x + \frac{x}{4} \sin x + \frac{1}{16} \cos 3x$

16.  $\frac{d^2y}{dx^2} + 4y = \sin^2 x$

Ans.  $y = C_1 \cos 2x + C_2 \sin 2x + \frac{1}{8} - \frac{x}{8} \sin 2x$

17.  $(D^2 - 4)y = x^2$

Ans.  $y = C_1 e^{2x} + C_2 e^{-2x} - \frac{1}{4} \left( x^2 + \frac{1}{2} \right)$

18. Solve  $(D^3 - 8)y = x^3$

Ans.  $y = C_1 e^{2x} + e^{-x} (C_2 \cos x\sqrt{3} + C_3 \sin x\sqrt{3}) - \frac{1}{8} \left( x^3 + \frac{3}{4} \right)$

19.  $(D^3 + 2D^2 + D)y = e^{2x} + x^2 + x$

Ans.  $y = C_1 + (C_2x + C_3)e^{-x} + \frac{1}{18}e^{2x} + \frac{1}{3}x^3 - \frac{3}{2}x^2 + 4x$

20.  $\frac{d^2y}{dx^2} = a + bx + cx^2$ , given that  $\frac{dy}{dx} = 0$ , when  $x = 0$  and  $y = d$ , when  $x = 0$

Ans.  $y = d + \frac{ax^2}{2} + \frac{bx^3}{6} + \frac{cx^4}{12}$

21.  $(D^2 + 4D - 12)y = (x - 1)e^{2x}$

Ans.  $y = C_1 e^{2x} + C_2 e^{-6x} + \frac{1}{64} (4x^2 - 9x) e^{2x}$

22.  $\left( \frac{d^2y}{dx^2} - 2 \frac{dy}{dx} + 5 \right) y = e^{2x} \sin x$



Ans.  $y = e^x (C_1 \cos 2x + C_2 \sin 2x) - \frac{1}{10} e^{2x} (\cos x - 2 \sin x)$

23.  $(D^2 - 4D + 4) y = e^{2x} \sin 3x$

Ans.  $y = (C_1 x + C_2) e^{2x} - \frac{1}{9} e^{2x} \sin 3x$

24.  $\left(\frac{d}{dx} + 1\right)^3 y = x^2 e^{-x}$

Ans.  $y = (C_1 x^2 + C_2 x + C_3) e^{-x} + \frac{1}{60} x^5 e^{-x}$

25.  $\frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + y = x \cos x$

Ans.  $y = (C_1 x + C_2) e^{-x} + \frac{1}{2} x \sin x - \frac{1}{2} \sin x + \frac{1}{2} \cos x$

26.  $\frac{d^4 y}{dx^4} - y = x \sin x$

Ans.  $y = C_1 e^x + C_2 e^{-x} + C_3 \cos x + C_4 \sin x + \frac{1}{8} (x^2 \cos x - 3 x \sin x)$

27.  $\frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + y = x \sin x$

**(I.A.S. 1998)**

Ans.  $y = (C_1 + C_2 x) e^{-x} + \frac{1}{2} (\sin x + \cos x - x \cos x)$

28. Solve  $(D^2 + 1)^2 y = 24 x \cos x$ , given that  $y = Dy = D^2 y = 0$  and  $D^3 y = 12$  when  $x = 0$

Ans.  $y = 3x^2 \sin x - x^3 \cos x$

29.  $(D^2 - 4D + 4) y = 8x^2 e^{2x} \sin 2x$

**(U.P.T.U. 2004, 2005)**

Ans.  $y = (C_1 + C_2 x) e^{2x} + e^{2x} (-2x^2 \sin 2x - 4x \cos 2x + 3 \sin 2x)$

## Objective Type of Questions

Choose a correct answer from the four answers given in each of the following questions.

1. The solution of the differential equation  $\frac{d^2y}{dx^2} + (3i - 1) \frac{dy}{dx} - 3iy = 0$  is

(I.A.S. 1998)

- (a)  $y = C_1 e^x + C_2 e^{3ix}$       (b)  $y = C_1 e^{-x} + C_2 e^{3ix}$   
(c)  $y = C_1 e^x + C_2 e^{-3ix}$       (d)  $y = C_1 e^{-x} + C_2 e^{-3ix}$

Ans. (c)

2. A particular integral of the differential equation  $(D^2 + 4)y = x$  is

(I.A.S. 1998)

- (a)  $xe^{-2x}$       (b)  $x \cos 2x$   
(c)  $x \sin 2x$       (d)  $x/4$

Ans. (d)

3. The particular integral of  $(D^2 + 1)y = e^{-x}$  is

(I.A.S. 1999)

- (a)  $\left(\frac{1}{4} - \frac{x}{2}\right)e^{-x}$       (b)  $\left(\frac{1}{4} + \frac{x}{2}\right)e^{-x}$   
(c)  $\frac{e^{-x}}{2}$       (d)  $\frac{e^{-x}}{-2}$

Ans. (c)

4. For the differential equation  $(D + 2)(D - 1)^3 y = e^x$  the particular integral is

(I.A.S. 1990, U.P.P.C.S. 2000)

- (a)  $\frac{1}{18} x^4 e^x$       (b)  $\frac{1}{18} x^3 e^x$   
(c)  $\frac{1}{18} x e^{3x}$       (d)  $\frac{1}{36} x e^{3x}$

Ans. (b)

5. The particular integral of the differential equation  $\frac{d^2y}{dx^2} + 9y = \sin 3x$  is

- (a)  $\frac{x \sin 3x}{18}$                       (b)  $\frac{x \sin 3x}{6}$   
(c)  $\frac{-x \cos 3x}{6}$                       (d)  $\frac{x \cos 3x}{18}$

**Ans. (c)**

6. The solution of the differential equation  $\frac{d^2y}{dx^2} - 3 \frac{dy}{dx} - 4y = 0$  is

(U.P.P.C.S. 2001)

- (a)  $y = C_1 e^{-x} + C_2 e^{-4x}$                       (b)  $y = C_1 e^x + C_2 e^{4x}$   
(c)  $y = C_1 e^{-x} + C_2 e^{4x}$                       (d)  $y = C_1 e^x + C_2 e^{-4x}$

**Ans. (c)**

7. The general solution of the differential equation  $(D^2 - 1)y = x^2$  is

- (a)  $y = C_1 e^x + C_2 e^{-x} - x^2$                       (b)  $y = C_1 e^x + C_2 e^{-x} + (x^2 + 2)$   
(c)  $y = C_1 e^x + C_2 e^{-x} - 2$                       (d)  $y = C_1 e^x + C_2 e^{-x} - (x^2 + 2)$

**Ans. (d)**

8. The P.I. of  $(D^2 - 2D)y = e^x \sin x$  is

- (a)  $-\frac{1}{2} e^x \sin x$                       (b)  $e^x \cos x$   
(c)  $-\frac{1}{2} e^x \cos x$                       (d) none of these

**Ans. (a)**

9. The general solution of the differential equation  $D^2(D + 1)^2 y = e^x$  is

(I.A.S. 1990)

- (a)  $y = C_1 + C_2 x + (C_3 + C_4 x) e^x$   
(b)  $y = C_1 + C_2 x + (C_3 + C_4 x) e^{-x} + \frac{1}{4} e^x$   
(c)  $y = (C_1 + C_2 e^{-x}) + (C_3 + C_4 x) e^{-x} + \frac{1}{4} e^x$   
(d) none of these

**Ans. (b)**

10.  $y = e^{-x} (C_1 \cos \sqrt{3} x + C_2 \sin \sqrt{3} x) + C_3 e^{2x}$  is the solution of

(I.A.S. 1994)

Linear Differential Equations with Constant Coefficients and Applications

- (a)  $\frac{d^3y}{dx^3} + 4y = 0$                       (b)  $\frac{d^3y}{dx^3} + 8y = 0$   
(c)  $\frac{d^3y}{dx^3} - 8y = 0$                       (d)  $\frac{d^3y}{dx^3} - 2\frac{d^2y}{dx^2} + \frac{dy}{dx} - 2 = 0$

**Ans. (c)**

11. The P.I of the differential equation  $(D^3 - D)y = e^x + e^{-x}$  is

**(I.A.S. 1993)**

- (a)  $\frac{1}{2}(e^x + e^{-x})$                       (b)  $\frac{1}{2}x(e^x + e^{-x})$   
(c)  $\frac{1}{2}x^2(e^x + e^{-x})$                       (d)  $\frac{1}{2}x^2(e^x - e^{-x})$

**Ans. (b)**

12. Given  $y = 1 + \cos x$  and  $y = 1 + \sin x$  are solutions of the differential equation  $\frac{d^2y}{dx^2} + y = 1$ , its solution will be also

**(R.A.S. 1994)**

- (a)  $y = 2(1 + \cos x)$                       (b)  $y = 2 + \cos x + \sin x$   
(c)  $y = \cos x - \sin x$                       (d)  $y = 1 + \cos x + \sin x$

**Ans. (d)**

13. The solution of the differential equation  $(D^3 - 6D^2 + 11D - 6)y = 0$  is

**(R.A.S. 1994)**

- (a)  $y = C_1 e^x + C_2 e^{2x} + C_3 e^{4x}$                       (b)  $y = C_1 e^{2x} + C_2 e^{3x} + C_3 e^{4x}$   
(c)  $y = C_1 e^{-x} + C_2 e^{2x} + C_3 e^{4x}$                       (d)  $y = C_1 e^x + C_2 e^{2x} + C_3 e^{3x}$

**Ans. (d)**

14. The solution of the differential equation  $\frac{d^2y}{dx^2} - \frac{dy}{dx} - 2y = 3e^{2x}$ , when  $y(0) = 0$  and  $y'(0) = -2$  is

**(R.A.S. 1994)**

- (a)  $y = e^{-x} - e^{2x} + x e^{2x}$                       (b)  $y = e^x - e^{-2x} - x e^{2x}$   
(c)  $y = e^{-x} + e^{2x} - \frac{x}{2} e^{2x}$                       (d)  $y = e^x - e^{-2x} - \frac{x}{2} e^{2x}$

**Ans. (a)**

15. The solution of the differential equation  $\frac{d^2y}{dx^2} + y = \cos 2x$  is

(U.P.P.C.S. 1995)

(a)  $A \cos x + B \sin x + \frac{1}{3} \cos 2x$       (b)  $A \cos x + B \sin x + \frac{1}{3} \sin 2x$

(c)  $A \cos x + B \sin x - \frac{1}{3} \cos 2x$       (d)  $A \cos x + B \sin x - \frac{1}{3} \sin 2x$

**Ans. (c)**

16. The general solution of the differential equation  $\frac{d^2y}{dx^2} + n^2 y = 0$  is

(R.A.S. 1995)

(a)  $C_1 \sqrt{\cos nx} + C_2 \sqrt{\sin nx}$       (b)  $C_1 \cos nx + C_2 \sin nx$

(c)  $C_1 \cos^2 nx + C_2 \sin^2 nx$       (d)  $C_1 \cos^3 nx + C_2 \sin^3 nx$

**Ans. (b)**

17. The particular integral of  $\frac{d^2y}{dx^2} + \frac{dy}{dx} = x^2 + 2x + 4$  is

(I.A.S. 1996)

(a)  $\frac{x^2}{3} + 4x$       (b)  $\frac{x^3}{3} + 4x$

(c)  $\frac{x^3}{3} + 4$       (d)  $\frac{x^3}{3} + 4x^2$

**Ans. (b)**

18. The general solution of the differential equation

$$\frac{d^4y}{dx^4} - 6 \frac{d^3y}{dx^3} + 12 \frac{d^2y}{dx^2} - 8 \frac{dy}{dx} = 0 \text{ is}$$

(I.A.S. 1996)

(a)  $y = C_1 + (C_2 + C_3 x + C_4 x^2) e^{2x}$       (b)  $y = (C_1 + C_2 x + C_3 x^2) e^{2x}$

(c)  $y = C_1 + C_2 x + C_3 x^2 + C_4 x^3$       (d)  $y = C_1 x + C_2 x^2 + C_3 x^3 + C_4 x^4$

**Ans. (a)**

19. The primitive of the differential equation  $(D^2 - 2D + 5)^2 y = 0$  is

(I.A.S. 1995)

Linear Differential Equations with Constant Coefficients and Applications

- (a)  $e^x \{(C_1 + C_2 x) \cos 2x + (C_3 + C_4 x) \sin 2x\}$   
(b)  $e^{2x} \{(C_1 + C_2 x) \cos x + (C_3 + C_4 x) \sin x\}$   
(c)  $(C_1 e^x + C_2 e^{2x}) \cos x + (C_3 e^x + C_4 e^{2x}) \sin x$   
(d)  $e^x \{C_1 \cos x + C_2 \cos 2x + C_3 \sin x + C_4 \sin 2x\}$

**Ans. (a)**

20. Which one of the following does not satisfy the differential equation

$$\frac{d^3 y}{dx^3} - y = 0?$$

(U.P.P.C.S. 1994)

- (a)  $e^x$  (b)  $e^{-x}$   
(c)  $e^{-x/2} \sin \left( \frac{\sqrt{3}}{2} x \right)$  (d)  $e^{-x/2} \cos \left( \frac{\sqrt{3}}{2} x \right)$

**Ans. (b)**

21. The particular integral of  $(D^2 + a^2) y = \sin ax$  is

(I.A.S. 1995)

- (a)  $-\frac{x}{2a} \cos ax$  (b)  $\frac{x}{2a} \cos ax$   
(c)  $-\frac{ax}{2} \cos ax$  (d)  $\frac{ax}{2} \cos ax$

**Ans. (a)**

# Chapter 4

## Equations Reducible To Linear Equations with Constant Coefficients

---

### INTRODUCTION

Now we shall study two such forms of linear differential equations with variable coefficients which can be reduced to linear differential equations with constant coefficients by suitable substitutions.

#### 1. Cauchy's Homogeneous Linear Equations

A differential equation of the form

$$x^n \frac{d^n y}{dx^n} + P_1 x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + P_n y = X$$

where  $P_1, P_2, \dots, P_n$  are constants and  $X$  is either a function of  $x$  or a constant is called Cauchy-Euler homogeneous linear differential equation.

The solution of the above homogeneous linear equation may be obtained after transforming it into linear equation with constant coefficients by using the substitution.

By the substitution  $x = e^z$  or  $z = \log_e x$ ;  $\therefore \frac{dz}{dx} = \frac{1}{x}$

$$\text{Now } \frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx} = \frac{1}{x} \frac{dy}{dz}$$

$$\therefore x \frac{dy}{dx} = \frac{dy}{dz} \tag{1}$$

$$\text{Again } \frac{d^2 y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d}{dx} \left( \frac{1}{x} \frac{dy}{dz} \right)$$

$$= \frac{x \frac{d^2 y}{dz^2} \frac{dz}{dx} - \frac{dy}{dz}}{x^2} = \frac{x \frac{d^2 y}{dz^2} \frac{1}{x} - \frac{dy}{dz}}{x^2}$$

$$\therefore x^2 \frac{d^2y}{dx^2} = \frac{d^2y}{dz^2} - \frac{dy}{dz} \tag{2}$$

Also 
$$\frac{d^3y}{dx^3} = \frac{d}{dx} \left( \frac{d^2y}{dx^2} \right) = \frac{d}{dx} \left[ \frac{1}{x^2} \left( \frac{d^2y}{dz^2} - \frac{dy}{dz} \right) \right]$$

$$= \frac{x^2 \left[ \frac{d^3y}{dz^3} \frac{dz}{dx} - \frac{d^2y}{dz^2} \frac{dz}{dx} \right] - 2x \left( \frac{d^2y}{dz^2} - \frac{dy}{dz} \right)}{x^4}$$

Substituting  $\frac{dz}{dx} = \frac{1}{x}$  and simplifying, we get

$$x^3 \frac{d^3y}{dx^3} = \frac{d^3y}{dz^3} - 3 \frac{d^2y}{dz^2} + 2 \frac{dy}{dz} \tag{3}$$

Using  $x \frac{d}{dx} \equiv \frac{d}{dz} \equiv D$ , in (1), (2) and (3)

we get 
$$x \frac{dy}{dx} = Dy$$

$$x^2 \frac{d^2y}{dx^2} = D(D-1)y$$

$$x^3 \frac{d^3y}{dx^3} = D(D-1)(D-2)y$$

In general, we have

$$x^n \frac{d^n y}{dx^n} = D(D-1)(D-2)\dots\dots\dots(D-n+1)y$$

Using these results in homogeneous linear equation, it will be transformed into a linear differential equation with constant coefficients.

**Example 1.** Solve  $x^2 \frac{d^2y}{dx^2} - x \frac{dy}{dx} - 3y = x^2 \log x$  (I.A.S. 2001)

**Solution.** On changing the independent variable by substituting

$$x = e^z \text{ or } z = \log_e x \text{ and } \frac{d}{dz} \equiv D$$



Equations Reducible To Linear Equations with Constant Coefficients

The differential equation becomes

$$[D(D-1) - D - 3]y = ze^{2z}$$

or  $(D^2 - 2D - 3)y = ze^{2z}$

Now the auxiliary equation is  $m^2 - 2m - 3 = 0$

$\Rightarrow m = 3, -1$

Hence, the C.F. =  $C_1 e^{3z} + C_2 e^{-z} = C_1 x^3 + \frac{C_2}{x}$

and P.I. =  $\frac{1}{D^2 - 2D - 3} ze^{2z}$

$$\begin{aligned} &= e^{2z} \frac{1}{(D+2)^2 - 2(D+2) - 3} z = e^{2z} \frac{1}{D^2 + 4 + 4D - 2D - 4 - 3} z \\ &= e^{2z} \frac{1}{D^2 + 2D - 3} z \\ &= e^{2z} \frac{1}{-3 \left( 1 - \frac{2D}{3} - \frac{D^2}{3} \right)} z \\ &= \frac{e^{2z}}{-3} \left[ 1 - \left( \frac{2D}{3} + \frac{D^2}{3} \right) \right]^{-1} z \\ &= \frac{e^{2z}}{-3} \left[ 1 + \frac{2D}{3} + \frac{D^2}{3} + \dots \right] z \\ &= \frac{e^{2z}}{-3} \left( z + \frac{2}{3} \right) = e^{2z} \left( -\frac{z}{3} - \frac{2}{9} \right) \\ &= x^2 \left( -\frac{1}{3} \log_e x - \frac{2}{9} \right) \end{aligned}$$

Hence solution of the given differential equation is

$$y = C_1 x^3 + \frac{C_2}{x} + x^2 \left( -\frac{1}{3} \log_e x - \frac{2}{9} \right)$$

**Example 2.** Solve  $x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} - 4y = x^4$

**Solution.** On changing the independent variable by substituting  $x = e^z$  or  $z = \log_e x$  and  $\frac{d}{dz} \equiv D$

The differential equation becomes

$$[D(D - 1) - 2D - 4] y = e^{4z}$$

Now, the auxiliary equation is

$$m^2 - 3m - 4 = 0 \text{ or } m = 4, -1$$

$$\therefore \text{C.F.} = C_1 e^{4z} + C_2 e^{-z}$$

$$= C_1 x^4 + C_2 \frac{1}{x} \quad \because e^z = x$$

and

$$\begin{aligned} \text{P.I.} &= \frac{1}{D^2 - 3D - 4} e^{4z} \\ &= e^{4z} \frac{1}{(D + 4)^2 - 3(D + 4) - 4} 1 \\ &= e^{4z} \frac{1}{D^2 + 16 + 8D - 3D - 12 - 4} 1 \\ &= e^{4z} \frac{1}{D^2 + 5D} 1 \\ &= e^{4z} \frac{1}{5D \left(1 + \frac{D}{5}\right)} 1 = e^{4z} \frac{1}{5D} \left(1 + \frac{D}{5}\right)^{-1} 1 \\ &= e^{4z} \frac{1}{5D} \left[1 - \frac{D}{5} + \dots\dots\dots\right] 1 \\ &= e^{4z} \frac{1}{5D} 1 = e^{4z} \frac{1}{5} z = \frac{1}{5} z e^{4z} \\ &= \frac{1}{5} x^4 \log_e x \end{aligned}$$

Hence the required solution is

$$y = C_1 x^4 + C_2 \frac{1}{x} + \frac{1}{5} x^4 \log_e x$$

Equations Reducible To Linear Equations with Constant Coefficients

**Example 3.** Solve  $x^3 \frac{d^3y}{dx^3} + 2x^2 \frac{d^2y}{dx^2} + 2y = 10 \left( x + \frac{1}{x} \right)$

(I.A.S. 2006, 1998, U.P.P.C.S. 1973)

**Solution.** On changing the independent variable by substituting  $x = e^z$  or  $z = \log_e x$  and  $\frac{d}{dz} \equiv D$  the given differential equation becomes

$$[D(D-1)(D-2) + 2D(D-1) + 2]y = 10(e^z + e^{-z})$$

or  $(D^3 - D^2 + 2)y = 10(e^z + e^{-z})$

The auxiliary equation is

$$m^3 - m^2 + 2 = 0 \text{ or } (m+1)(m^2 - 2m + 2) = 0$$

or  $D = -1, 1 \pm i$

$$\begin{aligned} \therefore \text{C.F} &= C_1 e^{-z} + C_2 e^z \cos(z + C_3) \\ &= C_1 x^{-1} + C_2 x \cos(\log_e x + C_3) \end{aligned}$$

and P.I. =  $\frac{1}{(D+1)(D^2-2D+2)} 10e^z + \frac{1}{(D+1)(D^2-2D+2)} 10e^{-z}$

$$= \frac{1}{(1+1)(1^2-2.1+2)} 10e^z + e^{-z} \frac{1}{\{(D-1+1)\}\{(D-1)^2-2(D-1)+2\}} 10$$

$$= \frac{1}{2} 10e^z + e^{-z} \frac{1}{D(D^2+1-2D-2D+2+2)} 10$$

$$= 5e^z + e^{-z} \frac{1}{D(D^2-4D+5)} 10$$

$$= 5e^z + e^{-z} \frac{1}{5D \left( 1 - \frac{4D}{5} + \frac{D^2}{5} \right)} 10$$

$$= 5e^z + e^{-z} \frac{1}{5D} \left\{ 1 - \left( \frac{4D}{5} - \frac{D^2}{5} \right) \right\}^{-1} 10$$

$$= 5e^z + e^{-z} \frac{1}{5D} 10$$

$$\begin{aligned} &= 5e^z + 2e^{-z} \frac{1}{D} 1 = 5e^z + 2e^{-z} z \\ &= 5e^z + 2z e^{-z} \\ &= 5x + 2 (\log_e x) \frac{1}{x} \end{aligned}$$

Hence the required solution is

$$y = C_1 x^{-1} + C_2 x (\log_e x + C_3) + 5x + (2 \log_e x) \frac{1}{x}$$

**Example 4.** Solve  $x^3 \frac{d^3 y}{dx^3} + 3x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + y = x + \log x$

(Bihar P.C.S. 2002, U.P.T.U. 2001)

**Solution.** On changing the independent variable by substituting

$$x = e^z \text{ or } z = \log_e x \text{ and } \frac{d}{dz} \equiv D, \text{ we have}$$

$$[D(D-1)(D-2) + 3D(D-1) + D + 1] y = e^z + z$$

$$\text{or } (D^3 + 1) y = e^z + z$$

The auxiliary equation is  $m^3 + 1 = 0$

$$\text{or } (m + 1)(m^2 - m + 1) = 0$$

$$\Rightarrow m = -1, \frac{1 \pm \sqrt{3} i}{2}$$

$$\text{so C.F.} = C_1 e^{-z} + e^{z/2} \left( C_2 \cos \frac{\sqrt{3}}{2} z + C_3 \sin \frac{\sqrt{3}}{2} z \right)$$

$$\text{and P.I.} = \frac{1}{D^3 + 1} (e^z + z)$$

Equations Reducible To Linear Equations with Constant Coefficients

$$\begin{aligned} &= \frac{1}{D^3 + 1} e^z + \frac{1}{1 + D^3} z \\ &= \frac{1}{1^3 + 1} e^z + (1 + D^3)^{-1} (z) \\ &= \frac{1}{2} e^z + (1 - D^3 + \dots\dots) z \\ &= \frac{e^z}{2} + z \end{aligned}$$

Therefore required solution is

$$y = C_1 e^z + e^{z/2} \left( C_2 \cos \frac{\sqrt{3}}{2} z + C_3 \sin \frac{\sqrt{3}}{2} z \right) + \frac{e^z}{2} + z$$

or 
$$y = C_1 x^{-1} + \sqrt{x} \left[ C_2 \cos \frac{\sqrt{3}}{2} (\log x) + C_3 \sin \frac{\sqrt{3}}{2} (\log x) \right] + \frac{x}{2} + \log x$$

**Example 5.** Solve  $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} + y = (\log x) \sin (\log x)$

(U.P.T.U. 2002)

**Solution.** On changing the independent variable by substituting  $x = e^z$  or  $z = \log_e x$  and  $\frac{d}{dz} = D$ , we have

$$[D(D - 1) + D + 1] y = z \sin z$$

or  $(D^2 + 1) y = z \sin z$

The auxiliary equation is

$$m^2 + 1 = 0$$

or  $m = \pm i$

Thus C.F. =  $C_1 \cos z + C_2 \sin z$   
 $= C_1 \cos (\log x) + C_2 \sin (\log x)$

& P.I. =  $\frac{1}{D^2 + 1} z \sin z$   
 $= \text{imaginary part of } \frac{1}{D^2 + 1} z e^{iz}$

$$\begin{aligned}
 &= \text{I.P. of } e^{iz} \frac{1}{(D+i)^2 + 1} z \\
 &= \text{I.P. of } e^{iz} \frac{1}{D^2 + 2iD - 1 + 1} z \\
 &= \text{I.P. of } e^{iz} \frac{1}{2iD} \left(1 + \frac{D}{2i}\right)^{-1} z \\
 &= \text{I.P. of } e^{iz} \frac{1}{2iD} \left(1 - \frac{D}{2i} + \dots\right) z \\
 &= \text{I.P. of } e^{iz} \frac{1}{2iD} \left(z - \frac{1}{2i}\right) \\
 &= \text{I.P. of } e^{iz} \frac{1}{2i} \int \left(z + \frac{i}{2}\right) dz \\
 &= \text{I.P. of } \frac{e^{iz}}{2i} \left(\frac{z^2}{2} + \frac{iz}{2}\right) \\
 &= \text{I.P. of } \frac{-i}{2} (\cos z + i \sin z) \left(\frac{z^2}{2} + \frac{i}{2} z\right) \\
 &= \text{I.P. of } -\frac{1}{2} (i \cos z - \sin z) \left(\frac{z^2}{2} + \frac{i}{2} z\right) \\
 &= -\frac{z^2}{4} \cos z + \frac{1}{4} z \sin z = \frac{z}{4} (\sin z - z \cos z) \\
 &= \frac{\log x}{4} [\sin (\log x) - \log x \cos (\log x)]
 \end{aligned}$$

Hence required solution is

$$y = C_1 \cos (\log x) + C_2 \sin (\log x) + \frac{\log x}{4} [\sin (\log x) - \log x \cos (\log x)]$$

**Example 6.** Solve  $x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + 4y = \cos (\log x) + x \sin (\log x)$

**Solution.** On changing the independent variable by substituting  $x = e^z$  or  $z = \log x$  and  $\frac{d}{dz} \equiv D$  we have

$$[D(D-1) - D + 4] y = \cos z + e^z \sin z$$

Equations Reducible To Linear Equations with Constant Coefficients

or  $(D^2 - 2D + 4) y = \cos z + e^z \sin z$

The auxiliary equation is

$$m^2 - 2m + 4 = 0$$

$$\Rightarrow m = 1 \pm i\sqrt{3}$$

Therefore C.F. =  $e^z (C_1 \cos \sqrt{3} z + C_2 \sin \sqrt{3} z)$

$$= x \left[ C_1 \cos \left\{ \sqrt{3} \log x + C_2 \sin \left( \sqrt{3} \log x \right) \right\} \right]$$

and

$$\begin{aligned} \text{P.I} &= \frac{1}{D^2 - 2D + 4} \cos z + \frac{1}{D^2 - 2D + 4} e^z \sin z \\ &= \frac{1}{-1^2 - 2D + 4} \cos z + e^z \frac{1}{(D + 1)^2 - 2(D + 1) + 4} \sin z \\ &= \frac{1}{3 - 2D} \cos z + e^z \frac{1}{D^2 + 3} \sin z \\ &= \frac{3 + 2D}{9 - 4D^2} \cos z + e^z \frac{1}{-1^2 + 3} \sin z \\ &= \frac{(3 + 2D)}{9 - 4(-1)^2} \cos z + \frac{e^z}{2} \sin z \\ &= \frac{1}{13} (3 + 2D) \cos z + \frac{1}{2} e^z \sin z \\ &= \frac{1}{13} (3 \cos z - 2 \sin z) + \frac{1}{2} e^z \sin z \\ &= \frac{1}{13} [3 \cos (\log x) - 2 \sin (\log x)] + \frac{1}{2} x \sin (\log x) \end{aligned}$$

Therefore required solution is

$$y = x \left[ C_1 \cos (\sqrt{3} \log x) + C_2 \sin (\sqrt{3} \log x) \right] + \frac{1}{13} [3 \cos (\log x) - 2 \sin (\log x)] + \frac{1}{2} x \sin (\log x)$$

**Example 7.** Solve  $x^2 \frac{d^2y}{dx^2} + 4x \frac{dy}{dx} + 2y = e^x$

[U.P.T.U. (CO) 2005]

**Solution.** Substituting  $x = e^z$  or  $z = \log_e x$  and putting  $\frac{d}{dz} \equiv D$ , we have

$$[D(D - 1) + 4D - 2] y = e^{e^z}$$

or  $(D + 2)(D + 1)y = e^{e^z}$

The auxiliary equation is  $(m + 2)(m + 1) = 0$

$\Rightarrow m = -2, -1$

$\therefore$  C.F. =  $C_1 e^{-2z} + C_2 e^{-z} = C_1 x^{-2} + C_2 x^{-1}$

and P.I. =  $\frac{1}{(D + 2)(D + 1)} e^{e^z} = \left[ \frac{1}{D + 1} - \frac{1}{D + 2} \right] e^{e^z}$

Let  $\frac{1}{D + 1} e^{e^z} = u \quad \therefore (D + 1)u = e^{e^z}$

or  $\frac{du}{dz} + u = e^{e^z}$ , which is linear

Integrating factor =  $e^z$ , Hence its solution is

$$\begin{aligned} u e^z &= \int e^z e^{e^z} dz \\ &= \int e^x dx \quad \because e^z = x \therefore e^z dz = dx \\ &= e^x \end{aligned}$$

or  $u = e^x \frac{1}{e^z} = \frac{1}{x} e^x \quad \because e^z = x$

Further, let  $\frac{1}{D + 2} e^{e^z} = v$

$\therefore (D + 2)v = e^{e^z}$

or  $\frac{dv}{dz} + 2v = e^{e^z}$ , which is linear

Integrating factor =  $e^{2z}$ , Hence its solution is

$$\begin{aligned} v e^{2z} &= \int e^{2z} e^{e^z} dz \\ &= \int e^z e^{e^z} e^z dz \\ &= \int x e^x dx \quad \because e^z = x, \therefore e^z dz = dx \\ &= e^x (x - 1) \end{aligned}$$



Equations Reducible To Linear Equations with Constant Coefficients

$$\therefore v = \frac{e^x (x-1)}{e^{2x}} = \frac{e^x (x-1)}{x^2} = \frac{e^x}{x} - \frac{e^x}{x^2}$$

Hence P.I. =  $u - v = \frac{1}{x} e^x - \left( \frac{e^x}{x} - \frac{e^x}{x^2} \right) = \frac{e^x}{x^2}$

Hence the required solution is

$$y = C_1 x^{-2} + C_2 x^{-1} + \frac{e^x}{x^2}$$

**2. Legendre's linear differential equation  
(Equation reducible to homogeneous form)**

An equation of the form

$$(ax + b)^n \frac{d^n y}{dx^n} + k_1 (ax + b)^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + k_n y = X \quad (1)$$

Where  $a, b, k_1, k_2, \dots, k_n$  are all constants and  $X$  is a function of  $x$ , is called Legendre's linear equation.

Such equations can be reduced to linear equations with constant coefficients by substituting  $ax + b = e^z$  i.e.  $z = \log(ax + b)$

Then if  $D = \frac{d}{dz}$ ,  $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx} = \frac{a}{ax + b} \frac{dy}{dz}$

i.e.  $(ax + b) \frac{dy}{dx} = aDy$

$$\begin{aligned} \frac{d^2 y}{dx^2} &= \frac{d}{dx} \left( \frac{a}{ax + b} \frac{dy}{dz} \right) = \frac{-a^2}{(ax + b)^2} \frac{dy}{dz} + \frac{a}{ax + b} \frac{d}{dz} \left( \frac{dy}{dz} \right) \frac{dz}{dx} \\ &= \frac{a^2}{(ax + b)^2} \left( \frac{d^2 y}{dz^2} - \frac{dy}{dz} \right) \end{aligned}$$

i.e.  $(ax + b)^2 \frac{d^2 y}{dx^2} = a^2 D(D - 1)y$

Similarly  $(ax + b)^3 \frac{d^3 y}{dx^3} = a^3 D(D - 1)(D - 2)y$  and so on.

After making these replacements in (1), there results a linear equation with constant coefficients.

**Example 8.** Solve  $(1+x)^2 \frac{d^2y}{dx^2} + (1+x) \frac{dy}{dx} + y = 4 \cos \log(1+x)$

**Solution.** put  $1+x = e^z$  and  $\frac{d}{dz} = D$

Hence the given differential equation becomes

$$[D(D-1) + D + 1]y = 4 \cos z$$

$\therefore$  Auxiliary equation is

$$D^2 + 1 = 0 \text{ or } D = \pm i$$

$\therefore$  C.F. =  $C_1 \cos(z + C_2) = C_1 \cos[\log(1+x) + C_2]$

$$\text{and P.I} = \frac{1}{D^2 + 1} 4 \cos z = 4 \cdot \frac{z}{2} \sin z$$

$$= 2z \sin z$$

$$= 2 \log(1+x) \sin \log(1+x)$$

Hence the required solution is

$$y = C_1 \cos[\log(1+x) + C_2] + 2 \log(1+x) \sin \log(1+x)$$

**Example 9 :** Solve

$$(1+2x)^2 \frac{d^2y}{dx^2} - 6(1+2x) \frac{dy}{dx} + 16y = 8(1+2x)^2$$

and  $y(0) = 0, y'(0) = 2$  (I.A.S. 1997)

**Solution :** Let  $1+2x = z$  then

$$\frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx} = 2 \frac{dy}{dz}$$
$$\therefore \frac{dz}{dx} = 2$$

$$\text{and } \frac{d^2y}{dx^2} = \frac{d}{dz} \left( 2 \frac{dy}{dz} \right) \frac{dz}{dx} = 4 \frac{d^2y}{dz^2}$$

Substituting these in given differential equation we have

Equations Reducible To Linear Equations with Constant Coefficients

$$4z^2 \frac{d^2y}{dz^2} - 6 \cdot 2z \frac{dy}{dz} + 16y = 8z^2$$

$$\text{or } z^2 \frac{d^2y}{dz^2} - 3z \frac{dy}{dz} + 4y = 2z^2$$

putting  $z = e^t$ , we have

$$\{\theta(\theta-1)-2\theta+4\} y = 2e^{2t}$$

$$\text{or } (\theta^2-4\theta+4) y = 2e^{2t}$$

its auxiliary equation is

$$m^2 - 4m + 4 = 0$$

$$\text{i.e. } (m - 2)^2 = 0$$

$$\text{or } m = 2 \text{ (twice)}$$

$$\therefore \text{C.F.} = (C_1 + C_2t)e^{2t}$$

$$= (C_1 + C_2 \log z) z^2$$

$$= \{C_1 + C_2 \log(1+2x)\} (1+2x)^2$$

$$\text{and P.I.} = \frac{1}{\theta^2 - 4\theta + 4} 2e^{2t} = 2e^{2t} \frac{1}{(\theta + 2)^2 - 4(\theta + 2) + 4} \cdot 1$$

$$= 2e^{2t} \frac{1}{\theta^2 + 4\theta - 4 - 4\theta - 8 + 4} \cdot 1 = 2e^{2t} \frac{1}{\theta^2} \cdot 1$$

$$= 2e^{2t} \frac{t^2}{2} = z^2 (\log z)^2$$

$$= (1+2x)^2 \{\log(1+2x)\}^2$$

Hence the complete solution is

$$y = \{C_1 + C_2 \log(1+2x)\} (1+2x)^2 + (1+2x)^2 \{\log(1+2x)\}^2$$

### METHOD OF VARIATION OF PARAMETERS

Method of variation of parameters enables to find solution of any linear non homogeneous differential equation of second order even (with variable coefficients also) provided its complimentary function is given (known). The particular integral of the non-homogeneous equation is obtained by varying the parameters i.e. by replacing the arbitrary constants in the C.F. by variable functions.

- Consider a linear non-homogeneous second order differential equation with variable coefficients

$$\frac{d^2y}{dx^2} + P(x) \frac{dy}{dx} + Q(x)y = X(x) \quad (1)$$

Suppose the complimentary function of (1) is  $= C_1y_1(x) + C_2y_2(x)$  (2)

so that  $y_1$  and  $y_2$  satisfy

$$\frac{d^2y}{dx^2} + P(x) \frac{dy}{dx} + Q(x)y = 0$$

In method of variation of parameters the arbitrary constants  $C_1$  and  $C_2$  in (2) are replaced by two unknown functions  $u(x)$  and  $v(x)$ .

Let us assume particular integral is  $= u(x)y_1(x) + v(x)y_2(x)$  (3)

where 
$$u = \int \frac{-X y_2}{y_1 y_2' - y_1' y_2} dx$$

and 
$$v = \int \frac{X y_1}{y_1 y_2' - y_1' y_2} dx$$

on putting the values of  $u$  and  $v$  in (3) we get P.I

Thus, required general solution  $= C.F + P.I$

**Example 10.** Apply the method of variation of parameters to solve

(U.P.T.U. 2009)

$$\frac{d^2y}{dx^2} + y = \tan x$$

**Solution.** The auxiliary equation is  $m^2 + 1 = 0 \Rightarrow m = \pm i$

$\therefore$  C.F.  $= C_1 \cos x + C_2 \sin x$  (1)

Here  $y_1 = \cos x, y_2 = \sin x$

Therefore  $y_1 y_2' - y_1' y_2 = \cos^2 x + \sin^2 x = 1$

Let us suppose P.I  $= u.y_1 + v.y_2$  (2)

where

$$u = \int \frac{-X y_2}{y_1 y_2' - y_1' y_2} dx = - \int \frac{\sin x \tan x}{1} dx$$

Equations Reducible To Linear Equations with Constant Coefficients

$$\begin{aligned}
 &= - \int \frac{\sin^2 x}{\cos x} dx = - \int \frac{1 - \cos^2 x}{\cos x} dx \\
 &= \int (\cos x - \sec x) dx \\
 &= \sin x - \log (\sec x + \tan x)
 \end{aligned}$$

$$\begin{aligned}
 \& \quad v = \int \frac{-X y_1}{y_1 y_2' - y_1' y_2} dx = \int \frac{\tan x \cos x}{1} dx \\
 &= \int \sin x dx = - \cos x
 \end{aligned}$$

Putting the values of u and v in (2), we get

$$\begin{aligned}
 \text{P.I} &= u y_1 + v y_2 \\
 &= [\sin x - \log (\sec x + \tan x)] \cos x - \cos x \sin x \\
 &= - \cos x \log (\sec x + \tan x)
 \end{aligned}$$

Therefore, complete solution is

$$y = C_1 \cos x + C_2 \sin x - \cos x \log (\sec x + \tan x)$$

**Example 11.** Use variation of parameters to solve

$$\frac{d^2 y}{dx^2} + y = \sec x$$

(U.P.T.U. 2002)

**Solution.** The auxiliary equation is

$$m^2 + 1 = 0$$

$$\Rightarrow m = \pm i$$

$$\therefore \text{C.F} = C_1 \cos x + C_2 \sin x \tag{1}$$

Here  $y_1 = \cos x, y_2 = \sin x$

$$\text{Let us suppose P.I.} = u y_1 + v y_2 \tag{2}$$

$$\text{where } u = \int \frac{-\sec x \sin x}{1} dx \quad \therefore u = \int \frac{-X y_2}{y_1 y_2' - y_1' y_2} dx$$

$$\text{As } y_1 y_2' - y_1' y_2 = \cos x \cos x - (-\sin x) \sin x$$

$$= \cos^2 x + \sin^2 x$$

$$= 1$$

$$= - \int \tan x \, dx$$

$$= \log \cos x$$

and 
$$v = \int \frac{Xy_1'}{y_1y_2' - y_1'y_2} \, dx$$

$$= \int \frac{\cos x \sec x}{1} \, dx = \int dx = x$$

putting the values of u and v in (2), we get

$$\text{P.I} = \log \cos x \cdot \cos x + x \cdot \sin x$$

Therefore, complete solution is

$$y = C_1 \cos x + C_2 \sin x + \cos x \cdot \log \cos x + x \sin x$$

**Example 12.** Using the method of variation of parameters solve

$$\frac{d^2y}{dx^2} + 4y = 4 \tan 2x$$

(I.A.S. 2001, U.P.T.U. 2006)

**Solution.** Here the auxiliary equations

$$m^2 + 4 = 0 \quad \Rightarrow \quad m = \pm 2i$$

$$\therefore \text{C.F} = C_1 \cos 2x + C_2 \sin 2x \quad (1)$$

Here  $y_1 = \cos 2x, y_2 = \sin 2x$

Let us suppose P.I. =  $uy_1 + vy_2$  (2)

where 
$$u = \int \frac{-X y_2 dx}{y_1 y_2' - y_1' y_2} = \int \frac{-4 \tan 2x \sin 2x}{2} \, dx$$

$$\therefore \text{As } y_1 y_2' - y_1' y_2 = 2 \cos 2x \cos 2x + 2 \sin 2x \sin 2x = 2$$

$$= - \int 2 \frac{\sin^2 2x}{\cos 2x} \, dx = - \int \frac{1 - \cos^2 2x}{\cos 2x} \, dx$$

$$= 2 \int (\cos 2x - \sec 2x) \, dx$$

$$= \sin 2x - \log (\sec 2x + \tan 2x)$$

Equations Reducible To Linear Equations with Constant Coefficients

and 
$$v = \int \frac{X y_1}{y_1 y_2' - y_1' y_2} dx = \int \frac{4 \tan 2x \cos 2x}{2} dx$$

$$= 2 \int \sin 2x dx = -\cos 2x$$

putting the values of u and v in (2) we get

$$P.I = \{\sin 2x - \log(\sec 2x + \tan 2x)\} \cos 2x - \cos 2x \sin 2x$$

$$= -\cos 2x \log(\sec 2x + \tan 2x)$$

Hence, the complete solution is

$$y = C_1 \cos 2x + C_2 \sin 2x - \cos 2x \log(\sec 2x + \tan 2x)$$

**Example 13.** Obtain general solution of the differential equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - y = x^3 e^x$$

(U.P.T.U. 2002)

**Solution.** On changing the independent variable by substituting  $x = e^z$  or  $z = \log_e x$  and  $\frac{d}{dz} \equiv D$  the differential equation becomes

$$[D(D-1) + D-1]y = e^{3z} e^{e^z}$$

or  $(D^2 - 1)y = e^{3z} e^{e^z}$

Here auxiliary equation is  $m^2 - 1 = 0 \Rightarrow m = \pm 1$

$\therefore$  C.F =  $C_1 e^z + C_2 e^{-z}$

$\Rightarrow$  C.F =  $C_1 x + \frac{C_2}{x}$

Let P.I =  $u y_1 + v y_2$

Here  $y_1 = x$  and  $y_2 = \frac{1}{x}$

Also 
$$u = \int \frac{-X y_2 dx}{y_1 y_2' - y_1' y_2} = \int \frac{-x^3 e^x \frac{1}{x} dx}{x \left( -\frac{1}{x^2} \right) - \frac{1}{x} (1)} = \int \frac{-x^2 e^x dx}{-\frac{2}{x}}$$

$$= \frac{1}{2} \int x^3 e^x dx$$

$$\begin{aligned}
 u &= \frac{1}{2} [x^3 e^x - 3x^2 e^x + 6xe^x - 6e^x] \\
 &= \frac{1}{2} [x^3 - 3x^2 + 6x - 6] e^x \\
 \& \ v &= \int \frac{X y_1 dx}{y_1 y_2' - y_1' y_2} = \int \frac{x^3 e^x x dx}{x \left(-\frac{1}{x^2}\right) - \frac{1}{x}} \quad (1) \\
 &= \int \frac{x^4 e^x}{-\frac{2}{x}} dx = -\frac{1}{2} \int x^5 e^x dx
 \end{aligned}$$

or 
$$v = -\frac{1}{2} [x^5 e^x - 5x^4 e^x + 20x^3 e^x - 60x^2 e^x + 120xe^x - 120e^x]$$

putting the values of u and v in equation (2) we get

$$\begin{aligned}
 P.I &= \frac{1}{2} (x^3 - 3x^2 + 6x - 6) e^x x - \frac{1}{2} (x^5 - 5x^4 + 20x^3 - 60x^2 + 120x - 120) e^x \frac{1}{x} \\
 &= \frac{e^x}{2} \left[ x^4 - 3x^3 + 6x^2 - 6x - x^4 + 5x^3 - 20x^2 + 60x - 120 + \frac{120}{x} \right] \\
 &= \frac{e^x}{2} \left[ 2x^3 - 14x^2 + 54x - 120 + \frac{120}{x} \right]
 \end{aligned}$$

Hence the required solution is  $y = C.F + P.I$

or 
$$y = C_1 x + \frac{C_2}{x} + (x^3 - 7x^2 + 27x - 60 + \frac{60}{x}) e^x$$

**Example 14.** Solve by method of variation of parameters

$$\frac{d^2y}{dx^2} - y = \frac{2}{1 + e^x} \quad (\text{U.P.T.U 2001})$$

**Solution.** Here auxiliary equation is  $m^2 - 1 = 0$

$\Rightarrow m = \pm 1$

$\therefore C.F = C_1 e^x + C_2 e^{-x}$

Here  $y_1 = e^x, y_2 = e^{-x}$

Let  $P.I = uy_1 + vy_2$



Equations Reducible To Linear Equations with Constant Coefficients

where 
$$u = \int \frac{-Xy_2}{y_1 y_2' - y_1' y_2} dx = \int \frac{2}{1+e^x} \frac{e^{-x}}{-2} dx$$

$$\therefore y_1 y_2' - y_1' y_2 = -e^x e^{-x} - e^x e^{-x} = -2$$

$$= \int \frac{e^{-x}}{1+e^x} dx = \int \frac{1}{e^x(1+e^x)} dx = \int \left( \frac{1}{e^x} - \frac{1}{1+e^x} \right) dx$$

$$= \int e^{-x} dx - \int \frac{e^{-x}}{e^{-x}+1} dx$$

$$= -e^{-x} + \log(e^{-x}+1)$$

$$v = \int \frac{Xy_1}{y_1 y_2' - y_1' y_2} dx = \int \frac{e^x}{-2} \frac{2}{1+e^x} dx$$

$$= - \int \frac{e^x}{1+e^x} dx = -\log(1+e^x)$$

$$P.I = u y_1 + v y_2$$

$$= [-e^{-x} + \log(e^{-x}+1)] e^x - e^{-x} \log(1+e^x)$$

$$= -1 + e^x \log(e^{-x}+1) - e^{-x} \log(e^x+1)$$

$$\therefore y = C_1 e^x + C_2 e^{-x} - 1 + e^x \log(e^{-x}+1) - e^{-x} \log(e^x+1)$$

**Example 15.** Apply the method of variation of parameters to solve

$$\frac{d^2y}{dx^2} - 3 \frac{dy}{dx} + 2y = \frac{e^x}{1+e^x}$$

(U.P.T.U. 2005)

**Solution.** Here auxiliary equation is

$$m^2 - 3m + 2 = 0$$

$$\Rightarrow (m-1)(m-2) = 0$$

$$\Rightarrow m = 1, 2$$

$$\therefore C.F. = C_1 e^x + C_2 e^{2x}$$

$$\text{Here } y_1 = e^x, y_2 = e^{2x}$$

$$P.I = u y_1 + v y_2$$

where 
$$u = \int -\frac{Xy_2}{y_1 y_2' - y_1' y_2} dx$$

$$= \int -\frac{\frac{e^x}{1+e^x} e^{2x}}{e^x (2e^{2x}) - e^{2x} (e^x)} dx = \int \frac{-e^{-3x}}{2e^{3x} - e^{3x}} dx$$

$$= \int \frac{-e^{-3x}}{e^{3x} (1+e^x)} dx = -\int \frac{1}{1+e^x} dx$$

$$= -\int \frac{e^{-x} dx}{e^{-x} + 1} = \log (e^{-x} + 1)$$

and 
$$v = \int \frac{X y_1}{y_1 y_2' - y_1' y_2} dx = \int \frac{\frac{e^x}{1+e^x} e^x}{e^x (2e^{2x}) - e^{2x} (e^x)} dx$$

$$= \int \frac{\frac{e^{2x}}{1+e^x}}{2e^{3x} - e^{3x}} dx = \int \frac{e^{2x}}{e^{3x} (1+e^x)} dx$$

$$= \int \frac{1}{e^x (1+e^x)} dx = \int \left( \frac{1}{e^x} - \frac{1}{1+e^x} \right) dx$$

$$= \int \left( e^{-x} - \frac{e^{-x}}{e^{-x} + 1} \right) dx = -e^{-x} + \log (e^{-x} + 1)$$

Therefore P.I =  $e^x \log (e^{-x} + 1) + e^{2x} \{-e^{-x} + \log (e^{-x} + 1)\}$   
 $= e^x \log (e^{-x} + 1) - e^{-x} + e^{2x} \log (e^{-x} + 1)$

Therefore, complete solution is  $y = C.F + P.I$

or  $y = C_1 e^x + C_2 e^{2x} + e^x \log (e^{-x} + 1) - e^{-x} + e^{2x} \log (e^{-x} + 1)$

**Example 16.** Solve by the method of variation of parameters

$$\frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} = e^x \sin x$$

(U.P.T.U. 2003)

**Solution.** Here auxiliary equation is  $m^2 - 2m = 0$

$\Rightarrow m(m - 2) = 0$

Equations Reducible To Linear Equations with Constant Coefficients

$$\begin{aligned} \Rightarrow \quad & m = 0, 2 \\ \therefore \quad & \text{C.F.} = C_1 + C_2 e^{2x} \end{aligned} \tag{1}$$

Here  $y_1 = 1, y_2 = e^{2x}$

$$\text{P.I} = uy_1 + vy_2 \tag{2}$$

$$\begin{aligned} \text{where } u &= \int \frac{-X y_2}{y_1 y_2' - y_1' y_2} dx = \int \frac{-e^x \sin x \cdot e^{2x}}{1(2e^{2x}) - 0(e^{2x})} dx \\ &= -\frac{1}{2} \int e^x \sin x dx \\ &= -\frac{1}{2} \frac{e^x}{(1)^2 + (1)^2} (\sin x - \cos x) \\ &= -\frac{1}{4} e^x (\sin x - \cos x) \end{aligned}$$

$$\begin{aligned} \text{and } v &= \int \frac{X y_1}{y_1 y_2' - y_1' y_2} dx \\ &= \int \frac{e^x \sin x \cdot 1}{1(2e^{2x}) - 0(e^{2x})} = \frac{1}{2} \int e^{-x} \sin x dx \\ &= \frac{1}{2} \frac{e^{-x}}{(-1)^2 + (1)^2} (-\sin x - \cos x) \\ &= -\frac{1}{2} \frac{e^{-x}}{2} (\sin x + \cos x) = -\frac{e^{-x}}{4} (\sin x + \cos x) \end{aligned}$$

putting the values of  $u$  and  $v$  in (2), we get

$$\begin{aligned} \text{P.I} &= \frac{e^x}{-4} (\sin x - \cos x) + \frac{e^{-x}}{4} (\sin x + \cos x) e^{2x} \\ &= \frac{e^x}{-4} (\sin x - \cos x + \sin x + \cos x) = \frac{e^{-x}}{-2} \sin x \end{aligned}$$

Hence, the complete solution is  $y = \text{C.F.} + \text{P.I}$

$$y = C_1 + C_2 e^{2x} - \frac{e^x}{2} \sin x$$

**Example 18.** Solve by method of variation of parameters

$$\frac{d^2y}{dx^2} + 2 \frac{dy}{dx} + y = e^{-x} \log x$$

(U.P.T.U. 2008)

**Solution.** Here auxiliary equation is  $m^2 + 2m + 1 = 0$

$$\Rightarrow (m + 1)^2 = 0$$

$$\Rightarrow m = -1, -1$$

$$\therefore \text{C.F} = (C_1 + C_2 x) e^{-x} \quad (1)$$

Here  $y_1 = e^{-x}$ ,  $y_2 = x e^{-x}$

$$\text{P.I} = uy_1 + vy_2 \quad (2)$$

where

$$\begin{aligned} u &= \int \frac{-Xy_2}{y_1 y_2' - y_1' y_2} dx \\ &= \int \frac{-e^{-x} \log x \cdot x e^{-x}}{-x e^{-2x} + e^{-2x} + x e^{-2x}} dx \\ &= \int \frac{-x e^{-2x} \log x}{e^{-2x}} dx = - \int x \log x dx \\ &= -\frac{x^2}{2} \log x + \int \frac{x^2}{2} \frac{1}{x} dx \\ &= -\frac{x^2}{2} \log x + \frac{x^2}{4} \end{aligned}$$

and

$$\begin{aligned} v &= \int \frac{X y_1}{y_1 y_2' - y_1' y_2} dx \\ &= \int \frac{e^{-x} \log x \cdot e^{-x}}{e^{-2x}} dx \\ &= \int \log x dx \\ &= x \log x - \int \frac{1}{x} x dx \\ &= x \log x - x \end{aligned}$$

Putting these values of  $u$  and  $v$  in (2) we get

Equations Reducible To Linear Equations with Constant Coefficients

$$P.I = -\frac{x^2 e^{-x}}{2} \log x + e^{-x} \frac{x^2}{4} + x^2 e^{-x} \log x - x^2 e^{-x}$$

or 
$$P.I = \frac{x^2 e^{-x}}{2} \log x - \frac{3}{4} x^2 e^{-x}$$

Hence, complete solution is  $y = C.F + P.I$

or 
$$y = (C_1 + C_2 x) e^{-x} + \frac{x^2 e^{-x}}{2} \log x - \frac{3}{4} x^2 e^{-x}$$

**Example 19.** Using variation of parameters method, solve

$$x^2 \frac{d^2 y}{dx^2} + 2x \frac{dy}{dx} - 12y = x^3 \log x$$

(U.P.T.U. 2004)

**Solution.** On changing the independent variable by substituting  $x = e^z$  or  $z = \log_e x$  and  $\frac{d}{dz} \equiv D$ , the differential equation becomes

$$\{D(D-1) + 2D - 12\} y = z e^{3z}$$

or 
$$(D^2 + D - 12) y = z e^{3z}$$

The auxiliary equation is  $m^2 + m - 12 = 0$

$\Rightarrow m = 3, -4$

$\therefore C.F = C_1 e^{3z} + C_2 e^{-4z}$

or 
$$C.F = C_1 x^3 + C_2 \frac{1}{x^4} \tag{1}$$

Here  $y_1 = x^3, y_2 = \frac{1}{x^4}$

$$P.I. = uy_1 + vy_2 \tag{2}$$

where

$$\begin{aligned} u &= \int \frac{-X y_2}{y_1 y_2' - y_1' y_2} dx \\ &= - \int \frac{x \log x \cdot x^{-4}}{x^3 (-4x^{-5}) - 3x^2 (x^{-4})} dx = - \int \frac{x^{-3} \log x}{-7x^{-2}} dx \\ &= \frac{1}{7} \int \frac{\log x}{x} dx \end{aligned}$$

$$\begin{aligned} &= \frac{1}{7} \frac{(\log x)^2}{2} \\ &= \frac{1}{14} (\log x)^2 \end{aligned}$$

and

$$\begin{aligned} v &= \int \frac{X y_1}{y_1 y_2' - y_1' y_2} dx \\ &= \int \frac{x \log x \cdot x^3}{-7 x^{-2}} dx = \frac{-1}{7} \int x^6 \log x dx \\ &= -\frac{1}{7} \left[ \log x \cdot \frac{x^7}{7} - \int \frac{1}{x} \frac{x^7}{7} dx \right] \\ &= -\frac{1}{7} \left[ \frac{x^7 \log x}{7} - \frac{1}{7} \left( \frac{x^7}{7} \right) \right] \\ &= \frac{x^7}{49} \left( \frac{1}{7} - \log x \right) \end{aligned}$$

putting the values of u and v in (1) we get

$$\begin{aligned} \text{P.I} &= \frac{1}{14} (\log x)^2 x^3 + \frac{1}{x^4} \frac{x^7}{49} \left( \frac{1}{7} - \log x \right) \\ &= \frac{x^3}{14} (\log x)^2 + \frac{x^3}{49} \left( \frac{1}{7} - \log x \right) \end{aligned}$$

Therefore, the required solution is  $y = \text{C.F} + \text{P.I}$

or 
$$y = C_1 x^3 + \frac{C_2}{x^4} + \frac{x^3}{14} (\log x)^2 + \frac{x^3}{343} - \frac{x^3}{49} \log x$$

or 
$$y = \left( C_1 + \frac{1}{343} \right) x^3 + \frac{C_2}{x^4} + \frac{x^3}{98} \log x (7 \log x - 2)$$

## SECOND ORDER LINEAR DIFFERENTIAL EQUATIONS WITH VARIABLE COEFFICIENTS (Solution by Changing dependent and independent variables)

### INTRODUCTION

The general form of linear differential equation of the second order may be written as

$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = R \quad (1)$$

where  $P$ ,  $Q$  and  $R$  are functions of  $x$  only. There is no general method for the solution of this type of equations. Some particular methods used to solve these equations are, change of independent variables, Variation of parameters and removal of first order derivatives etc. As this kind of differential equations are of great significance in physics, especially in connection with vibrations in mechanics and theory of electric circuit. In addition many profound and beautiful ideas in pure mathematics have grown out to the study of these equations.

**Method I: Complete solution is terms of known integral belonging to the complementary function (i.e. part of C.F. is known or one solution is known).**

Let  $u$  be a part of complementary function of equation (1) and  $v$  is remaining solution of differential equation (1)

Then the complete solution of equation (1) is

$$y = u v \quad (2)$$

$$\Rightarrow \frac{dy}{dx} = v \frac{du}{dx} + u \frac{dv}{dx} \text{ and } \frac{d^2y}{dx^2} = v \frac{d^2u}{dx^2} + 2 \frac{du}{dx} \frac{dv}{dx} + u \frac{d^2v}{dx^2}$$

Putting these values in equation (1) then, we get

$$v \frac{d^2u}{dx^2} + 2 \frac{du}{dx} \frac{dv}{dx} + u \frac{d^2v}{dx^2} + P \left( v \frac{du}{dx} + u \frac{dv}{dx} \right) + Quv = R$$

$$\text{or } v \left[ \frac{d^2u}{dx^2} + P \frac{du}{dx} + Qu \right] + u \left[ \frac{d^2v}{dx^2} + P \frac{dv}{dx} \right] + 2 \frac{du}{dx} \frac{dv}{dx} = R \quad (3)$$

Since  $u$  is a part of C.F. i.e. solution of (1)

$$\therefore \frac{d^2u}{dx^2} + P \frac{du}{dx} + Qu = 0$$

Hence equation (3) becomes

$$u \left( \frac{d^2v}{dx^2} + P \frac{dv}{dx} \right) + 2 \frac{du}{dx} \frac{dv}{dx} = R$$

or 
$$\frac{d^2v}{dx^2} + \left( P + \frac{2}{u} \frac{du}{dx} \right) \frac{dv}{dx} = \frac{R}{u} \quad (4)$$

Let 
$$\frac{dv}{dx} = z, \text{ so that } \frac{d^2v}{dx^2} = \frac{dz}{dx}$$

Equation (4) becomes

$$\frac{dz}{dx} + \left( P + \frac{2}{u} \frac{du}{dx} \right) z = \frac{R}{u},$$

which is linear in  $z$ , Hence  $z$  can be determined

We obtain  $v$ , by integration the relation  $\frac{dv}{dx} = z$

$$\Rightarrow v = \int z \, dx + C_1$$

Therefore, the solution of (1) is  $y = u \left[ \int z \, dx + C_1 \right]$

i.e.  $y = uv$

**Remark.** Solving by the above method,  $u$  determined by inspection of the following rules

- (1) If  $P + Qx = 0$ , then  $u = x$
- (2) If  $1 + P + Q = 0$ , then  $u = e^x$
- (3) If  $1 - P + Q = 0$ , then  $u = e^{-x}$
- (4) If  $1 + \frac{P}{a} + \frac{Q}{a^2} = 0$ , then  $u = e^{ax}$
- (5) If  $2 + 2Px + Qx^2 = 0$ , then  $u = x^2$
- (6) If  $m(m - 1) + Pmx + Qx^2 = 0$ , then  $u = x^m$

**Example 20.** Solve  $y'' - 4xy' + (4x^2 - 2)y = 0$  given that  $y = e^{x^2}$  is an integral induced in the complementary function.

(U.P.T.U. 2004)

**Solution.** The given equation may be written as



Equations Reducible To Linear Equations with Constant Coefficients

$$\frac{d^2y}{dx^2} - 4x \frac{dy}{dx} + (4x^2 - 2)y = 0$$

Here  $P = -4x$ ,  $Q = 4x^2 - 2$ ,  $R = 0$

and  $u = e^{x^2}$ , so that  $\frac{du}{dx} = 2xe^{x^2}$

Let  $y = uv \Rightarrow y = e^{x^2} v$  (1)

we know that

$$\frac{d^2v}{dx^2} + \left( P + \frac{2}{u} \frac{du}{dx} \right) \frac{dv}{dx} = 0, \text{ As } R = 0$$

$$\Rightarrow \frac{d^2v}{dx^2} + \left( \frac{2}{e^{x^2}} 2xe^{x^2} - 4x \right) \frac{dv}{dx} = 0$$

$$\Rightarrow \frac{d^2v}{dx^2} = 0$$

$$\Rightarrow \frac{dv}{dx} = C$$

$$\Rightarrow v = C_1x + C_2$$

Hence the complete solution is  $y = e^{x^2} v$

or  $y = e^{x^2} (C_1x + C_2)$

**Example 21.** By the method of variation of parameters, solve the differential equation (U.P.T.U. 2004)

$$\frac{d^2y}{dx^2} + (1 - \cot x) \frac{dy}{dx} - y \cot x = \sin^2 x$$

**Solution.** Here  $P = 1 - \cot x$ ,  $Q = -\cot x$

Therefore  $1 - P + Q = 1 - (1 - \cot x) - \cot x = 0$

That is  $y = e^{-x}$  is a part of the C.F. putting  $y = ve^{-x}$

$$\frac{dy}{dx} = -ve^{-x} + e^{-x} \frac{dv}{dx}$$

and 
$$\frac{d^2y}{dx^2} = e^{-x} \frac{d^2v}{dx^2} - 2e^{-x} \frac{dv}{dx} + ve^{-x}$$

on putting these values in the given differential equation, we have

$$\frac{d^2v}{dx^2} - (1 + \cot x) \frac{dv}{dx} = 0$$

or 
$$\frac{dp}{dx} - (1 + \cot x) p = 0 \quad \text{where } p = \frac{dv}{dx}$$

$$\Rightarrow \frac{dp}{p} = (1 + \cot x) dx$$

on integrating we get

$$\log p = x + \log \sin x + \log C_1$$

$$\Rightarrow p = C_1 e^x \sin x$$

Substituting for p

$$\frac{dv}{dx} = C_1 e^x \sin x$$

or 
$$dv = C_1 e^x \sin x dx$$

Integrating

$$\begin{aligned} v &= C_1 \int e^x \sin x dx + C_2 \\ &= C_1 \frac{1}{2} e^x (\sin x - \cos x) + C_2 \end{aligned}$$

Therefore, solution of the given differential equation i.e. C.F. is given by

$$y = ve^{-x} = C_1 \frac{1}{2} (\sin x - \cos x) + C_2 e^{-x}$$

Let  $y = Au + Bv$  be the complete solution of the given differential equation where A and B are the functions of x, i.e.

$$y = A (\sin x - \cos x) + Be^{-x} \tag{1}$$

Differentiating on both sides

$$\frac{dy}{dx} = A (\cos x + \sin x) - Be^{-x} + \frac{dA}{dx} (\sin x - \cos x) + \frac{dB}{dx} e^{-x}$$

Equations Reducible To Linear Equations with Constant Coefficients

Let us choose A and B such that

$$\frac{dA}{dx} (\sin x - \cos x) + \frac{dB}{dx} e^{-x} = 0 \quad (2)$$

$$\Rightarrow \frac{dy}{dx} = A (\cos x + \sin x) - Be^{-x}$$

$$\text{and } \frac{d^2y}{dx^2} = \frac{dA}{dx} (\cos x + \sin x) - \frac{dB}{dx} e^{-x} + A (-\sin x + \cos x) + Be^{-x}$$

putting these values of  $\frac{d^2y}{dx^2}$ ,  $\frac{dy}{dx}$  and y in the given equation, we get

$$\frac{dA}{dx} (\cos x + \sin x) - \frac{dB}{dx} e^{-x} = \sin^2 x \quad (3)$$

on solving equation (2) and (3), we get

$$\frac{dA}{dx} = \frac{1}{2} \sin x$$

Integrating,

$$A = -\frac{1}{2} \cos x + C_1$$

$$\text{and } \frac{dB}{dx} = \frac{1}{2} e^x (\sin x \cos x - \sin^2 x)$$

$$= \frac{e^x}{4} (\sin 2x + \cos 2x - 1)$$

on integration, we have

$$B = \frac{e^x}{20} (3 \sin 2x - \cos 2x) - \frac{e^x}{4} + C_2$$

putting the values of A and B in equation (1) we get

$$y = \left( -\frac{1}{2} \cos x + C_1 \right) (\sin x - \cos x) \left( \frac{e^x}{20} (3 \sin 2x - \cos 2x) - \frac{e^x}{4} + C_2 \right) e^{-x}$$

$$= -\frac{1}{2} \cos x \sin x + \frac{1}{2} \cos^2 x + C_1 \sin x - C_1 \cos x + \frac{3}{20} \sin 2x - \frac{1}{20} \cos 2x - \frac{1}{4} + C_2 e^{-x}$$

or  $y = C_1 (\sin x - \cos x) + C_2 e^{-x} - \frac{1}{10} (\sin 2x - 2 \cos 2x)$

**Method II. Normal form (Removal of first derivative)**

Let  $\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = R$  (1)

putting  $y = uv$ , we get

$$v \left( \frac{d^2u}{dx^2} + P \frac{du}{dx} + Qu \right) + u \left( \frac{d^2v}{dx^2} + P \frac{dv}{dx} \right) + 2 \frac{du}{dx} \frac{dv}{dx} = R$$

$$\Rightarrow \frac{d^2v}{dx^2} + \left( \frac{2}{u} \frac{du}{dx} + P \right) \frac{dv}{dx} + v \left( \frac{1}{u} \frac{d^2u}{dx^2} + \frac{P}{u} \frac{du}{dx} + Q \right) = \frac{R}{u}$$
 (2)

But the first order derivative must be remove

so  $\frac{2}{u} \frac{du}{dx} + P = 0 \Rightarrow \frac{du}{u} = -\frac{1}{2} P dx$

$$\Rightarrow \log u = -\int \frac{P}{2} dx$$

$$\Rightarrow u = e^{-\int \frac{P}{2} dx}$$

Since  $\frac{du}{dx} = -\frac{Pu}{2} \Rightarrow \frac{d^2u}{dx^2} = -\frac{1}{2} \left[ P \frac{du}{dx} + u \frac{dP}{dx} \right]$

$$\Rightarrow \frac{d^2u}{dx^2} = -\frac{1}{2} \left[ P \left( -\frac{Pu}{2} \right) + u \frac{dP}{dx} \right] = \frac{P^2u}{4} - \frac{u}{2} \frac{dP}{dx}$$

From (2)  $\frac{d^2v}{dx^2} + v \left( \frac{P^2}{4} - \frac{1}{2} \frac{dP}{dx} - \frac{P^2}{2} + Q \right) = \frac{R}{u}$

$$\Rightarrow \frac{d^2v}{dx^2} + v \left( Q - \frac{1}{2} \frac{dP}{dx} - \frac{P^2}{4} \right) = \frac{R}{u}$$

$$\Rightarrow \frac{d^2v}{dx^2} + Iv = \frac{R}{u}$$
 (3)

This equation is called normal form of equation (1)

Equations Reducible To Linear Equations with Constant Coefficients

where  $I = Q - \frac{1}{2} \frac{dP}{dx} - \frac{P^2}{4}$

**Example 22.** Solve

$$\frac{d^2y}{dx^2} - 4x \frac{dy}{dx} + (4x^2 - 1)y = -3e^{x^2} \sin 2x$$

[I.A.S. 2000, U.P.T.U. (C.O.) 2004]

**Solution.** Here  $P = -4x$ ,  $Q = 4x^2 - 1$ ,  $R = -3e^{x^2} \sin 2x$

so 
$$I = Q - \frac{1}{2} \frac{dP}{dx} - \frac{1}{4} P^2 = 4x^2 - 1 - \frac{1}{2} (-4) - \frac{1}{4} (-4x)^2$$
$$= 4x^2 - 1 + 2 - 4x^2 = 1$$

$$u = e^{-\frac{1}{2} \int P dx} = e^{-\frac{1}{2} \int (-4x) dx}$$
$$= e^{2 \int x dx} = e^{x^2}$$

Then substituting these values in the equation

$$\frac{d^2v}{dx^2} + Iv = \frac{R}{u}, \text{ We have}$$
$$\frac{d^2v}{dx^2} + v = \frac{-3e^{x^2} \sin 2x}{e^{x^2}} = -3 \sin 2x$$

its C.F =  $C_1 \cos x + C_2 \sin x$

and P.I =  $-\frac{1}{D^2 + 1} 2 \sin 2x = -3 \frac{1}{-2^2 + 1} \sin 2x$ 
$$= \sin 2x$$

Thus  $v = C_1 \cos x + C_2 \sin x + \sin 2x$

Therefore required solution is  $y = uv$

or  $y = e^{x^2} (C_1 \cos x + C_2 \sin x + \sin 2x)$

**Example 23.** Solve

$$\frac{d^2y}{dx^2} - 4x \frac{dy}{dx} + (4x^2 - 3)y = e^{x^2}$$

by removing first derivative

**Solution.** Here  $P = -4x$ ,  $Q = 4x^2 - 3$ ,  $R = e^{x^2}$

$$\begin{aligned} I &= Q - \frac{1}{2} \frac{dP}{dx} - \frac{P^2}{4} = 4x^2 - 3 - \frac{1}{2} (-4) - \frac{1}{4} (-4x)^2 \\ &= 4x^2 - 3 + 2 - 4x^2 = -1 \\ w &= e^{-\frac{1}{2} \int P dx} \\ &= e^{-\frac{1}{2} \int (-4x) dx} = e^{2 \int x dx} = e^{x^2} \end{aligned}$$

Then substituting these values in the equation

$$\frac{d^2v}{dx^2} + Iv = \frac{R}{u}, \text{ we get}$$

$$\frac{d^2v}{dx^2} - v = \frac{e^{x^2}}{e^{x^2}}$$

or 
$$\frac{d^2v}{dx^2} - v = 1$$

its C.F =  $C_1 e^x + C_2 e^{-x}$

and 
$$\begin{aligned} P.I &= \frac{1}{D^2 - 1} 1 = -(1 - D^2)^{-1} 1 \\ &= -(1 + D^2 + D^4 + \dots) 1 \\ &= -1 \end{aligned}$$

Thus  $v = C_1 e^x + C_2 e^{-x} - 1$

Hence the general solution of the given equation is

$$y = uv$$

or 
$$y = e^{x^2} (C_1 e^x + C_2 e^{-x} - 1)$$

### Method III. Change of independent variable

consider 
$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = R \tag{1}$$

Let us change the independent variable  $x$  to  $z$  and  $z = f(x)$ .

Equations Reducible To Linear Equations with Constant Coefficients

Then 
$$\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx} \tag{2}$$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dz} \frac{dz}{dx} \right) = \frac{dy}{dz} \frac{d^2z}{dx^2} + \left( \frac{dz}{dx} \right)^2 \frac{d^2y}{dx^2} \tag{3}$$

Putting the values of  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$  in (1) we get

$$\frac{d^2y}{dz^2} \left( \frac{dz}{dx} \right)^2 + \frac{dy}{dz} \frac{d^2z}{dx^2} + P \frac{dy}{dz} \frac{dz}{dx} + Qy = R$$

or 
$$\frac{d^2y}{dz^2} \left( \frac{dz}{dx} \right)^2 + \left( P \frac{dz}{dx} + \frac{d^2z}{dx^2} \right) \frac{dy}{dz} + Qy = R$$

or 
$$\frac{d^2y}{dz^2} + \frac{\left( P \frac{dz}{dx} + \frac{d^2z}{dx^2} \right)}{\left( \frac{dz}{dx} \right)^2} \frac{dy}{dz} + \frac{Q}{\left( \frac{dz}{dx} \right)^2} y = \frac{R}{\left( \frac{dz}{dx} \right)^2}$$

$$\Rightarrow \frac{d^2y}{dz^2} + P_1 \frac{dy}{dz} + Q_1 y = R_1 \tag{4}$$

where 
$$P_1 = \frac{\left( P \frac{dz}{dx} + \frac{d^2z}{dx^2} \right)}{\left( \frac{dz}{dx} \right)^2}, Q_1 = \frac{Q}{\left( \frac{dz}{dx} \right)^2} \text{ and } R_1 = \frac{R}{\left( \frac{dz}{dx} \right)^2}$$

Equation (4) is solved either by taking  $P_1 = 0$  or  $Q_1 = \text{a constant}$

**Example 24.** Solve by changing the independent variable

$$x \frac{d^2y}{dx^2} - \frac{dy}{dx} + 4x^3y = x^5 \quad (\text{U.P.T.U. 2002, 2003})$$

**Solution.** Given equation is

$$\frac{d^2y}{dx^2} - \frac{1}{x} \frac{dy}{dx} + 4x^2y = x^4 \tag{1}$$

Here  $P = -\frac{1}{x}$ ,  $Q = 4x^2$  and  $R = x^4$

On changing the independent variable  $x$  to  $z$ , the equation (1) transformed as

$$\frac{d^2y}{dz^2} + P_1 \frac{dy}{dz} + Q_1 y = R_1 \quad (2)$$

where 
$$Q_1 = \frac{Q}{\left(\frac{dz}{dx}\right)^2} = \frac{4x^2}{\left(\frac{dz}{dx}\right)^2} = \text{constant} = 1 \text{ say}$$

or 
$$\left(\frac{dz}{dx}\right)^2 = 4x^2$$

$$\Rightarrow \frac{dz}{dx} = 2x$$

$$\Rightarrow z = x^2$$

$$\Rightarrow \frac{d^2z}{dx^2} = 2$$

$$P_1 = \frac{P \frac{dz}{dx} + \frac{d^2z}{dx^2}}{\left(\frac{dz}{dx}\right)^2} = \frac{\left\{2 + \left(\frac{-1}{x}\right) 2x\right\}}{4x^2} = 0$$

$$R_1 = \frac{R}{\left(\frac{dz}{dx}\right)^2} = \frac{x^4}{4x^2} = \frac{x^2}{4} = \frac{z}{4}$$

on putting the values of  $P_1$ ,  $Q_1$  and  $R_1$  in (2), we get

$$\frac{d^2y}{dz^2} + y = \frac{z}{4}$$

or 
$$(D^2 + 1)y = \frac{z}{4}$$

its A.E. is  $m^2 + 1 = 0 \Rightarrow m = \pm i$

$\therefore$  C.F =  $C_1 \cos z + C_2 \sin z$

or C.F =  $C_1 \cos x^2 + C_2 \sin x^2$

and P.I =  $\frac{1}{D^2 + 1} \frac{z}{4} = \frac{1}{4} (1 + D^2)^{-1} z$



Equations Reducible To Linear Equations with Constant Coefficients

$$\begin{aligned} &= \frac{1}{4} (1 - D^2 + \dots\dots\dots) z \\ &= \frac{z}{4} \\ &= \frac{x^2}{4} \end{aligned}$$

Hence the complete solution is  $y = C.F + P.I$

or 
$$y = C_1 \cos x^2 + C_2 \sin x^2 + \frac{x^2}{4}$$

**Example 25.** Solve the following differential equation by changing the independent variable  $x$   $\frac{d^2y}{dx^2} + (4x^2 - 1) \frac{dy}{dx} + 4x^3y = 2x^3$  (U.P.T.U. 2006)

**Solution.** The given differential equation may be written as

$$\frac{d^2y}{dx^2} + \left(4x - \frac{1}{x}\right) \frac{dy}{dx} + 4x^2y = 2x^2 \quad (1)$$

Here  $P = 4x - \frac{1}{x}$ ,  $Q = 4x^2$ ,  $R = 2x^2$

on changing the independent variable  $x$  to  $z$ , the equation (1) is transformed as

$$\frac{d^2y}{dz^2} + P_1 \frac{dy}{dz} + Q_1y = R_1 \quad (2)$$

where 
$$Q_1 = \frac{Q}{\left(\frac{dz}{dx}\right)^2} = \frac{4x^2}{\left(\frac{dz}{dx}\right)^2} = 1 \text{ (constant) say}$$

$\Rightarrow \frac{dz}{dx} = 2x$

$\Rightarrow z = x^2$

$\Rightarrow \frac{d^2z}{dx^2} = 2$

$$P_1 = \frac{P \frac{dz}{dx} + \frac{d^2z}{dx^2}}{\left(\frac{dz}{dx}\right)^2} = \frac{8x^2 - 2 + 2}{4x^2} = 2$$

$$\text{and } R_1 = \frac{R}{\left(\frac{dz}{dx}\right)^2} = \frac{2x^2}{4x^2} = \frac{1}{2}$$

Putting the values of  $P_1, Q_1$  &  $R_1$  in (2), we get

$$\frac{d^2y}{dz^2} + 2 \frac{dy}{dz} + y = \frac{1}{2}$$

its Auxiliary equation is  $m^2 + 2m + 1 = 0$

$$\Rightarrow (m + 1)^2 = 0$$

$$\Rightarrow m = -1, -1$$

$$\therefore \text{C.F.} = (C_1 + C_2 z) e^{-z}$$

$$= (C_1 + C_2 x^2) e^{-x^2}$$

$$\begin{aligned} \text{and } P.I &= \frac{1}{D^2 + 2D + 1} \left(\frac{1}{2}\right) \\ &= \frac{1}{2} \frac{1}{D^2 + 2D + 1} e^{0z} \\ &= \frac{1}{2} \frac{1}{(0)^2 + 2.0 + 1} 1 = \frac{1}{2} \end{aligned}$$

$\therefore$  Complete solution is  $y = \text{C.F.} + \text{P.I}$

$$= (C_1 + C_2 x^2) e^{-x^2} + \frac{1}{2}$$

**Example 26.** Solve  $\frac{d^2y}{dx^2} + \cot x \frac{dy}{dx} + 4y \operatorname{cosec}^2 x = 0$

**Solution.** Here  $P = \cot x, Q = 4 \operatorname{cosec}^2 x$  and  $R = 0$  on changing the independent variable  $x$  to  $z$ , the given differential equation transformed to

$$\frac{d^2y}{dz^2} + P_1 \frac{dy}{dz} + Q_1 y = 0$$

Equations Reducible To Linear Equations with Constant Coefficients

where  $P_1 = \frac{P \frac{dz}{dx} + \frac{d^2z}{dx^2}}{\left(\frac{dz}{dx}\right)^2}, Q_1 = \frac{Q}{\left(\frac{dz}{dx}\right)^2}$

**Case I.** Let us take  $P_1 = 0$

$$\frac{P \frac{dz}{dx} + \frac{d^2z}{dx^2}}{\left(\frac{dz}{dx}\right)^2} = 0 \text{ or } P \frac{dz}{dx} + \frac{d^2z}{dx^2} = 0$$

$$\Rightarrow \frac{d^2z}{dx^2} + \cot x \frac{dz}{dx} = 0 \quad (2)$$

put  $\frac{dz}{dx} = v, \frac{d^2z}{dx^2} = \frac{dv}{dx}$

Using these, (2) becomes  $\frac{dv}{dx} + (\cot x) v = 0$

$$\Rightarrow \frac{dv}{v} = -\cot x \, dx$$

$$\Rightarrow \log v = -\log \sin x + \log C = \log C \operatorname{cosec} x$$

$$\Rightarrow v = C \operatorname{cosec} x$$

$$\frac{dz}{dx} = c \operatorname{cosec} x$$

or  $dz = (C \operatorname{cosec} x) \, dx$

$$\Rightarrow z = c \log \tan \frac{x}{2}$$

**Case II.** Now, let us take  $Q_1 = \text{constant}$

$$Q_1 = \frac{Q}{\left(\frac{dz}{dx}\right)^2} = \frac{4 \operatorname{cosec}^2 x}{c^2 \operatorname{cosec}^2 x} = \frac{4}{c^2} \text{ which is constant}$$

Hence the equation (1) reduce to

$$\frac{d^2y}{dz^2} + 0 \frac{dy}{dz} + \frac{4}{c^2} y = 0$$

or  $\frac{d^2y}{dz^2} + \frac{4}{c^2} y = 0 \quad \because P_1 = 0, Q_1 = \frac{4}{c^2}$

$$\Rightarrow \left( D^2 + \frac{4}{c^2} \right) y = 0$$

its auxiliary equation is  $m^2 + \frac{4}{c^2} = 0 \Rightarrow m = \pm i \frac{2}{c}$

$$\therefore \text{C.F} = c_1 \cos \frac{2z}{c} + c_2 \sin \frac{2z}{c}$$

$$\Rightarrow y = c_1 \cos \left( 2 \log \tan \frac{x}{2} \right) + C_2 \sin \left( 2 \log \tan \frac{x}{2} \right) \because z = c \log \tan \frac{x}{2}$$

### SIMULTANEOUS LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS:

In Several applied mathematics problems, there are more than one dependent variables, each of which is a function of one independent variable, usually say time t. The formulation of such problems leads to a system of simultaneous linear differential equation with constant coefficients. Such a system can be solved by the method of elimination. Laplace transform method, using matrices and short cut operator methods.

**Example 27.** Solve  $\frac{dx}{dt} + y = \sin t, \frac{dy}{dt} + x = \cos t$

$$x(0) = 2, y(0) = 0$$

(U.P.T.U. 2004)

**Solution.** We have

$$\frac{dx}{dt} + y = \sin t \tag{1}$$

$$\frac{dy}{dt} + x = \cos t \tag{2}$$

Differentiating (1) w.r.t. 't' we have

$$\frac{d^2x}{dt^2} + \frac{dy}{dt} = \cos t \tag{3}$$

Equations Reducible To Linear Equations with Constant Coefficients

Using (2) in (3) we get

$$\frac{d^2x}{dt^2} - x = 0 \quad \Rightarrow \quad (D^2 - 1)x = 0$$

its auxiliary equation is

$$m^2 - 1 = 0 \quad \Rightarrow \quad m = \pm 1$$

$$\therefore \quad x = C_1 e^t + C_2 e^{-t} \quad (4)$$

$$\Rightarrow \quad \frac{dx}{dt} = C_1 e^t - C_2 e^{-t}$$

putting this value of  $\frac{dx}{dt}$  in (1) we get

$$y = \sin t - C_1 e^t + C_2 e^{-t} \quad (5)$$

Using given conditions

$$\left. \begin{array}{l} \text{from (iv) } C_1 + C_2 = 2 \\ \text{from (v) } -C_1 + C_2 = 0 \end{array} \right\} \Rightarrow C_1 = C_2 = 1$$

putting these values of  $C_1$  and  $C_2$  in (4) & (5) we get

$$x = e^t + e^{-t}$$

$$\text{and} \quad y = \sin t - e^t + e^{-t}$$

is the required solution

**Example 28.** Solve  $\frac{dx}{dt} + 4x + 3y = t$

$$\frac{dy}{dt} + 2x + 5y = e^t \quad (\text{U.P.T.U. 2006})$$

**Solution.** The given equation can be written as

$$(D + 4)x + 3y = t \quad (1)$$

$$2x + (D + 5)y = e^t \quad (2)$$

operating  $(D + 5)$  on equation (1) and multiplied equation (2) by 3, we get

$$(D + 5)(D + 4)x + 3(D + 5)y = (D + 5)t \quad (3)$$

$$6x + 3(D + 5)y = 3e^t \quad (4)$$

Subtracting (4) from (3) we get

$$(D^2 + 9D + 20 - 6) x = 1 + 5t - 3e^t$$

$$\Rightarrow (D^2 + 9D + 14) x = 5t - 3e^t + 1$$

Here auxiliary equation is  $m^2 + 9m + 14 = 0$

$$\Rightarrow (m + 7)(m + 2) = 0$$

$$\Rightarrow m = -7, -2$$

$$\therefore \text{C.F} = C_1 e^{-7t} + C_2 e^{-2t}$$

$$\text{and P.I} = \frac{1}{(D^2 + 9D + 14)} (5t - 3e^t + 1)$$

$$= \frac{5}{14} \left( 1 + \frac{D^2 + 9D}{14} \right)^{-1} t - \frac{3e^t}{(1)^2 + 9(1) + 14} + \frac{1}{(0)^2 + 9(0) + 14} e^{0t}$$

$$= \frac{5}{14} \left( 1 - \frac{9D}{14} \right) t - \frac{3e^t}{24} + \frac{1}{14}$$

$$= \frac{5}{14} \left( t - \frac{9}{14} \right) - \frac{e^t}{8} + \frac{1}{14}$$

$$\Rightarrow \text{P.I} = \frac{5t}{14} - \frac{e^t}{8} - \frac{31}{196}$$

$$\therefore x = C_1 e^{-7t} + C_2 e^{-2t} + \frac{5t}{14} - \frac{e^t}{8} - \frac{31}{196}$$

$$\text{Now } (D + 4) x = -7C_1 e^{-7t} - 2C_2 e^{-2t} + \frac{5}{14} - \frac{e^t}{8} + 4C_1 e^{-7t} + 4C_2 e^{-2t} + \frac{10}{7} t - \frac{e^t}{2} - \frac{31}{49}$$

$$\Rightarrow (D + 4) x = -3C_1 e^{-7t} + 2C_2 e^{-2t} - \frac{5e^t}{8} + \frac{10}{7} t - \frac{27}{98}$$

Using this value in equation (1) we get

$$3y = t + 3C_1 e^{-7t} - 2C_2 e^{-2t} + \frac{5}{8} e^t - \frac{10}{7} t + \frac{27}{98}$$

$$\Rightarrow y = -\frac{1}{7} t + C_1 e^{-7t} - \frac{2}{3} C_2 e^{-2t} + \frac{5}{24} e^t + \frac{9}{98}$$

Thus the required solution is

Equations Reducible To Linear Equations with Constant Coefficients

$$x = C_1 e^{-7t} + C_2 e^{-2t} + \frac{5t}{14} - \frac{e^t}{8} - \frac{31}{196}$$

$$\text{and } y = -\frac{1}{7}t + C_1 e^{-7t} - \frac{2}{3}C_2 e^{-2t} + \frac{5}{24}e^t + \frac{9}{98}$$

**Example 29.** The equation of motion of a particle are given by  $\frac{dx}{dt} + wy = 0$ ,

$\frac{dy}{dt} - wx = 0$ . Find the path of the particle and show that it is a circle.

(U.P.T.U. 2009)

**Solution.** Writing D for  $\frac{d}{dt}$ , the equations are

$$Dx + wy = 0 \tag{1}$$

and  $-wx + Dy = 0 \tag{2}$

Differentiating (1) w.r.t. 't' we have

$$D^2x + wDy = 0 \Rightarrow D^2w + w(wx) = 0 \Rightarrow (D^2 + w^2)x = 0 \text{ using (2)}$$

$$\Rightarrow x = C_1 \cos wt + C_2 \sin wt$$

Putting this value of x in (1) we have  $y = -\frac{1}{w} \frac{d}{dt} (C_1 \cos wt + C_2 \sin wt)$

$$\text{we get } y(t) = C_1 \cos wt + C_2 \sin wt \tag{3}$$

$$\text{and } x(t) = C_2 \cos wt - C_1 \sin wt \tag{4}$$

Squaring (3) and (4) their adding, we get

$$x^2 + y^2 = C_1^2 + C_2^2$$

or  $x^2 + y^2 = R^2$

which is a circle

### Applications to Engineering Problems

#### INTRODUCTION


Differential equations have many numerical applications in Physics, Chemistry, electrical engineering, mechanical engineering, biological sciences, social sciences etc. In this section, we discuss some applications.

### Simple Harmonic Motion

A particle moving in a straight line, is said to execute simple harmonic motion, if its acceleration is always directed towards a fixed point in line and is proportional to the distance of the particle from the fixed point.

Since the acceleration is always directed towards a fixed point, the differential equation of the motion of the particle is given by

$$\frac{d^2x}{dt^2} = -w^2x \quad (1)$$



where  $x$  is the displacement of the particle from a fixed point  $o$  at any time  $t$ .

The solution of (1) is

$$x = C_1 \cos wt + C_2 \sin wt \quad (2)$$

If the particle starts from rest at a point  $A$ , where

$OA = a$  i.e. ( $x = a$ , when  $t = 0$ ) then, from (2), we get

$$C_1 = a$$

Differentiating (2) with respect to  $t$ , we get

$$v = \frac{dx}{dt} = w (-C_1 \sin wt + C_2 \cos wt) \quad (3)$$

Since  $\frac{dx}{dt} = 0$ , at  $t = 0$ , from (3), we get

$$0 = C_2$$

Hence, the displacement of the particle is

$$x = a \cos wt \text{ (a is amplitude)} \quad (4)$$

such that

$$\text{Velocity} = v = \frac{dx}{dt} = -aw \sin wt$$

$$= -w\sqrt{a^2 - x^2} \quad (5)$$



Equations Reducible To Linear Equations with Constant Coefficients

Equation (5) gives the velocity of the particle at any time  $t$ , when its displacement from a fixed point  $O$  is  $x$ . Particle time (time for one complete oscillation) is denoted by  $T$  and is given by  $T = \frac{2\pi}{\omega}$ . The number of complete oscillations per

second is called the frequency of motion and we have  $n = \frac{1}{T} = \frac{\omega}{2\pi}$

In the figure  $O$  is the fixed point

we have  $OA = a$

The acceleration is directed towards  $O$ . The particle moves towards  $O$  from  $A$ . The acceleration gradually decreases and vanishes at  $O$ . At  $O$  particle acquired maximum acceleration. Under retardation the particle further moves towards  $A'$  and comes to rest at  $A'$  such that

$$OA' = OA$$


The point  $O$  is called mean position.

**Example 30.** A point moves in a straight line towards a centre of force  $\mu/(\text{distance})^3$ , starting from rest at a distance  $a$  from the centre of force. Show that the time of reaching a point distance  $b$  from the centre of force is

$$\frac{a}{\sqrt{\mu}} \sqrt{a^2 - b^2} \text{ and that its velocity is } \frac{\sqrt{\mu}}{ab} \sqrt{a^2 - b^2}$$

(U.P.T.U. 2001)

**Solution.** Let  $O$  is the centre of force and let a point moves from  $P$  towards the centre of force  $O$ .



The equation of motion is

$$\frac{d^2x}{dt^2} = \frac{-\mu}{x^3} \tag{1}$$

$$\Rightarrow 2 \frac{dx}{dt} \frac{d^2x}{dt^2} = -2 \frac{\mu}{x^3} \frac{dx}{dt}$$

on integrating, we get

$$\left(\frac{dx}{dt}\right)^2 = -2\mu \frac{1}{-2x^2} + C = \frac{\mu}{x^2} + C$$

$$\Rightarrow \frac{dx}{dt} = \pm \sqrt{\frac{\mu}{x^2} + C} \quad (2)$$

At P,  $\frac{dx}{dt} = 0$  and  $x = a$

$$\Rightarrow 0 = \sqrt{\frac{\mu}{a^2} + C} \Rightarrow C = -\frac{\mu}{a^2}$$

$$\text{From (2), } \frac{dx}{dt} = \pm \sqrt{\frac{\mu}{x^2} - \frac{\mu}{a^2}} = \pm \sqrt{\mu} \frac{\sqrt{a^2 - x^2}}{ax} \quad (3)$$

The velocity at  $x = b$  is

$$v = \pm \sqrt{\mu} \frac{\sqrt{a^2 - b^2}}{ab}$$

or 
$$v = -\sqrt{\mu} \frac{\sqrt{a^2 - b^2}}{ab} \quad (\text{As the point P is moving towards O})$$

$$\text{From (3) } \frac{dx}{dt} = -\sqrt{\mu} \frac{\sqrt{a^2 - x^2}}{ax}$$

$$\Rightarrow dt = -\frac{1}{\sqrt{\mu}} \frac{xa}{\sqrt{a^2 - x^2}} dx$$

on integration, we get

$$t = \frac{a}{\sqrt{\mu}} \sqrt{a^2 - x^2} + C \quad (4)$$

At P,  $t = 0$ ,  $x = a$ , in (4), we get

$$C = 0$$

Putting this value of C in (4), we have

$$t = \frac{a}{\sqrt{\mu}} \sqrt{a^2 - x^2}$$

At  $x = b$

Equations Reducible To Linear Equations with Constant Coefficients

$$t = \frac{a}{\sqrt{\mu}} \sqrt{(a^2 - b^2)}$$

**Vertical Motion In Resisting Medium**

**Example 31.** A particle falls under gravity in a resisting medium whose resistance varies with velocity. Find the relation between distance and velocity if initially the particle starts from rest.

(Bihar P.C.S. 1997, U.P.T.U. 2003)

**Solution.** Let  $m$  be the mass of a particle falls from rest from a fixed point  $O$ . Let  $P$  be the position of a particle such that  $OP = x$ . The forces acting on the particle at  $P$  are:

- (1) The weight  $mg$  of a particle acting vertically downwards.
- (2) The resistance  $mkv$  acting vertically upwards.

Now by Newton's second law of motion the equation of the motion of the body

$$\frac{md^2x}{dt^2} = mg - mkv$$

or 
$$\frac{d^2x}{dt^2} = g - kv$$

or 
$$v \frac{dv}{dx} = g - kv$$


$$\therefore \frac{d^2x}{dt^2} = \frac{dv}{dt} = \frac{dv}{dx} \frac{dx}{dt} = v \frac{dv}{dx}$$

or 
$$\frac{v dv}{g - kv} = dx$$

$$\Rightarrow \frac{1}{k} \left( \frac{-g + kv + g}{g - kv} \right) dv = dx$$

$$\Rightarrow -\frac{dv}{k} + \frac{g}{k(g - kv)} dv = dx$$

Integrating, we get



$$-\frac{v}{k} + \frac{g}{k} \left(-\frac{1}{k}\right) \log(g - kv) = x + C$$

or 
$$-\frac{v}{k} - \frac{g}{k^2} \log(g - kv) = x + C \quad (1)$$

Initially, at point O,  $x = 0, v = 0$

$$\Rightarrow -\frac{g}{k^2} \log g = C$$

putting this value of C in (1) we have

$$-\frac{v}{k} - \frac{g}{k^2} \log(g - kv) = x - \frac{g}{k^2} \log g$$

$$\Rightarrow -\frac{v}{k} - \frac{g}{k^2} \log \frac{g - kv}{g} = x$$

**Example 32.** A 4 kg object falls from rest of time  $t = 0$  in a medium offering a resistance in kg numerically equal to twice its instantaneous velocity in m/sec. Find the velocity and distance travelled at any time  $t > 0$  and also the limiting velocity.

(U.P.T.U. 2007)

**Solution.** Air resistance =  $2v$

$$\therefore \text{Upthrust} = 2 \times 4v = 8v \quad \text{As } m = 4\text{kg}$$

By Newton's second law of motion the equation of motion of body

$$4 \frac{d^2x}{dt^2} = 4g - 8v$$

$$\Rightarrow \frac{d^2x}{dt^2} = g - 2 \frac{dx}{dt}$$

$$\Rightarrow \frac{d^2x}{dt^2} + 2 \frac{dx}{dt} = g \quad (1)$$

$$\text{Let } \frac{dx}{dt} = p \quad \Rightarrow \quad \frac{d^2x}{dt^2} = \frac{dp}{dt}$$

$\therefore$  From (1), we get

Equations Reducible To Linear Equations with Constant Coefficients

$$\frac{dp}{dt} + 2p = g \quad (2)$$

which is linear in p.

its I.F =  $e^{\int 2 dt} = e^{2t}$

So, the solution of equation (2) is

$$p \cdot e^{2t} = \int g e^{2t} dt + C = \frac{g}{2} e^{2t} + C$$

$$\Rightarrow p = \frac{g}{2} + Ce^{-2t}$$

$$\Rightarrow \frac{dx}{dt} = \frac{g}{2} + c e^{-2t} \quad (3)$$

At  $t = 0$ ,  $\frac{dx}{dt} = 0$  which gives  $C = -\frac{g}{2}$

From (3)  $\frac{dx}{dt} = \frac{g}{2} (1 - e^{-2t})$

$$\Rightarrow \text{velocity} = \frac{g}{2} (1 - e^{-2t})$$

Again integrating above equation, we get

$$x = \frac{gt}{2} + \frac{g}{4} e^{-2t} + C_1 \quad (4)$$

At  $t = 0$ ,  $x = 0 \Rightarrow 0 = \frac{g}{4} + C_1 \Rightarrow C_1 = -g/4$

From (4),  $x = \frac{gt}{2} + \frac{g}{4} (e^{-2t} - 1)$

$$\Rightarrow \text{distance} = \frac{gt}{2} + \frac{g}{4} (e^{-2t} - 1)$$

and Limiting velocity =  $\left( \lim_{t \rightarrow \infty} \frac{dx}{dt} \right) = \lim_{t \rightarrow \infty} \frac{g}{2} (1 - e^{-2t})$   
 $= \frac{g}{2}$

**Example 33.** A mass  $M$  suspended from the end of a helical spring is subjected to a periodic force  $f = F \sin wt$  in the direction of its length. The force  $f$  is measured positive vertically downwards and at zero time  $M$  is at rest. If the spring stiffness is  $S$ , prove that the displacement of  $M$  at time  $t$  from the commencement of motion is given by  $x = \frac{F}{M(p^2 - w^2)} \left( \sin wt - \frac{w}{p} \sin pt \right)$ , where  $p^2 = \frac{S}{M}$  and damping effects are neglected.

(U.P.T.U. 2000)

**Solution.** Let  $x$  be the displacement from the equilibrium position at any time  $t$  then the equation of the motion is

$$M \frac{d^2x}{dt^2} = -Sx + F \sin wt$$

or 
$$\frac{d^2x}{dt^2} + \frac{S}{M} x = \frac{F}{M} \sin wt$$

or 
$$\frac{d^2x}{dt^2} + p^2x = \frac{F}{M} \sin wt \tag{1}$$

$$\left( \text{As } \frac{S}{M} = p^2 \right)$$

The A.E. is  $m^2 + p^2 = 0 \Rightarrow m = \pm ip$

C.F. =  $C_1 \cos pt + C_2 \sin pt$

and P.I. =  $\frac{1}{D^2 + p^2} \left( \frac{F}{M} \sin wt \right)$

$$= \frac{F}{M} \frac{1}{-w^2 + p^2} \sin wt$$

$\therefore x = C_1 \cos pt + C_2 \sin pt + \frac{F}{M} \frac{1}{(p^2 - w^2)} \sin wt \tag{2}$

Initially, at  $t = 0, x = 0 \therefore C_1 = 0$

Differentiating equation (2) w.r.t. 't' we get

$$\frac{dx}{dt} = -p C_1 \sin pt + p C_2 \cos pt + \frac{F}{M} \frac{w}{p^2 - w^2} \cos wt$$

Equations Reducible To Linear Equations with Constant Coefficients

At  $t = 0$ ,  $\frac{dx}{dt} = 0$

$$\therefore p C_2 + \frac{F}{M} \frac{w}{p^2 - w^2} = 0$$

$$\text{or } C_2 = -\frac{w}{p} \frac{F}{M(p^2 - w^2)}$$

From (2), we have

$$x = -\frac{w}{p} \frac{F}{M(p^2 - w^2)} \sin pt + \frac{F}{M} \frac{1}{p^2 - w^2} \sin wt$$

$$\text{or } x = \frac{F}{M(p^2 - w^2)} \left( \sin wt - \frac{w}{p} \sin pt \right)$$

### Problems Related to Electric Circuit

There are some formulae which are useful to solve such type of problems

(1)  $i = \frac{dq}{dt}$

(2) Voltage drop across resistance R is  $V_R = Ri$

(3) Voltage drop across inductance L is  $V_L = L \frac{di}{dt}$

(4) Voltage drop across capacitance C is  $V_C = \frac{q}{C}$

### Electro-Mechanical Analogy

The following correspondences between the electrical and mechanical quantities should be kept in mind

Mechanical system	Series Circuit	Parallel circuit
Displacement	Current i	Voltage E
Force or Couple	Voltage E	Current i
Mass m or M.I.	Inductance L	Capacitance C
Damping force	Resistance R	Conductance 1/R
Spring modulus	Elastance 1/C	Susceptance 1/L

**Example 34.** An uncharged condenser of capacity C is charged by applying an e.m.f.  $E \sin \frac{t}{\sqrt{LC}}$  through leads of self-inductance L and negligible resistance. Prove that at time t, the charge on one of the plates is

$$\frac{EC}{2} \left[ \sin \frac{t}{\sqrt{LC}} - \frac{t}{\sqrt{LC}} \cos \frac{t}{\sqrt{LC}} \right]$$

(U.P.T.U. 2003)

**Solution.** If q be the charge on the condenser, the differential equation of the circuit is

$$L \frac{d^2q}{dt^2} + \frac{q}{C} = E_0$$

$$L \frac{d^2q}{dt^2} + \frac{q}{C} = E \sin \frac{t}{\sqrt{LC}} \quad \text{As } E_0 = E \sin \frac{t}{\sqrt{LC}}$$

or 
$$\frac{d^2q}{dt^2} + \frac{1}{LC} q = \frac{E}{L} \sin \frac{t}{\sqrt{LC}} \quad (1)$$

Here auxiliary equation is  $m^2 + \frac{1}{LC} = 0$

$$\Rightarrow m = \pm \frac{i}{\sqrt{LC}}$$

$$\therefore C.F = C_1 \cos \frac{1}{\sqrt{LC}} t + C_2 \sin \frac{1}{\sqrt{LC}} t$$



Equations Reducible To Linear Equations with Constant Coefficients

$$\begin{aligned}
 P.I &= \frac{1}{\left(D^2 + \frac{1}{LC}\right)} \frac{E}{L} \sin \frac{t}{\sqrt{LC}} \quad (\text{Case of failure}) \\
 &= \frac{E}{L} \left( -\frac{t}{2\sqrt{LC}} \cos \frac{t}{\sqrt{LC}} \right) \quad \because \frac{1}{D^2 + a^2} \sin ax = \frac{-x}{2a} \cos ax \\
 &= \frac{E}{L} \left( \frac{-t\sqrt{LC}}{2} \cos \frac{t}{\sqrt{LC}} \right) \\
 &= -\frac{Et}{2} \sqrt{\left(\frac{C}{L}\right)} \cos \frac{t}{\sqrt{LC}}
 \end{aligned}$$

Therefore, the solution of the equation is

$$q = C_1 \cos \frac{t}{\sqrt{LC}} + C_2 \sin \frac{t}{\sqrt{LC}} - \frac{Et}{2} \sqrt{\left(\frac{C}{L}\right)} \cos \frac{t}{\sqrt{LC}} \quad (2)$$

At  $t = 0, q = 0 \therefore C_1 = 0$

Differentiating equation (2) w.r.t. "t" we get

$$\frac{dq}{dt} = -\frac{C_1}{\sqrt{LC}} \sin \frac{t}{\sqrt{LC}} + \frac{C_2}{\sqrt{LC}} \cos \frac{t}{\sqrt{LC}} + \frac{Et}{2} \sqrt{\left(\frac{C}{L}\right)} \sin \left(\frac{t}{\sqrt{LC}}\right) \frac{1}{\sqrt{LC}} - \frac{E}{2} \sqrt{\frac{C}{L}} \cos \frac{t}{\sqrt{LC}}$$

Initially  $\frac{dq}{dt} = 0$ , when  $t = 0$

$$\therefore \frac{C_2}{\sqrt{LC}} - \frac{E}{2} \sqrt{\frac{C}{L}} = 0 \quad \Rightarrow \quad C_2 = \frac{EC}{2}$$

From equation (2) we get

$$q = \frac{EC}{2} \sin \frac{t}{\sqrt{LC}} - \frac{Et}{2} \sqrt{\left(\frac{C}{L}\right)} \cos \frac{t}{\sqrt{LC}}$$

or 
$$q = \frac{EC}{2} \left( \sin \frac{t}{\sqrt{LC}} - \frac{t}{\sqrt{LC}} \cos \frac{t}{\sqrt{LC}} \right)$$

**Example 35.** The equation of electromotive force in terms of current  $i$  for an electrical circuit having resistant  $R$  and a condenser of capacity  $C$ , in series is

$$E = Ri + \int \frac{i}{C} dt. \text{ Find the current } i \text{ at any time } t, \text{ when } E = E_0 \sin wt$$

(U.P.T.U. 2006)

**Solution.** The given equation is

$$Ri + \int \frac{i}{C} dt = E_0 \sin wt \text{ as } E = E_0 \sin wt$$

· Differentiating w.r.t. 't', we get

$$R \frac{di}{dt} + \frac{i}{C} = E_0 w \cos wt$$

$$\Rightarrow \frac{di}{dt} + \frac{i}{RC} = \frac{E_0 w}{R} \cos wt \quad (1)$$

which is a linear differential equation

its 
$$I.F = e^{\int \frac{1}{RC} dt}$$

$$= e^{\frac{t}{RC}}$$

The solution of (1) is

$$i \cdot e^{\frac{t}{RC}} = \int \frac{E_0 w}{R} \cos wt \cdot e^{t/RC} dt + C_1$$

$$= \frac{E_0 w}{R} \frac{e^{t/RC}}{\frac{1}{R^2 C^2} + w^2} \left[ \frac{1}{RC} \cos wt + w \sin wt \right] + C_1$$

∴ 
$$\int e^{ax} \cos bx dx = \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx)$$

or 
$$i = \frac{wE_0 RC^2}{1 + w^2 R^2 C^2} \left[ \frac{1}{RC} \cos wt + w \sin wt \right] + ke^{-t/RC}$$

**Example 36.** The damped LCR circuit is governed by the equation

$$L \frac{d^2Q}{dt^2} + R \frac{dQ}{dt} + \frac{1}{C} Q = 0$$

Equations Reducible To Linear Equations with Constant Coefficients

where  $L, C, R$  are positive constants. Find the conditions under which the circuit is overdamped, underdamped and critically damped. Find also the critical resistance.

(U.P.T.U. 2005)

**Solution.** The given equation is

$$L \frac{d^2Q}{dt^2} + R \frac{dQ}{dt} + \frac{1}{C} Q = 0$$

or 
$$\frac{d^2Q}{dt^2} + 2k \frac{dQ}{dt} + w^2 Q = 0 \quad (1)$$

Where  $2k = \frac{R}{L}$  and  $w^2 = \frac{1}{LC}$

Here auxiliary equation is

$$m^2 + 2km + w^2 = 0$$

$\Rightarrow m = -k \pm \sqrt{k^2 - w^2} \quad (2)$

**Case I.** when  $k < w$  i.e.  $\frac{R}{2L} < \frac{1}{\sqrt{LC}}$ , the roots of A.E. given by (2) are imaginary.

The general solution of (1) is

$$Q = e^{-kt} (C_1 \cos \sqrt{w^2 - k^2} t + C_2 \sin \sqrt{w^2 - k^2} t)$$

where  $C_1$  and  $C_2$  being arbitrary constants.

Time period =  $\frac{2\pi}{\sqrt{w^2 - k^2}}$  which is greater than  $\frac{2\pi}{w}$

Thus the effect of damping increases the period of oscillation and motion ultimately dies away. In this condition when  $\frac{R}{2L} < \frac{1}{\sqrt{LC}}$  the circuit is under damped.

**Case II.** When  $k = w$ , then roots of A.E. (2) are equal, each being equal to  $-k$ . The general solution of (1) is

$$Q = (C_1 + C_2 t) e^{-kt}$$

In this case charge  $Q$  is always positive and decreases to zero as  $t \rightarrow \infty$ . In this case circuit is called critically damped and the resistance  $R$  is called critical resistance.

Thus  $k = w \Rightarrow \frac{R}{2L} = \frac{1}{\sqrt{LC}}$

$\Rightarrow R = 2\sqrt{\frac{L}{C}}$

which is required critical resistance.

**Case III.** when  $k > w$ , the roots of A.E. are real and unequal.


Also, the roots

$$m = -k + \sqrt{k^2 - w^2} \text{ and } m = -k - \sqrt{k^2 - w^2}$$

are both negative. The general solution of (1) is

$$Q = C_1 e^{\{-k + \sqrt{k^2 - w^2}\}t} + C_2 e^{\{-k - \sqrt{k^2 - w^2}\}t}$$

In this case also charge  $Q$  is positive and decreases to zero as  $t \rightarrow \infty$ , since exponential terms having negative powers approach to zero. In this case the circuit is called overdamped.



**Example 37.** The voltage  $V$  and the current  $i$  at a distance  $x$  from the sending end of the transmission line satisfying the equations.

$$-\frac{dv}{dx} = Ri, \quad -\frac{di}{dx} = GV$$

where  $R$  and  $G$  are constants. If  $V = V_0$  at the sending end ( $x = 0$ ) and  $V = 0$  at receiving end ( $x = l$ ), show that

$$V = V_0 \left\{ \frac{\sinh n(l-x)}{\sinh nl} \right\}$$

Equations Reducible To Linear Equations with Constant Coefficients

(U.P.T.U. 2006)

**Solution.** We have  $-\frac{dV}{dx} = Ri$  (1)

and  $-\frac{di}{dx} = GV$  (2)

when  $x = 0, V = V_0$ , when  $x = l, V = 0$

From (1) and (2), we have

$$-\frac{d}{dx} \left( -\frac{dV}{dx} \frac{1}{R} \right) = GV$$

$$\Rightarrow \frac{d^2V}{dx^2} = RGV$$

$$\Rightarrow \frac{d^2V}{dx^2} - (RG) V = 0$$

or  $(D^2 - RG) V = 0, D \equiv \frac{d}{dx}$  (3)

Here auxiliary equation is  $m^2 - RG = 0$

$$\Rightarrow m = \pm n, n^2 = RG$$

The solution of (3) is  $V = C_1 e^{nx} + C_2 e^{-nx}$  (4)

where  $C_1$  and  $C_2$  are arbitrary constants.

putting  $x = 0$  and  $V = V_0$  is (4), we get

$$V_0 = C_1 + C_2$$
 (5)

Again putting  $x = l$  and  $V = 0$  is (4), we get

$$0 = C_1 e^{nl} + C_2 e^{-nl}$$
 (5)

Solving equations (5) and (6), we have

$$C_1 = \frac{V_0}{1 - e^{2nl}}, C_2 = \frac{-V_0 e^{2nl}}{1 - e^{2nl}}$$

Substituting the values of  $C_1$  and  $C_2$  in (4), we get

$$V = \frac{V_0}{1 - e^{2nl}} e^{nx} - \frac{V_0 e^{2nl}}{1 - e^{2nl}} e^{-nx}$$

$$= \frac{V_0 (e^{nx} - e^{2nl - nx})}{1 - e^{2nl}}$$

or 
$$V = \frac{V_0 \{e^{(nl - nx)} - e^{-(nl - nx)}\}}{e^{nl} - e^{-nl}} = V_0 \left\{ \frac{\sin h n(l - x)}{\sin h nl} \right\}$$

**Example 38.** An inductance of 2 henries and a resistance of 20 ohms are connected in series with an emf E volts. If the current is zero when  $t = 0$ , find the current at the end of 0.01 sec if  $E = 100$  volts, using the following differential equation.

(U.P.T.U. 2008)


$$L \frac{di}{dt} + iR = E$$

**Solution.** we have  $L \frac{di}{dt} + iR = E$

or 
$$\frac{di}{dt} + \frac{R}{L} i = \frac{E}{L} \quad (1)$$

Equation (1) is linear differential equation of first order.

$$I.F = e^{\int \frac{R}{L} dt} = e^{(R/L)t}$$



∴ Solution of (1) is

$$i e^{(R/L)t} = \int \frac{E}{L} e^{(R/L)t} dt + C$$

where C is an arbitrary constant

or 
$$i e^{(R/L)t} = \frac{E}{R} e^{(R/L)t} + C$$

Equations Reducible To Linear Equations with Constant Coefficients

$$\Rightarrow i = \frac{E}{R} + C e^{-(R/L)t} \quad (2)$$

Initially  $i = 0$ , when  $t = 0$   $\therefore$  From (2), we have

$$C = -\frac{E}{R}$$

$$\therefore \text{From (2), we have } i = \frac{E}{R} [1 - e^{-(R/L)t}] \quad (3)$$

on putting  $E = 100$  volts,  $R = 20$  ohms and  $L = 2$  henries in (3) we have

$$i = \frac{100}{5} \left[ 1 - e^{-\frac{20}{2}t} \right] = 5 (1 - e^{-10t})$$

$$\begin{aligned} \text{At } t = 0.01 \text{ sec, } i &= 5 (1 - e^{-0.1}) \\ &= 0.475 \text{ amp (approximately)} \end{aligned}$$

**Example 39.** In an LCR circuit, the charge  $q$  on a plate of a condenser is given by  $L \frac{d^2q}{dt^2} + R \frac{dq}{dt} + \frac{q}{C} = E \sin pt$ . The circuit is tuned to resonance so that  $p^2 = \frac{1}{LC}$ , if initially the current  $i$  and the charge  $q$  be zero, show that for small values of  $\frac{R}{L}$ , the current in the circuit at time  $t$  is given by  $\frac{Et}{2L} \sin pt$

(U.P.T.U. 2004) (C.O.)

**Solution.** The given differential equation is

$$L \frac{d^2q}{dt^2} + R \frac{dq}{dt} + \frac{q}{C} = E \sin pt \quad (1)$$

Here A.E. is  $Lm^2 + Rm + \frac{1}{C} = 0$

$$\Rightarrow m = \frac{-R + \sqrt{R^2 - (4L/C)}}{2L} = -\frac{R}{2L} \pm \frac{1}{2} \sqrt{\frac{R^2}{L^2} - \frac{4}{CL}}$$

$$\Rightarrow m = -\frac{R}{2L} \pm \frac{1}{2} \sqrt{-\frac{4}{CL}} \quad \left( \text{neglected } \frac{R^2}{L^2} \text{ as } \frac{R}{LC} \text{ is small} \right)$$

$$\Rightarrow m = -\frac{R}{2L} \pm \frac{i}{\sqrt{CL}} = -\frac{R}{2L} \pm ip \quad \left( \text{since } p^2 = \frac{1}{LC} \text{ given} \right)$$

$$\therefore \text{C.F} = e^{-Rt/2L} (C_1 \cos pt + C_2 \sin pt)$$

$$\text{But } e^{-Rt/2L} = 1 - \frac{Rt}{2L} + \frac{1}{L^2} \frac{R^2 t^2}{4L^2} - \dots$$

$$= 1 - \frac{Rt}{2L} \quad \text{neglecting } \frac{R^2}{L^2} \text{ etc}$$

$$\therefore \text{C.F.} = \left( 1 - \frac{Rt}{2L} \right) (C_1 \cos pt + C_2 \sin pt)$$

where  $C_1$  and  $C_2$  are arbitrary constants

$$\text{P.I} = \frac{1}{LD^2 + RD + \frac{1}{C}} E \sin pt \quad \text{where } D \equiv \frac{d}{dt}$$

$$= E \frac{1}{L(-p^2) + RD + \frac{1}{C}} \sin pt$$

$$= E \frac{1}{RD} \sin pt \quad \text{since } p^2 = \frac{1}{LC}$$

$$= \frac{E}{R} \int \sin pt \, dt = \frac{-E}{pR} \cos pt$$

Hence the general solution of (1) is given by

$$q = \left( 1 - \frac{Rt}{2L} \right) (C_1 \cos pt + C_2 \sin pt) - \frac{E}{pR} \cos pt \quad (2)$$

Differentiating (2) w.r.t. 't' we have

$$i = \frac{dq}{dt} = \left( 1 - \frac{Rt}{2L} \right) (-pC_1 \sin pt + pC_2 \cos pt) - \frac{R}{2L} (C_1 \cos pt + C_2 \sin pt) + \frac{E}{R} \sin pt \quad (3)$$

Initially given that  $t = 0, q = 0$

$\therefore$  (2) gives

$$0 = C_1 - \frac{E}{pR} \quad \Rightarrow \quad C_1 = \frac{E}{pR}$$

and (3) gives



Equations Reducible To Linear Equations with Constant Coefficients

$$0 = pC_2 - \frac{RC_1}{2L} \Rightarrow C_2 = \frac{RC_1}{2pL} = \frac{E}{2Lp^2}$$

Now putting values of  $C_1$  and  $C_2$  in (3), the current  $i$  in the circuit at any time  $t$  is given by

$$i = \left(1 - \frac{Rt}{2L}\right) \left(-\frac{E}{R} \sin pt + \frac{E}{2pL} \cos pt\right) - \frac{R}{2L} \left(\frac{E}{pR} \cos pt + \frac{E}{2Lp^2} \sin pt\right) + \frac{E}{R} \sin pt$$

$$= \frac{Et}{2L} \sin pt - \frac{ERt}{4pL^2} \cos pt - \frac{ER}{4L^2p^2} \sin pt$$

$= \frac{Et}{2L} \sin pt$  since  $\frac{R}{L}$  is small, also  $\frac{R}{L^2} = \frac{1}{R} \left(\frac{R}{L}\right)^2$ , so neglecting second and third terms

**BEAM**

A bar whose length is much greater than its cross-section and its thickness is called a beam

**Cantilever:** If one end of a beam is fixed and the other end is loaded, it is called a cantilever.

**Bending of Beam:** Let a beam be fixed at one end and the other end is loaded. Then the upper surface is elongated and therefore under tension and the lower surface is shortened so under compression.

**Bending Moment:** Whenever a beam is loaded it deflects from its original position. If  $M$  is the bending moment of the forces acting on it, then

$$M = \frac{EI}{R} \tag{1}$$

where  $E$  = Modulus of elasticity of the beam

$I$  = Moment of inertia of the cross-section of beam about neutral axis

$R$  = Radius of curvature of the curved beam

$$R = \frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{3/2}}{\frac{d^2y}{dx^2}} = \frac{1}{\frac{d^2y}{dx^2}} \text{ neglecting } \frac{dy}{dx}$$


Thus equation (1) becomes  $M = EI \frac{d^2y}{dx^2}$

**Example 40.** A beam of length  $l$  is clamped horizontally at its end  $x = 0$  and is free at the end  $x = l$ . A point load  $W$  is applied at the end  $x = l$ , in addition of a uniform load  $w$  per unit length from  $x = 0$  to  $x = \frac{l}{2}$ . Find the deflection at any point.

(U.P.T.U. 2002)

**Solution.** Let OA be a beam, clamped (i.e. fixed) at one end O and free at end A. Let B be its mid point. The weight  $\frac{wl}{2}$  of the beam OB acts at C (the mid point of OB). The weight  $w$  acts at A.

Let R be the force acting at O. The directions of all the force acting on the beam are as shown in figure.



∴ From balance equation, we have

$$R = W + \frac{wl}{2} \tag{1}$$

Now we choose a random axis NN', if  $(x, y)$  are the co-ordinates of N, then taking moments about N, we get

$$EI \frac{d^2y}{dx^2} = -Rx + \frac{Wl}{2} \left( x - \frac{l}{4} \right)$$

or 
$$EI \frac{d^2y}{dx^2} = - \left( W + \frac{wl}{2} \right) x + \frac{wl}{2} x - \frac{wl^2}{8} \text{ using (1) for R}$$

Equations Reducible To Linear Equations with Constant Coefficients

or 
$$EI \frac{d^2y}{dx^2} = -\frac{Wl^2}{8} - Wx$$

Integrating both sides w.r.t.  $x$ , we get

$$\text{E.I. } \frac{dy}{dx} = -\frac{wl^2}{8} x - \frac{1}{2} Wx^2 + C_1 \quad (2)$$

Applying boundary conditions at the fixed (i.e. clamped) end O i.e., at  $x = 0$ ,  $dy/dx = 0$ , we get from (2),  $C_1 = 0$

$\therefore$  (2) becomes, 
$$EI \frac{dy}{dx} = -\frac{wl^2}{8} x - \frac{1}{2} Wx^2$$

Again integrating  $EIy = -\frac{wl^2x^2}{16} - \frac{1}{6} Wx^3 + C_2$  (3)

Again boundary conditions, at  $x = 0$ ,  $y = 0$  gives  $C_2 = 0$

$\therefore$  (3) becomes,

$$EIy = \frac{-wl^2x^2}{16} - \frac{Wx^3}{6}$$

or 
$$y = -\frac{1}{EI} \left( \frac{wl^2x^2}{16} + \frac{Wx^3}{6} \right)$$

which gives the deflection at any point.

**Example 41.** The deflection of a strut of length  $l$  with one end ( $x = 0$ ) built in and the other end supported and subjected to end thrust  $P$ , satisfies the equation

$$\frac{d^2y}{dx^2} + a^2y = \frac{a^2R}{P} (l - x)$$

Prove that the deflection curve  $y = \frac{R}{P} \left( \frac{\sin ax}{a} - l \cos ax + l - x \right)$

where  $a l = \tan a l$

(U.P.T.U. 2001)


**Solution.** we have

$$\frac{d^2y}{dx^2} + a^2y = \frac{a^2R}{P} (l - x) \quad (1)$$

its auxiliary equation is  $m^2 + a^2 = 0 \Rightarrow m = \pm ai$

$\therefore$  C.F. =  $C_1 \cos ax + C_2 \sin ax$

where  $C_1$  and  $C_2$  are arbitrary constants.



$$\begin{aligned}
 P.I &= \frac{1}{D^2 + a^2} \frac{a^2 R}{P} (l - x) \\
 &= \frac{R}{P} \left( 1 + \frac{D^2}{a^2} \right) (l - x) \\
 &= \frac{R}{P} \left( 1 - \frac{D^2}{a^2} \right) (l - x) \\
 &= \frac{R}{P} (l - x)
 \end{aligned}$$

$\therefore$  The general solution is

$$y = C_1 \cos ax + C_2 \sin ax + \frac{R}{P} (l - x) \quad (2)$$

Differentiating (2) w.r.t.  $x$ , we get

$$\frac{dy}{dx} = -C_1 a \sin ax + C_2 a \cos ax - \frac{R}{P} \quad (3)$$

The end O of the strut is built in, so at  $x = 0$ ,  $y = dy/dx = 0$

$\therefore$  (2) gives

$$0 = C_1 + \frac{Rl}{P} \Rightarrow C_1 = -\frac{Rl}{P}$$

and (3) gives  $0 = C_2 a - \frac{R}{P} \Rightarrow C_2 = \frac{R}{aP}$

Equations Reducible To Linear Equations with Constant Coefficients

Putting for  $C_1$  and  $C_2$  is (2)

$$y = -\frac{Rl}{P} \cos ax + \frac{R}{aP} \sin ax + \frac{R}{P} (l - x)$$

$$\Rightarrow y = \frac{R}{P} \left( \frac{\sin ax}{a} - l \cos ax + l - x \right) \quad (4)$$

Also, the end A of the strut is supported, so at  $x = l, y = 0$ , so that (4) becomes

$$0 = \frac{R}{P} \left( \frac{\sin al}{a} - l \cos al + l - l \right)$$

or  $\frac{\sin al}{a} = l \cos al$

or  $al = \tan al$

Hence the required equation of deflection curve is given by (4) where  $al = \tan al$ .

### EXERCISE

**Solve the following differential equations**

1. (i)  $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} + y = \sin(\log x^2)$

Ans.  $y = C_1 \cos(\log x) + C_2 \sin(\log x) - \frac{1}{3} \sin(\log x^2)$

(ii)  $(1+x)^2 \frac{d^2y}{dx^2} + (1+x) \frac{dy}{dx} + y = \sin 2 \{\log(1+x)\}$  (I.A.S. 2003)

Ans.  $y = C_1 \cos \{\log(1+x)\} + C_2 \sin \{\log(1+x)\} - \frac{1}{3} \sin 2 \{\log(1+x)\}$

2.  $x^2 \frac{d^3y}{dx^3} + 3x \frac{d^2y}{dx^2} + \frac{dy}{dx} = x^2 \log x$

Ans.  $y = C_1 + C_2 \log x + C_3 (\log x)^2 + \frac{x^3}{27} (\log x - 1)$

3.  $x^4 \frac{d^4y}{dx^4} + 6x^3 \frac{d^3y}{dx^3} + 4x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} - 4y = 2 \cos(\log x)$

Ans.  $y = C_1 x^2 + C_2 x^{-2} + C_3 \cos \log x + C_4 \sin \log x - \frac{1}{5} \log x \sin \log x$

4.  $x^2 \frac{d^2y}{dx^2} - x \frac{dy}{dx} - 3y = x^2 \log x$  (I.A.S. 2001)

Ans.  $y = C_1 x^3 + C_2 x^{-1} - \frac{1}{3} x^3 \left( \log x + \frac{2}{3} \right)$

5.  $\frac{d^2y}{dx^2} + y = \operatorname{cosec} x$  by the method of variation of Parameters

Ans.  $y = C_1 \cos x + C_2 \sin x - x \cos x + \sin x \cdot \log \sin x$

6.  $\frac{d^2y}{dx^2} + (1 - \cot x) \frac{dy}{dx} - y \cot x = \sin^2 x$  by variation of parameters.

(U.P.T.U. Special Exam 2001)

Ans.  $y = C_1 (\sin x - \cos x) + C_2 e^{-x} - \frac{1}{10} \sin 2x + \frac{1}{5} \cos 2x$

7.  $x^2 \frac{d^2y}{dx^2} - (x^2 + 2x) \frac{dy}{dx} + (x + 2)y = x^3 e^x$  of which  $y = x$  is a solution.

Ans.  $y = x (C_1 + C_2 e^x + x e^x)$

8.  $(x \sin x + \cos x) \frac{d^2y}{dx^2} - x \cos x \frac{dy}{dx} + y \cos x = 0$  of which  $y = x$  is a solution.

Ans.  $y = -C_1 \cos x + C_2 x$

9.  $\frac{d^2y}{dx^2} - 2 \tan x \frac{dy}{dx} + 5y = e^x \cdot \sec x$  by reducing normal form.

Ans.  $y = \sec x \left( C_1 \cos \sqrt{6} x + C_2 \sin \sqrt{6} x + \frac{1}{7} e^x \right)$

10.  $(1 + x^2)^2 \frac{d^2y}{dx^2} + 2x(1 + x^2) \frac{dy}{dx} + 4y = 0$  by changing independent variable.

Ans.  $y = C_1 \cos (2 \tan^{-1} x) + C_2 \sin (2 \tan^{-1} x)$

11.  $x \frac{d^2y}{dx^2} - \frac{dy}{dx} + 4x^3 y = 8x^3 \sin x^2$  by changing independent variable.

Ans.  $y = C_1 \cos x^2 + C_2 \sin x^2 - x^2 \cos x^2$

12.  $\frac{d^2y}{dx^2} + (3 \sin x - \cot x) \frac{dy}{dx} + 2y \sin^2 x = e^{-\cos x} \cdot \sin^2 x$  by changing independent variable.

Equations Reducible To Linear Equations with Constant Coefficients

Ans.  $y = C_1 e^{\cos x} + C_2 e^{2 \cos x} + \frac{1}{6} e^{-\cos x}$

13.  $\frac{d^2x}{dt^2} + \frac{dy}{dt} + 3x = e^{-t}, \frac{d^2y}{dt^2} - 4 \frac{dx}{dt} + 3y = \sin 2t$

(U.P.T.U. 2007)

Ans.  $x = -\frac{C_1}{2} \sin 3t + \frac{C_2}{2} \cos 3t + \frac{C_3}{2} \sin t - \frac{C_4}{2} \cos t + \frac{1}{5} e^{-t} + \frac{1}{5} \sin 2t + A$  where  $A = -\frac{C_5}{4}$

$$y = C_1 \cos 3t + C_2 \sin 3t + C_3 \cos t + C_4 \sin t + \frac{1}{5} \left( -e^{-t} + \frac{1}{3} \sin 2t \right)$$

14.  $\frac{dx}{dt} + \frac{dy}{dt} - 2y = 2 \cos t - 7 \sin t$

$$\frac{dx}{dt} - \frac{dy}{dt} + 2x = 4 \cos t - 3 \sin t$$

(U.P.T.U. 2001)

Ans.  $x = -\left(1 - \sqrt{2}\right) C_1 e^{\sqrt{2}t} - \left(1 + \sqrt{2}\right) C_2 e^{-\sqrt{2}t} + 3 \cos t + C_3$

$$y = C_1 e^{\sqrt{2}t} + C_2 e^{-\sqrt{2}t} + 2 \sin t$$

15.  $\frac{d^2x}{dt^2} - 3x - 4y = 0, \frac{d^2y}{dt^2} + x + y = 0$

(U.P.T.U. 2005)

Ans.  $x = -[(2 C_1 + 2 C_2 + 2 C_2 t) e^t + 2 C_3 - 2 C_4 + 2 C_4 t] e^{-t}$

$$y = (C_1 + C_2 t) e^t + (C_3 + C_4 t) e^{-t}$$

16. Solve  $\frac{dx}{dt} + 5x - 2y = t$

$$\frac{dy}{dt} + 2x + y = 0$$

Also show that  $x = y = 0$  when  $t = 0$  for some definite values of constants

(U.P.T.U. 2008)

Ans.  $x = C_1 e^{-3t} + C_2 t e^{-3t} - \frac{1}{2} C_2 e^{-3t} - \frac{1}{2} C_1 e^{-3t} + \frac{t}{9} + \frac{1}{27}, y = (C_1 + C_2 t) e^{-3t} - \frac{2t}{9} + \frac{4}{27}$

17. A particle moving in a straight line with S.H.M. has velocities  $v_1$  and  $v_2$  when its distances from the centre are  $x_1$  and  $x_2$  respectively. Show that the period of motion is  $2\pi \sqrt{\frac{x_1^2 - x_2^2}{v_2^2 - v_1^2}}$  and its amplitude is  $\sqrt{\frac{(v_1^2 x_2^2 - v_2^2 x_1^2)}{(v_1^2 - v_2^2)}}$   
(Bihar P.C.S. 2005)
18. A particle is performing a simple harmonic motion of period  $T$  about a centre  $O$  and it passes through a point  $P$ , where  $OP = b$  with velocity  $v$  in the direction  $OP$ . Prove that the time which elapses before it returns to  $P$  is  $\frac{T}{\pi} \tan^{-1} \left( \frac{vT}{2\pi b} \right)$ .  
(I.A.S. 2007)
19. A particle of mass  $m$  is projected vertically under gravity, the resistance of the air being  $mk$  times the velocity. Show that the greatest height attained by the particle is  $\frac{V^2}{g} [\lambda - \log(1 + \lambda)]$  where  $V$  is terminal velocity of the particle and  $\lambda V$  is the initial velocity.  
(U.P.P.C.S. 2004)
20. If  $u$  and  $V$  are the velocity of projection and the terminal velocity respectively of a particle rising vertically against a resistance varying as the square of the velocity. Prove that the time taken by the particle to reach the highest point is  $\frac{V}{g} \tan^{-1} \left( \frac{u}{V} \right)$ .  
(I.A.S. 2006)
21. In the LCR circuit, the charge  $q$  on a plate of a condenser is given by  $L \frac{d^2q}{dt^2} + R \frac{dq}{dt} + \frac{q}{C} = E \sin pt$ . The circuit is tuned to resonance so that  $p^2 = \frac{1}{LC}$ . If initially the current  $i$  and the charge  $q$  be zero, show that for small value of  $\frac{R}{L}$ , the current in the circuit at time  $t$  is given by  $\left( \frac{Et}{2L} \right) \sin pt$ .  
(U.P.T.U. 2004)



## Objective Type of Questions

Choose a correct answer from the four answers given in each of the following questions.

1. The solution of the differential equation

$$x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} + y = \sin(\log x^2) \text{ is}$$

(a)  $C_1 \cos(\log x) + C_2 \sin(\log x) + \frac{1}{3} \sin(\log x^2)$

(b)  $C_1 \cos(\log x) + C_2 \sin(\log x) - \frac{1}{3} \sin(\log x^2)$

(c)  $C_1 \cos(\log x) + C_2 \sin(\log x) - \frac{1}{3} \sin(\log x)$

(d)  $C_1 \cos(\log x^2) + C_2 \sin(\log x^2) - \frac{1}{3} \sin(\log x)$

Ans. (b)

2. A particular integral of the differential equation

$$x^2 \frac{d^3y}{dx^3} + 3x \frac{d^2y}{dx^2} + \frac{dy}{dx} = x^2 \log x$$

(a)  $\frac{x^2}{27} (\log x - 1)$       (b)  $\frac{x^2}{27} (\log x + 1)$

(c)  $\frac{x^3}{27} (\log x - 1)$       (d)  $\frac{x^3}{27} (\log x + 1)$

Ans. (c)

3. A particular integral of  $x^2 \frac{d^2y}{dx^2} + 4x \frac{dy}{dx} + 2y = e^x$  is

(a)  $\frac{1}{x^2} e^x$       (b)  $\frac{1}{x} e^x$

(c)  $\frac{1}{x^2} e^{2x}$       (d)  $\frac{1}{x^2} e^{3x}$

Ans. (a)

4. The C.F. of the differential equation  $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} - y = x^3 e^x$

(a)  $C_1 x^2 + \frac{C_2}{x^2}$                       (b)  $C_1 x^2 + \frac{C_2}{x}$

(c)  $C_1 x + \frac{C_2}{x^2}$                       (d)  $C_1 x + \frac{C_2}{x}$

Ans. (d)

5. On putting  $x = e^z$ , the transformed differential equation of

$$x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} + y = x \text{ is}$$

(a)  $\frac{d^2y}{dz^2} = e^z$                       (b)  $\frac{d^2y}{dz^2} + y = e^z$

(c)  $\frac{d^2y}{dz^2} + y = e^{2z}$                       (d)  $\frac{d^2y}{dz^2} + y = z$

Ans. (b)

6. The equation of motion of a particle are given by simultaneous differential

equations  $\frac{dx}{dt} + wy = 0, \frac{dy}{dt} - wx = 0$ , Then the path of the particle is

(a) Straight line                      (b) Circle

(c) Ellipse                      (d) Parabola

Ans. (b)

7. A Particular integral of  $\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = 0$  is  $y = e^{mx}$  if

(a)  $m^2 + Pm + Q = 0$                       (b)  $m^2 - Pm + Q = 0$

(c)  $m + Pm^2 + Q = 0$                       (d)  $m^2 + Pm - Q = 0$

Ans. (a)

8.  $y = e^x$  is a part of C.F. of differential equation  $\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = 0$  if

(a)  $1 + P + Q = 0$                       (b)  $1 - P + Q = 0$

(c)  $P + Qx = 0$                       (d)  $P - Qx = 0$

Ans. (a)

Equations Reducible To Linear Equations with Constant Coefficients

9. In a differential equation

$$x^2 \frac{d^2y}{dx^2} - (x^2 + 2x) \frac{dy}{dx} + (x + 2)y = x^3e^x, y = x \text{ is a part of C.F. because}$$

- (a)  $P - Qx = 0$                       (b)  $P + Qx = 0$   
(c)  $1 + P + Q = 0$                 (d)  $1 - P - Q = 0$

Ans. (b)

10. The solution for  $\frac{d^2y}{dx^2} - 4x \frac{dy}{dx} + (4x^2 - 2)y = 0$ , given that  $y = e^{x^2}$  is an integral included in the complementary function is

- (a)  $y = (C_1 x + C_2)$                 (b)  $e^x (C_1 x + C_2)$   
(c)  $e^{x^2} C_1 x$                         (d)  $e^{x^2} (C_1 x + C_2)$

Ans. (d)

11. A resistance of 100 ohms, an inductance of 0.5 henry are connected in series with a battery of 20 volts. The current in the circuit is

- (a)  $i = \frac{1}{5} (1 - e^{-20t})$                 (b)  $i = \frac{1}{5} (1 - e^{20t})$   
(c)  $i = \frac{1}{5} (1 - e^{-200t})$                 (d)  $i = \frac{1}{5} (1 - e^{200t})$

Ans. (c)

12. The solution of the differential equation  $L \frac{di}{dt} + Ri = E_0 \sin wt$  is

- (a)  $i = \frac{E_0 L}{\sqrt{R^2 + L^2 w^2}} \sin \left( wt + \tan^{-1} \frac{wL}{R} \right)$   
(b)  $i = \frac{E_0 L}{\sqrt{R^2 + w^2 L^2}} \sin \left( wt - \tan^{-1} \frac{wL}{R} \right)$   
(c)  $i = \frac{E_0 L}{\sqrt{R^2 + w^2 L^2}} \sin \left( wt + \tan^{-1} \frac{R}{wL} \right)$   
(d)  $i = \frac{E_0 L}{\sqrt{R^2 + w^2 L^2}} \sin \left( wt - \tan^{-1} \frac{R}{wL} \right)$

Ans. (b)

13. A particle executes S.H.M. Such that in two of its positions, the velocities are  $u$ ,  $v$  and the corresponding accelerations  $\alpha$ ,  $\beta$ . The distance between the position is

(a)  $\frac{\alpha + \beta}{u^2 - v^2}$       (b)  $\frac{\alpha^2 + \beta^2}{u^2 - v^2}$

(c)  $\frac{u^2 - v^2}{\alpha + \beta}$       (d)  $\frac{u^2 - v^2}{\alpha - \beta}$

Ans. (c)

# Chapter

## Partial Differential Equations

---

### Introduction

Partial differential equations arise, in cases when a dependent variable is a function of two or more independent variables. An ordinary differential equation can be formed by eliminating arbitrary constants from a relation between two variables such as  $f(x, y) = 0$ , and in general the order of the differential equation is equal to the number of arbitrary constants eliminated. A partial differential equation, on the other hand, can be formed by eliminating not-arbitrary constants, but arbitrary functions, from a relation involving three or more variables, provided such an elimination is possible. In many problems of science and engineering a dependent variable is connected implicitly or explicitly with two or more independent variables. If  $z = z(x, y)$  is a dependent variable where  $x$  and  $y$  are the independent variables, then the first order partial derivative of  $z$  with respect to  $x$  and  $y$  are denoted by  $p = \frac{\partial z}{\partial x}$  and  $q = \frac{\partial z}{\partial y}$ . The second order

partial derivatives of  $z$  are given by

$$r = \frac{\partial^2 z}{\partial x^2}, s = \frac{\partial^2 z}{\partial x \partial y}, t = \frac{\partial^2 z}{\partial y^2}$$

**Definition:** An equation involving one or more partial derivatives of an unknown function of two or more independent variables is called a partial differential equation.

**Example 1.**  $\frac{\partial^2 u}{\partial x^2} = a^2 \frac{\partial^2 u}{\partial t^2}$  is a partial differential equation.

**Example 2.**  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$  is a partial differential equation.

**Definition 2** The order of the highest derivative occurring in a partial differential equation is called the order of the equation.

**Example 1.**  $x^2 \frac{\partial z}{\partial x} + y^2 \frac{\partial z}{\partial y} = z$  is a first order partial differential equation.

**Example 2.** The equation

$\left(\frac{\partial^2 u}{\partial x^2}\right)^2 + u^2 \left(\frac{\partial u}{\partial x}\right) = f(x, y)$  is a second order partial differential equation.

**Definition 3.** The degree of a partial differential equation is the degree of the highest order partial derivative occurring in the equation.

**Example** The degree of the equation

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \text{ is one}$$

**Notation**

If  $z$  is a function of two independent variables say  $x$  and  $y$  then, we shall use the following notation for the partial derivatives of  $z$ .

$$\frac{\partial z}{\partial x} = p, \quad \frac{\partial z}{\partial y} = q, \quad \frac{\partial^2 z}{\partial x^2} = r, \quad \frac{\partial^2 z}{\partial x \partial y} = s, \quad \frac{\partial^2 z}{\partial y^2} = t$$

i.e.  $z_x = p, z_y = q, z_{xx} = r, z_{xy} = s, z_{yy} = t$

**Formation of Partial Differential Equations**

Partial differential equation can be formed in two ways- (1) eliminating arbitrary constants and (2) eliminating arbitrary functions

**1. By Elimination of Arbitrary constants:**

We can form partial differential equation by eliminating arbitrary constants from the given equations.

If the number of arbitrary constants is equal to the number of variables, in the given equation of a curve, we get a first order partial differential equation

Consider the equation

$$f(x, y, z, a, b) = 0 \tag{1}$$

Where  $a$  and  $b$  are arbitrary constants

Differentiating (1) partially with respect to  $x$  and  $y$ , we get

$$\frac{\partial f}{\partial x} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial x} = 0 \text{ or } \frac{\partial f}{\partial x} + \frac{\partial f}{\partial z} p = 0 \tag{2}$$

and  $\frac{\partial f}{\partial y} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial y} = 0 \text{ or } \frac{\partial f}{\partial y} + \frac{\partial f}{\partial z} q = 0 \tag{3}$

Eliminating  $a$  and  $b$  from the equations (1), (2) and (3), we get an equation of the form

$$\phi(x, y, z, p, q) = 0$$

Which is the required partial differential equation of (1)

**Example 1** Eliminate the constants  $a$  and  $b$  from the following equations

(A)  $z = (x + a)(y + b)$

Partial Differential Equations

(B)  $z = (x - a)^2 + (y - b)^2$

(C)  $ax^2 + by^2 + z^2 = 1$

**Solution (A)** We have  $z = (x+a)(y+b)$  (1)

Differentiating equation (1) partially with respect to x and y, we get

$$\frac{\partial z}{\partial x} = p = (y + b) \text{ \& } \frac{\partial z}{\partial y} = q = (x + a)$$

Substituting in (1) we have  $z = pq$  which is the required differential equation

(B) The given equation is

$$z = (x - a)^2 + (y - b)^2 \quad (1)$$

Differentiating equation (1) partially with respect to x and y, we get

$$\frac{\partial z}{\partial x} = 2(x - a) \text{ \& } \frac{\partial z}{\partial y} = 2(y - b)$$

On squaring and adding these equations, we get

$$\begin{aligned} \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 &= 4[(x - a)^2 + (y - b)^2] \\ &= 4z \text{ using (1)} \\ \Rightarrow p^2 + q^2 &= 4z \end{aligned}$$

Which is the required differential equation

(C) The given equation is

$$ax^2 + by^2 + z^2 = 1 \quad (1)$$

Differentiating equation (1) partially with respect to x and y, we get

$$2ax + 2z \frac{\partial z}{\partial x} = 0 \Rightarrow 2ax + 2zp = 0$$

$$\Rightarrow ax = -zp$$

$$\Rightarrow a = -zp/x$$

$$\text{and } 2by + 2z \frac{\partial z}{\partial y} = 0 \Rightarrow 2by + 2zq = 0$$

$$\Rightarrow by = -zq$$

$$\Rightarrow b = -zq/y$$

putting the values of a and b in equation (1) we get

$$-\frac{z}{x} p x^2 + \left(-\frac{z}{y} q y^2\right) + z^2 = 1$$

$$\text{or } z(px + qy) = z^2 - 1$$

Which is the required differential equation.

**Example 2.** Find the partial differential equation of all planes cutting of equal intercepts with x and y axes

**Solution** Let  $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$  (1)

be a equation of a plane making equal intercepts on x and y axis, so in this case a = b.

Differentiating (1) partially with respect to x and y, we get

$$\frac{1}{a} + \frac{1}{c} \frac{\partial z}{\partial x} = 0 \text{ i.e. } \frac{-c}{a} = p \quad (2)$$

$$\text{and } \frac{1}{b} + \frac{1}{c} \frac{\partial z}{\partial y} = 0 \text{ i.e. } \frac{-c}{a} = q \because a = b \quad (3)$$

From (2) and (3), we have

$$p = q \\ \text{i.e. } p - q = 0$$

Which is the required differential equation

**Example 3** Find the partial differential equation of all spheres whose centres lie on the Z-axis (U.P.T.U. 2009)

**Solution** Let the equation of the sphere having its centre on z-axis be

$$x^2 + y^2 + (z-c)^2 = r^2 \quad (1)$$

Differentiating (1) partially with respect to x and y, we get

$$2x + 2(z-c) \frac{\partial z}{\partial x} = 0 \\ \text{or } x + (z-c)p = 0 \\ \text{or } z-c = -x/p \quad (2)$$

$$\text{and } 2y + 2(z-c) \frac{\partial z}{\partial y} = 0 \\ \text{or } y + (z-c)q = 0 \\ \text{or } z-c = -y/q \quad (3)$$

From (2) and (3), we have

$$-\frac{x}{p} = -\frac{y}{q}$$

$$\text{i.e. } xq - yp = 0$$

which is the required differential equation



## 2. Formation of Partial Differential Equations by The Elimination of Arbitrary Function of Specific Function

When one Arbitrary Function is involved

In this case the resulting partial differential equation is a first order partial differential equation

Let the arbitrary function be of the form

$$z = f(u) \tag{1}$$

where  $u$  is function of  $x, y, z$

Differentiating (1) partially with respect to  $x$  and  $y$ , we get

$$\frac{\partial z}{\partial x} = \frac{\partial f}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial f}{\partial u} \frac{\partial u}{\partial z} \frac{\partial z}{\partial x} \tag{2}$$

$$\text{and } \frac{\partial z}{\partial y} = \frac{\partial f}{\partial u} \frac{\partial u}{\partial y} + \frac{\partial f}{\partial u} \frac{\partial u}{\partial z} \frac{\partial z}{\partial y} \tag{3}$$

By eliminating the arbitrary function  $f$ , from (1), (2) and (3) we get a first order partial differential equation

**Example 4** Form the partial differential equation by eliminating the arbitrary function  $f$  from the relation

$$z = y^2 + 2f\left(\frac{1}{x} + \log y\right) \tag{I.A.S. 2007, Bihar, P.C.S. 1995}$$

**Solution** We have

$$z = y^2 + 2f\left(\frac{1}{x} + \log y\right) \tag{1}$$

Differentiating (1) partially with respect to  $x$  and  $y$ , We get

$$\frac{\partial z}{\partial x} = p = 2f'\left(\frac{1}{x} + \log y\right) \left(-\frac{1}{x^2}\right)$$

$$\text{or } -px^2 = 2f'\left(\frac{1}{x} + \log y\right) \tag{2}$$

$$\text{and } \frac{\partial z}{\partial y} = q = 2y + 2f'\left(\frac{1}{x} + \log y\right) \left(\frac{1}{y}\right)$$

$$\text{or } qy - 2y^2 = 2f'\left(\frac{1}{x} + \log y\right) \tag{3}$$

From (2) and (3) we have

$$-px^2 = qy - 2y^2$$

$$\text{i.e. } x^2p + qy = 2y^2$$

Which is the required partial differential equation.

**Example 5.** Eliminate the arbitrary function  $f$  from the equation

$$z = f\left(\frac{xy}{z}\right)$$

Solution. We have  $z = f\left(\frac{xy}{z}\right)$  (1)

Differentiating (1) partially with respect to  $x$  and  $y$ , we get

$$\frac{\partial z}{\partial x} = p = f'\left(\frac{xy}{z}\right) y \left(\frac{z - xp}{z^2}\right)$$
 (2)

$$\frac{\partial z}{\partial y} = q = f'\left(\frac{xy}{z}\right) x \left(\frac{z - yq}{z^2}\right)$$
 (3)

Dividing (2) by (3) we have

$$\frac{p}{q} = \frac{y(z - xp)}{x(z - yq)}$$

$$\text{or } \frac{p}{q} = \frac{yz - xyp}{xz - xyq}$$

$$\text{or } px - qy = 0$$

Which is the required partial differential equation.

### When Two Arbitrary Functions are Involved

When two arbitrary functions are to be eliminated from the given relation to form a partial differential equation, we differentiate twice or more number of times and eliminate the arbitrary functions from the relations obtained.

**Example 6.** Form a partial differential equation by eliminating the function  $f$  and  $F$  from

$$z = f(x + iy) + F(x - iy)$$

**Solution.** The given equation is

$$z = f(x + iy) + F(x - iy)$$
 (1)

Differentiating (1) partially with respect to  $x$  and  $y$ , we get

$$\frac{\partial z}{\partial x} = f'(x + iy) + F'(x - iy)$$
 (2)

$$\text{and } \frac{\partial z}{\partial y} = i f'(x + iy) - i F'(x - iy)$$
 (3)

Differentiating equations (2) and (3) partially once again w.r.t.  $x$  and  $y$ , we get

Partial Differential Equations

$$\frac{\partial^2 z}{\partial x^2} = f''(x + iy) + F''(x - iy) \quad (4)$$

and  $\frac{\partial^2 z}{\partial y^2} = -f''(x + iy) - F''(x - iy) \quad (5)$

Adding (4) and (5), we get

$$\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2} = 0$$

Which is required partial differential equation of second order.

**EXERCISE**

1. Form Partial differential equations by eliminating the arbitrary constants from the following:

- (i)  $z = (x+a)(y+b)$                       Ans.  $pq = z$
- (ii)  $2z = (ax + y)^2 + b$                       Ans.  $q^2 = px + qy$
- (iii)  $z = ax + (1-a)y + b$                       Ans.  $p+q = 1$
- (iv)  $az + b = a^2x + y$                       Ans.  $pq = 1$
- (v)  $z = a \log \left( \frac{b(y-1)}{1-x} \right)$                       Ans.  $px + qy = p + q$

2. Form the partial differential equation by eliminating the arbitrary functions from

- (i)  $z = xy + f(x^2 + y^2)$                       Ans.  $py - qx = y^2 - x^2$
- (ii)  $z = x + y + f(xy)$                       Ans.  $px - qy = x - y$
- (iii)  $z = f(xy)$                       Ans.  $p+q = 0$
- (iv)  $z = f(x^2 + y^2)$                       Ans.  $xq - yp = 0$
- (v)  $z = f(x^2 - y^2)$                       Ans.  $yp + xq = 0$
- (vi)  $f(x^2 + y^2 + z^2, z^2 - 2xy) = 0$                       Ans.  $(p-q)z = y - x$

Hint. The given equation is  $f(x^2 + y^2 + z^2, z^2 - 2xy) = 0 \quad (1)$

Let  $u = x^2 + y^2 + z^2, v = z^2 - 2xy \quad (2)$

so equation (1) becomes  $f(u, v) = 0 \quad (3)$

Differentiating (3) partially w.r.t  $x$ , we get

$$\frac{\partial f}{\partial u} \left( \frac{\partial u}{\partial x} + p \frac{\partial u}{\partial z} \right) + \frac{\partial f}{\partial v} \left( \frac{\partial v}{\partial x} + p \frac{\partial v}{\partial z} \right) = 0 \quad (4)$$

From (2) we have

$$\left. \begin{aligned} \frac{\partial u}{\partial x} = 2x, \quad \frac{\partial u}{\partial z} = 2z, \quad \frac{\partial v}{\partial y} = -2x \\ \frac{\partial v}{\partial z} = 2z, \quad \frac{\partial u}{\partial y} = 2y, \quad \frac{\partial v}{\partial x} = -2y \end{aligned} \right\} \quad (5)$$

From (4) and (5) we have

$$\frac{\partial f / \partial u}{\partial f / \partial v} = \frac{-\left(\frac{\partial v}{\partial x} + p \frac{\partial v}{\partial z}\right)}{\left(\frac{\partial u}{\partial x} + p \frac{\partial u}{\partial z}\right)} \quad (6)$$

Similarly differentiating w.r.t. y, we get

$$\frac{\partial f / \partial u}{\partial f / \partial v} = -\frac{\left(\frac{\partial v}{\partial y} + q \frac{\partial v}{\partial z}\right)}{\left(\frac{\partial u}{\partial y} + q \frac{\partial u}{\partial z}\right)} \quad (7)$$

putting the values from (5) in equation (6) and (7), we get

$$\frac{\partial f / \partial u}{\partial f / \partial v} = -\frac{(-2y + 2pz)}{(2x + 2z)}$$

$$\& \frac{\partial f / \partial u}{\partial f / \partial v} = -\frac{(-2x + 2qz)}{(2y + 2z)}$$

$$\therefore \frac{pz - y}{pz + x} = \frac{qz - x}{qz + y}$$

$$\Rightarrow pz(x + y) - qz(x + y) = y^2 - x^2$$

$$\text{or } (p - q)z = y - x$$

3. Form a partial differential equation by eliminating the arbitrary function  $\phi$  from  $\phi(x + y + z, x^2 + y^2 - z^2) = 0$ . What is the order of this partial differential equation? (U.P.P.C.S. 1993, Bihar P.C.S. 2007)

Hint. Given

$$\phi(x + y + z, x^2 + y^2 - z^2) = 0 \quad (1)$$

$$\text{Let } u = x + y + z \text{ \& } v = x^2 + y^2 - z^2 \quad (2)$$

$$\text{Then (1) becomes } f(u, v) = 0 \quad (3)$$

Differentiating (3) partially w.r.t. x, we get

$$\frac{\partial \phi}{\partial u} \left(\frac{\partial u}{\partial x} + p \frac{\partial u}{\partial z}\right) + \frac{\partial \phi}{\partial v} \left(\frac{\partial v}{\partial x} + p \frac{\partial v}{\partial z}\right) = 0 \quad (4)$$

Partial Differential Equations

$$\text{From (2) } \left. \begin{aligned} \frac{\partial u}{\partial x} = 1, \frac{\partial u}{\partial z} = 1, \frac{\partial v}{\partial x} = 2x \\ \frac{\partial v}{\partial z} = -2z, \frac{\partial u}{\partial y} = 1, \frac{\partial v}{\partial y} = 2y \end{aligned} \right\} \quad (5)$$

$$\text{From (4) and (5) } \frac{\partial \phi}{\partial u} (1+p) + 2 \frac{\partial \phi}{\partial v} (x - pz) = 0$$

$$\text{or } \frac{\partial \phi}{\partial u} = -2(x - pz) / (1+p) \quad (6)$$

Again differentiating (3) partially w.r.t y, we get

$$\frac{\partial \phi}{\partial u} \left( \frac{\partial u}{\partial y} + q \frac{\partial u}{\partial z} \right) + \frac{\partial \phi}{\partial v} \left( \frac{\partial v}{\partial y} + q \frac{\partial v}{\partial z} \right) = 0$$

$$\text{or } \frac{\partial \phi}{\partial u} (1 + q) + 2 \frac{\partial \phi}{\partial v} (y - qz) = 0 \text{ using (5)}$$

$$\text{or } \frac{\partial \phi}{\partial u} = -2(y - qz) / (1 + q) \quad (7)$$

From (6) and (7) by eliminating  $\phi$ , we get

$$\frac{(x-pz)}{1+p} = \frac{y-qz}{1+q}$$

$$\text{or } (1 + q)(x - pz) = (1 + p)(y - qz)$$

$$\text{or } (y + z)p - (x + z)q = x - y$$

which is the desired partial differential equation of first order.

**Solution of Partial Differential Equation by Direct integration:**

**Example 1.** Solve  $\frac{\partial^2 z}{\partial x^2} = xy$

**Solution.** Given  $\frac{\partial^2 z}{\partial x^2} = xy \quad (1)$

treating y as constant and integrating (1) with respect to x, we get

$$\frac{\partial z}{\partial x} = y \frac{x^2}{2} + f(y) \text{ (say)} \quad (2)$$

Integrating (2) with respect to x keeping y as constant, we get

$$z = y \frac{x^3}{6} + x f(y) + g(y)$$

Hence the required solution is  $z = \frac{x^3 y}{6} + x f(y) + g(y)$

**Example 2** Solve  $\frac{\partial^2 z}{\partial x \partial y} = \frac{1}{xy}$

**Solution.** Given  $\frac{\partial^2 z}{\partial x \partial y} = \frac{1}{xy}$  (1)

treating  $y$  as constant and integrating (1) w.r.t  $x$ , we get

$$\frac{\partial z}{\partial y} = \frac{1}{y} \log x + \phi(y)$$
 (2)

Now keeping  $x$  as constant and integrating (2) w.r.t  $y$ , we get

$$z = \log x \log y + \int \phi(y) dy + g(x)$$

or  $z = \log x \log y + f(y) + g(x)$

which is the required solution

**Example 3.** Solve  $\frac{\partial^3 z}{\partial x^2 \partial y} = \cos(2x + 3y)$

**Solution.** we have  $\frac{\partial^3 z}{\partial x^2 \partial y} = \cos(2x + 3y)$  (1)

treating  $y$  as constant and integrating (1) w.r.t.  $x$ , we get

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{1}{2} \sin(2x + 3y) + f(y)$$

Integrating w.r.t.  $x$ , we get

$$\begin{aligned} \frac{\partial z}{\partial y} &= -\frac{1}{4} \cos(2x + 3y) + x \int f(y) dx + g(y) \\ &= -\frac{1}{4} \cos(2x + 3y) + x \phi(y) + g(y) \end{aligned}$$

Integrating w.r.t.  $y$  we get

$$z = -\frac{1}{12} \sin(2x + 3y) + x \int f(y) dy + \int g(y) dy$$

$$\Rightarrow z = -\frac{1}{12} \sin(2x + 3y) + x \phi_1(y) + \phi_2(y)$$

**Lagrange's Linear Equation**

The partial differential equation of the form

$$Pp + Qq = R \tag{1}$$

where P, Q and R are functions of x, y, z is called Lagrange's linear partial differential equation. Lagrange's linear equation is a first order partial differential equation.

**Method of solving Lagrange's equation**

Equation (1) i.e. Lagrange's equation is obtain by eliminating arbitrary function from  $\phi(u, v) = 0$  where u and v are functions of x, y and z

Differentiating

$$\phi(u, v) = 0 \tag{2}$$

partially with respect to x and y, we get

$$\frac{\partial \phi}{\partial u} \left( \frac{\partial u}{\partial x} + \frac{\partial u}{\partial z} p \right) + \frac{\partial \phi}{\partial v} \left( \frac{\partial v}{\partial x} + \frac{\partial v}{\partial z} p \right) = 0 \tag{3}$$

$$\text{and } \frac{\partial \phi}{\partial u} \left( \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} q \right) + \frac{\partial \phi}{\partial v} \left( \frac{\partial v}{\partial y} + \frac{\partial v}{\partial z} q \right) = 0 \tag{4}$$

Eliminating  $\frac{\partial \phi}{\partial u}$  and  $\frac{\partial \phi}{\partial v}$  from equation (3) and (4) we get

$$\begin{vmatrix} \frac{\partial u}{\partial x} + \frac{\partial u}{\partial z} p & \frac{\partial v}{\partial x} + \frac{\partial v}{\partial z} p \\ \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} q & \frac{\partial v}{\partial y} + \frac{\partial v}{\partial z} q \end{vmatrix} = 0$$

$$\text{i.e. } \left( \frac{\partial u}{\partial x} + \frac{\partial u}{\partial z} p \right) \left( \frac{\partial v}{\partial y} + \frac{\partial v}{\partial z} q \right) - \left( \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} q \right) \left( \frac{\partial v}{\partial x} + \frac{\partial v}{\partial z} p \right) = 0$$

$$\text{or } \left( \frac{\partial u}{\partial y} \frac{\partial v}{\partial z} - \frac{\partial u}{\partial z} \frac{\partial v}{\partial y} \right) p + \left( \frac{\partial u}{\partial z} \frac{\partial v}{\partial x} - \frac{\partial u}{\partial x} \frac{\partial v}{\partial z} \right) q = \frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \frac{\partial v}{\partial x} \tag{5}$$

Comparing (1) and (5), we get

$$p = \frac{\partial u}{\partial x} \frac{\partial v}{\partial z} - \frac{\partial u}{\partial z} \frac{\partial v}{\partial y}$$

$$Q = \frac{\partial u}{\partial z} \frac{\partial v}{\partial x} - \frac{\partial u}{\partial x} \frac{\partial v}{\partial z}$$

$$R = \frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \frac{\partial v}{\partial x}$$

Now let us suppose  $u = C_1$ ,  $v = C_2$  are two solutions of the Lagrange's equation  $Pp + Qq = R$

Differentiating  $u = C_1$  and  $v = C_2$  partial with respect to  $x$  and  $y$ , we have

$$\frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy + \frac{\partial u}{\partial z} dz = du = 0 \quad (6)$$

$$\text{and } \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy + \frac{\partial v}{\partial z} dz = dv = 0 \quad (7)$$

From (6) and (7) by cross-multiplication we have

$$\frac{dx}{\frac{\partial u}{\partial y} \frac{\partial v}{\partial z} - \frac{\partial u}{\partial z} \frac{\partial v}{\partial y}} = \frac{dy}{\frac{\partial u}{\partial z} \frac{\partial v}{\partial x} - \frac{\partial u}{\partial x} \frac{\partial v}{\partial z}} = \frac{dz}{\frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \frac{\partial v}{\partial x}}$$

$$\text{or } \frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R}$$

The solutions of these equations are

$$u = C_1 \text{ and } v = C_2$$

Hence  $\phi(u, v) = 0$  is a solution of equation (1)

**Working rule:**

**Step 1:** Form auxiliary equations  $\frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R}$

**Step 2:** Solve the above auxiliary equations Let the two solutions obtained be denoted by  $u = C_1$  and  $v = C_2$

**Step 3:** The required solution of the equation

$$Pp + Qq = R \text{ is}$$

$$\phi(u, v) = 0$$

**Example 1.** Solve  $yzp + zxq = xy$

**Solution.** Given  $yzp + zxq = xy$  (1)

Lagrange's auxiliary equations for (1) are

$$\frac{dx}{yz} = \frac{dy}{zx} = \frac{dz}{xy} \quad (2)$$

Taking first two members, we have



Partial Differential Equations

$$x dx - y dy = 0$$

Integrating, we have

$$x^2 - y^2 = C_1$$

Similarly taking the first and the last members

we get

$$x^2 - z^2 = C_2$$

Therefore, the required solutions is

$$f(x^2 - y^2, x^2 - z^2) = 0$$

**Example 2.** Solve  $p \tan x + q \tan y = \tan z$  (U.P.P.C.S. 1990)

**Solution.** Here Lagrange's auxiliary equations are

$$\frac{dx}{\tan x} = \frac{dy}{\tan y} = \frac{dz}{\tan z} \quad (1)$$

From first two fraction of (1) we get

$$\cot x dx = \cot y dy$$

Integrating, we get

$$\log \sin x = \log \sin y + \log C_1$$

$$\text{or } \log \left( \frac{\sin x}{\sin y} \right) = \log C_1$$

$$\text{or } \sin x / \sin y = C_1 \quad (2)$$

Similarly, from the last two fractions,

we have

$$\sin y / \sin z = C_2 \quad (3)$$

From (2) and (3) required general solution of the given equation is

$$\phi \left( \frac{\sin x}{\sin y}, \frac{\sin y}{\sin z} \right) = 0$$

**Example 3.** Find the general integral of

$$(mz - ny) p + (nx - lz) q = ly - mx \quad (\text{I.A.S. 1977})$$

**Solution.** The auxiliary equations are

$$\frac{dx}{mz - ny} = \frac{dy}{nx - lz} = \frac{dz}{ly - mx} \quad (1)$$

Choosing  $x, y, z$  as multipliers, each fraction of (1)

$$= \frac{xdx + y dy + z dz}{x(mz - ny) + y(nx - lz) + z(ly - mx)}$$

$$= \frac{x \, dx + y \, dy + z \, dz}{0}$$

$$\therefore x \, dx + y \, dy + z \, dz = 0$$

Which on integration gives

$$x^2 + y^2 + z^2 = C_1 \tag{2}$$

choosing  $l, m, n$  as multipliers, each fraction of (1)

$$= \frac{l \, dx + m \, dy + n \, dz}{l(mz - ny) + m(nx - lz) + n(ly - mx)}$$

$$= \frac{l \, dx + m \, dy + n \, dz}{0}$$

$$\therefore l \, dx + m \, dy + n \, dz = 0$$

which on integration, we have

$$lx + my + nz = C_2 \tag{3}$$

$\therefore$  Required general solution is

$$\phi(x^2 + y^2 + z^2, lx + my + nz) = 0$$

where  $\phi$  is an arbitrary function

**Example 4.** Find the general integral of  $(y + zx) p - (x + yz) q = x^2 - y^2$

(U.P.P.C.S. 2002, Bihar P.C.S. 2002)

**Solution.** The Lagrange's auxiliary equations are

$$\frac{dx}{y+zx} = \frac{dy}{-(x+yz)} = \frac{dz}{x^2-y^2} \tag{1}$$

$$\text{i.e. } \frac{y \, dx + x \, dy}{y^2 - x^2} = \frac{dz}{x^2 - y^2}$$

$$\text{or } d(xy) + dz = 0$$

on integration, we get

$$xy + z = C_1 \tag{2}$$

$$\text{Again } \frac{dx + dy}{(y-x) - z(y-x)} = \frac{dz}{x^2 - y^2}$$

$$\text{or } \frac{dx + dy}{-(1-z)} = \frac{dz}{x+y}$$

$$\text{or } (x+y)(dx + dy) + (1-z) dz = 0$$

$$\text{or } d\left\{\frac{1}{2}(x+y)^2\right\} + (1-z) dz = 0$$

Integrating above, we get

$$\frac{1}{2} (x+y)^2 + \left( z - \frac{z^2}{2} \right) = C_2$$

or  $x^2 + y^2 + 2xy + 2z - z^2 = C_3$  where  $2C_2 = C_3$

or  $x^2 + y^2 - z^2 = C_4$  where  $C_3 - 2C_1 = C_4$

Therefore, the general solution is

$$f(x^2 + y^2 - z^2, xy + z) = 0$$

where  $f$  is an arbitrary function

**Example 5.** Solve

$$(2x^2 + y^2 + z^2 - 2yz - zx - xy) p + (x^2 + 2y^2 + z^2 - yz - 2zx - xy) q = x^2 + y^2 + 2z^2 - yz - zx - 2xy \quad (\text{I.A.S. 1992})$$

**Solution.** The Lagrange's auxiliary equations are

$$\frac{dx}{2x^2 + y^2 + z^2 - 2yz - zx - xy} = \frac{dy}{x^2 + 2y^2 + z^2 - yz - 2zx - xy} = \frac{dz}{x^2 + y^2 + 2z^2 - yz - zx - 2xy}$$

$$\therefore \frac{dx - dy}{(x-y)(x+y+z)} = \frac{dy - dz}{(y-z)(x+y+z)} = \frac{dz - dx}{(z-x)(x+y+z)}$$

Taking first two fractions we have

$$\frac{dx - dy}{x - y} = \frac{dy - dz}{y - z}$$

Integrating, we get

$$\log(x-y) = \log(y-z) + \log C_1$$

$$\text{or } \frac{x-y}{y-z} = C_1$$

Similarly, we have

$$\frac{z-x}{y-z} = C_2$$

$$\therefore \text{The required solution is } \phi \left( \frac{x-y}{y-z}, \frac{z-x}{y-z} \right) = 0$$

**Example 6.** Solve

$$(y+z+t) \frac{\partial t}{\partial x} + (z+x+t) \frac{\partial t}{\partial y} + (x+y+t) \frac{\partial t}{\partial z} = x+y+z \quad (\text{I.A.S. 1995})$$

**Solution.** The Lagrange's auxiliary equations are

$$\frac{dx}{y+z+t} = \frac{dy}{z+x+t} = \frac{dz}{x+y+t} = \frac{dt}{x+y+z} \quad (1)$$

$$\therefore \frac{dx-dy}{-(x-y)} = \frac{dy-dz}{-(y-z)} = \frac{dz-dt}{-(z-t)} = \frac{dx+dy+dz+dt}{3(x+y+z+t)}$$

Taking first and Second fraction, we have

$$\frac{dx - dy}{x-y} = \frac{dy-dz}{y-z}$$

Integrating above we have

$$\log (x-y) = \log (y-z) + \log C_1$$

$$\frac{x-y}{y-z} = C_1 \tag{2}$$

Similarly taking second and third fraction, we have

$$\frac{z-t}{y-z} = C_2 \tag{3}$$

Again taking third and fourth fraction, we have

$$\frac{dz - dt}{z-t} + \frac{dx+dy+dz+dt}{3(x+y+z+t)} = 0$$

Integrating, above we have

$$\log (z-t) + \frac{1}{3} \log (x+y+z+t) = \log C_3$$

$$\text{or } (z-t) (x+y+z+t)^{1/3} = C_3$$

$\therefore$  The general integral is

$$f \left[ \frac{x-y}{y-z}, \frac{z-t}{y-z}, (z-t) (x+y+z+t)^{1/3} \right] = 0$$

**Example 7.** Find the surface whose tangent planes cut off an intercept of constant length  $k$  from the axis of  $z$ . (IAS 1993)

**Solution.** Equation of the tangent plane at  $(x, y, z)$  is

$$Z-z = p (X-x) + q (Y-y)$$

Since  $k$  is the intercept on the axis of  $z$

$$\therefore \text{when } X = 0 = Y, Z = k$$

$$\therefore k-z = p (-x) + q (-y)$$

$$\text{or } xp + yq = z-k$$

The subsidiary equations are

$$\frac{dx}{x} = \frac{dy}{y} = \frac{dz}{z-k}$$

Taking the first two members, we have

Partial Differential Equations

$$\frac{dx}{x} = \frac{dy}{y}$$

$$\therefore \log x = \log y - \log C_1$$

$$\text{or } \frac{x}{y} = C_1$$

Again taking the first and last members.

we have

$$\frac{dx}{x} = \frac{dz}{z-k}$$

$$\therefore \log x = \log (z-k) - \log C_2$$

$$\text{or } \frac{z-k}{x} = C_2$$

Therefore, the general solution is  $\phi \left( \frac{y}{x}, \frac{z-k}{x} \right) = 0$  which represents the required surface.

**Example 8.** Solve

$$(z^2 - 2yz - y^2)p + (xy + xz)q = xy - xz$$

If the solution of the above equation represents a sphere. What will be the coordinate of its centre. (Bihar P.C.S. 1999, Roorkee 1975)

**Solution.** Lagrange's auxiliary equations are

$$\frac{dx}{z^2 - 2yz - y^2} = \frac{dy}{x(y+z)} = \frac{dz}{x(y-z)} \quad (1)$$

From the last two fractions of (1), we get

$$\frac{dy}{y+z} = \frac{dy}{y-z}$$

$$\text{or } (y-z) dy = (y+z) dz$$

$$\text{or } ydy - (zdy + ydz) - zdz = 0$$

$$\text{or } ydy - d(yz) - zdz = 0$$

Integrating, we get

$$y^2 - 2yz - z^2 = C_1 \quad (2)$$

Again choosing  $x, y, z$  as multipliers, each fractions of (1)

$$= \frac{xdx + ydy + zdz}{x(z^2 - 2yz - y^2) + y(xy + xz) + z(xy - xz)}$$

$$= \frac{xdx + ydy + zdz}{0}$$

$$xdx + ydy + zdz = 0$$

Integrating we get

$$x^2 + y^2 + z^2 = C_2 \quad (3)$$

From (2) and (3), the required general integral is

$$\phi(y^2 - 2yz - z^2, x^2 + y^2 + z^2) = 0 \quad (4)$$

Where  $\phi$  is an arbitrary function

From (4) we observe that if the solution represent a sphere, then co-ordinates of its centre must be (0, 0, 0) i.e. origin

**NON-LINEAR PARTIAL DIFFERENTIAL EQUATIONS** Those equations in which p and q occur other than in the first degree are called non-linear partial differential equations of the first order. In other words partial differential equations which contains p and q with powers higher than unity and the product of p and q are called non-linear partial differential equations.

### Special Types of Equations

**Standard I.** Equations involving only p and q and no x, y, z That is, equation of the form  $f(p, q) = 0$  (1)

i.e. equation which are independent of x, y, z.

Let the required solution be

$$z = ax + by + c \quad (2)$$

where a and b are connected by

$$f(a, b) = 0 \quad (3)$$

Where a, b, c are constants,

Differentiating (1) partially with respect to x and y, we get

$$\frac{\partial z}{\partial x} = a \text{ and } \frac{\partial z}{\partial y} = b$$

Which when substituted in (3), gives (1)

From (3) we may find b in terms of a

i.e.  $b = \phi(a)$  say

The required solution of (1) is

$$z = ax + \phi(a)y + C$$

**Example 1** Solve  $p^2 + q^2 = 1$

**Solution.** The equation is of the form  $f(p, q) = 0$

The solution is given by

$$z = ax + by + C$$

where  $a^2 + b^2 = 1$

$$\text{or } b = \sqrt{1-a^2}$$

Hence, the required solution is

$$z = ax + \sqrt{1-a^2} y + C$$

**Equation Reducible to  $f(p, q) = 0$**

In some cases, we transform the equations into  $f(p, q) = 0$  by making suitable substitutions

**Example 2.** Solve  $(x + y)(p + q)^2 + (x - y)(p - q)^2 = 1$  (I.A.S. 1991)

**Solution.** putting  $x + y = X^2, x - y = Y^2$

so that

$$p = \frac{\partial z}{\partial x} = \frac{\partial z}{\partial X} \frac{\partial X}{\partial x} + \frac{\partial z}{\partial Y} \frac{\partial Y}{\partial x} = \frac{1}{2X} \frac{\partial z}{\partial X} + \frac{1}{2Y} \frac{\partial z}{\partial Y}$$

$$\text{and } q = \frac{\partial z}{\partial y} = \frac{\partial z}{\partial X} \frac{\partial X}{\partial y} + \frac{\partial z}{\partial Y} \frac{\partial Y}{\partial y} = \frac{1}{2X} \frac{\partial z}{\partial X} - \frac{1}{2Y} \frac{\partial z}{\partial Y}$$

$$\therefore p + q = \frac{1}{X} \frac{\partial z}{\partial X} \text{ and } p - q = \frac{1}{Y} \frac{\partial z}{\partial Y}$$

putting in the given equation, we have

$$\left(\frac{\partial z}{\partial X}\right)^2 + \left(\frac{\partial z}{\partial Y}\right)^2 = 1$$

Which is of the form of standard I

$\therefore$  The complete integral is given by

$$z = aX + bY + c$$

$$\text{Where } a^2 + b^2 = 1 \text{ or } b^2 = \sqrt{1-a^2}$$

$\therefore$  The required complete integral is

$$z = a\sqrt{x+y} + \sqrt{1-a^2} \sqrt{x-y} + c$$

**Example 3.** Find the complete integral of  $(y - x)(qy - px) = (p - q)^2$  (I.A.S. 1992)

**Solution.** Let us put  $X = x + y$  and  $Y = xy$

$$\text{so that } p = \frac{\partial z}{\partial x} = \frac{\partial z}{\partial X} \frac{\partial X}{\partial x} + \frac{\partial z}{\partial Y} \frac{\partial Y}{\partial x} = \frac{\partial z}{\partial X} + y \frac{\partial z}{\partial Y}$$

$$\text{and } q = \frac{\partial z}{\partial y} = \frac{\partial z}{\partial X} \frac{\partial X}{\partial y} + \frac{\partial z}{\partial Y} \frac{\partial Y}{\partial y} = \frac{\partial z}{\partial X} + x \frac{\partial z}{\partial Y}$$

Substituting in the given equation we have

$$(y-x)(y-x) \frac{\partial z}{\partial X} = (y-x)^2 \left( \frac{\partial z}{\partial Y} \right)^2$$

$$\text{or } \frac{\partial z}{\partial X} = \left( \frac{\partial z}{\partial Y} \right)^2$$

Which is of the form of standard I

∴ The complete integral is given by

$$z = aX + bY + c$$

where  $a = b^2$

∴ The complete integral is

$$z = b^2(x+y) + bxy + c$$

**Standard II Equations involving only p, q and z i.e. equations of the form  $f(z, p, q) = 0$**

Equations of the form  $f(z, p, q) = 0$  (1)

Let us assume  $z = f(x+ay)$  as a trial solution of given equation (1), where a is an arbitrary constant

∴  $z = f(X)$  where  $X = x + ay$

$$\therefore p = \frac{\partial z}{\partial x} = \frac{\partial z}{\partial X} \frac{\partial X}{\partial x} = \frac{\partial z}{\partial X} \cdot 1 = \frac{\partial z}{\partial X}$$

$$\text{and } q = \frac{\partial z}{\partial y} = \frac{\partial z}{\partial X} \frac{\partial X}{\partial y} = a \frac{\partial z}{\partial X} = a \frac{dz}{dX}$$

∴ Equation (1) reduces to the form

$$f\left(z, \frac{dz}{dX}, a \frac{dz}{dX}\right) = 0$$

Which is an ordinary differential equation of order one. Integrating it we may get the complete integral

**Example 4** Find the complete integral of

$$z^2 (p^2 z^2 + q^2) = 1$$

(I.A.S, 1997)

**Solution** Putting  $z = f(x + ay) = f(X)$

where  $X = x + ay$

$$\text{so that } p = \frac{\partial z}{\partial x} = \frac{dz}{dX}$$

$$\text{and } q = \frac{\partial z}{\partial y} = a \frac{dz}{dX}$$

The equation becomes



$$z^2 \left[ \left( \frac{dz}{dX} \right)^2 z^2 + a^2 \left( \frac{dz}{dX} \right)^2 \right] = 1$$

$$\text{or } z^2 (z^2 + a^2) \left( \frac{dz}{dX} \right)^2 = 1$$

$$\text{or } z\sqrt{(z^2 + a^2)} dz = dX$$

Integrating, we have

$$\frac{1}{3} (z^2 + a^2)^{3/2} = X + b$$

$$\text{or } 9(x + ay + b)^2 = (z^2 + a^2)^3$$

which is the required complete integral

**Example 5.** Solve  $pq = x^m y^n z^l$

(I.A.S. 1989, 1994)

**Solution.** Putting  $\frac{x^{m+1}}{m+1} = X, \frac{y^{n+1}}{n+1} = Y$

$$\text{so that } p = \frac{\partial z}{\partial x} = \frac{\partial z}{\partial X} \frac{dX}{dx} = x^m \frac{\partial z}{\partial X}$$

$$\text{and } q = \frac{\partial z}{\partial y} = \frac{\partial z}{\partial Y} \frac{dY}{dy} = y^n \frac{\partial z}{\partial Y}$$

Then the given equation reduce to

$$\frac{\partial z}{\partial X} \frac{\partial z}{\partial Y} = z^l \tag{1}$$

Which is the form of standard II

$\therefore$  putting  $z = f(X + aY) = f(u)$

where  $u = X + aY$

$$\frac{\partial z}{\partial X} = \frac{dz}{du} \frac{\partial u}{\partial X} = \frac{dz}{du}$$

$$\text{and } \frac{\partial z}{\partial Y} = \frac{dz}{du} \frac{\partial u}{\partial Y} = a \frac{dz}{du}$$

Equation (1) becomes

$$a \left( \frac{dz}{du} \right)^2 = z^l$$

$$\text{or } z^{-l/2} dz = \frac{du}{\sqrt{a}}$$

Integrating, we have

$$\frac{z^{-\frac{l}{2}+1}}{-\frac{l}{2}+1} = \frac{u}{\sqrt{a}} + b$$

$$\text{or } \frac{z^{-\frac{l}{2}+1}}{-\frac{l}{2}+1} = \frac{1}{\sqrt{a}} \left( \frac{x^{m+1}}{m+1} + a \frac{y^{n+1}}{n+1} \right) + b$$

**Standard III i.e. Equation of the form  $f(x, p) = F(y, q)$**

As a trial solution, let us put each side equal to a arbitrary constant

i.e.  $f(x, p) = F(y, q) = a$

from which we obtain

$$p = f_1(x, a) \text{ and } q = f_2(y, a)$$

Now from  $dz = p dx + q dy$

we have  $dz = f_1(x, a) dx + f_2(y, a) dy$

$$z = \int f_1(x, a) dx + \int f_2(y, a) dy + b$$

Which is the complete integral

**Example 6.** Solve  $p^2 + q^2 = x + y$

**Solution.** Let  $p^2 - x = y - q^2 = a$

$$\therefore p = \sqrt{(x+a)} \text{ and } q = \sqrt{(y-a)}$$

putting in  $dz = p dx + q dy$ , we have

$$dz = \sqrt{(x+a)} dx + \sqrt{(y-a)} dy$$

Integrating, we get

$$z = \frac{2}{3} (x+a)^{3/2} + \frac{2}{3} (y-a)^{3/2} + b$$

**Example 7.** Solve  $z(p^2 - q^2) = x - y$

(I.A.S. 1989, U.P.P.C.S. 1992)

**Solution** The given equation can be written as

$$\left( \sqrt{z} \frac{\partial z}{\partial x} \right)^2 - \left( \sqrt{z} \frac{\partial z}{\partial y} \right)^2 = x - y$$

putting  $\sqrt{z} dz = dZ$ , so that  $Z = \frac{2}{3} z^{3/2}$

The equation becomes  $\left( \frac{\partial Z}{\partial x} \right)^2 - \left( \frac{\partial Z}{\partial y} \right)^2 = x - y$

or  $P^2 - Q^2 = x - y$

Partial Differential Equations

where  $P = \frac{\partial Z}{\partial x}$ ,  $Q = \frac{\partial Z}{\partial y}$

or  $P^2 - x = Q^2 - y = a$

Which is of the form of standard III

$\therefore P^2 - x^2 = Q^2 - y = a$

$\therefore P = \sqrt{(x+a)}$  and  $Q = \sqrt{(y+a)}$

Putting in  $dZ = Pdx + Qdy$ , we have

$dZ = \sqrt{(x+a)} dx + \sqrt{(y+a)} dy$

Integrating  $Z = \frac{2}{3} (x+a)^{3/2} + \frac{2}{3} (y+a)^{3/2} + b$

or  $z^{3/2} = (x+a)^{3/2} + (y+a)^{3/2} + c$

**Standard IV Equation of the form  $Z = px + qy + f(p, q)$   
(Clairaut's form)**

An equation of the form  $z = px + qy + f(p, q)$ , Which is linear in  $x$  and  $y$  is called Clairaut's equation

The complete solution of the Clairaut's equation is

$z = ax + by + f(a, b)$

i.e. the solution of Clairaut's equation is obtained putting  $p = a$  and  $q = b$

**Example 8** Solve  $z = px + qy + c\sqrt{1+p^2+q^2}$

(I.A.S. 1989, Bihar P.C.S. 2007; U.P.P.C.S. 2005)

**Solution.** This is of the form of Standard IV

$\therefore$  The complete integral is

$z = ax + by + c\sqrt{(1+a^2+b^2)}$

**EXERCISE**

1. Solve the following partial differential equations

(a)  $y^2p - xyq = x(z-2y)$

Ans.  $\phi(x^2-y^2, zy-y^2) = 0$

(b)  $xzp + yzq = xy$

Ans.  $\phi(x/y, xy-z^2) = 0$

(c)  $\left(\frac{y^2z}{x}\right)p + xzq = y^2$

Ans.  $\phi(x^3 - y^3, x^2 - y^2) = 0$

(d)  $z(xp - yq) = y^2 - x^2$

Ans.  $f(xy, x^2 + y^2 + z^2) = 0$

(e)  $p + 3q = 5z + \tan(y - 3x)$

Ans.  $\phi[y-3x, e^{-5x} \{5z + \tan(y-3x)\}] = 0$

(f)  $x(y+z)p - y(x^2+z)q = z(x^2-y^2)$

Ans.  $f(x^2 + y^2 - 2z, xyz) = 0$

(g)  $x^2(y-z)p + (z-x)y^2q = z^2(x-y)$

Ans.  $f\left(xyz, \frac{1}{x} + \frac{1}{y} + \frac{1}{z}\right) = 0$

(h)  $(x+2z)p + (4zx-y)q = 2x^2 + y$

(Roorkee 1976)

Ans.  $\phi(xy - z^2, x^2 - y - z) = 0$

(i)  $px(z-2y^2) = (z-xy)(z-y^2-2x^3)$

Ans.  $\phi\left(\frac{z}{x} + x^2 - \frac{y^2}{x}, \frac{y}{z}\right) = 0$

(j)  $(x^2 - yz)p + (y^2 - xz)q = z^2 - xy$

Ans.  $\phi\left(\frac{x-y}{y-z}, \frac{y-z}{z-x}\right) = 0$

2. Solve the following partial differential equations.

(a)  $p = 2q^2 + 1$

Ans.  $z = ax + \sqrt{(2b^2 + 1)}y + c$

(b)  $x^2p^2 + y^2q^2 = z^2$

(Raj SLET 1997)

Ans.  $\log z = a \log x + \sqrt{(1-a^2)} \log y + c$

(c)  $(x^2 + y^2)(p^2 + q^2) = 1$

Ans.  $z = \frac{a}{2} \log(x^2 + y^2) + \sqrt{(1-a^2)} \tan^{-1} \frac{y}{x} + C$

(d)  $9(p^2z + q^2) = 4$

Ans.  $(z+a)^3 = (x+ay+b)^2$

(e)  $z^2(p^2z^2 + q^2) = 1$

Ans.  $9(x+ay+b)^2 = (z^2+a^2)^3$

(f)  $p(1+q^2) = q(z-a)$

Ans.  $4(bz - ab - 1) = (x + by + c)^2$

(g)  $p^2 = z^2(1 - pq)$

Ans.  $\frac{1}{\sqrt{a}} \log \left[ z\sqrt{a} + \sqrt{a(1+az^2)} + \sqrt{(1+az^2)} \right] = x + c$

Partial Differential Equations

(h)  $p(1 + q^2) = q(z-a)$

Ans.  $4a(z-a) = 4 + (x+ay+b)^2$

(i)  $q^2y^2 = z(z - px)$

Ans.  $z^{2a^2/\{-1 \pm \sqrt{1+4a^2}\}} = bxy^a$

(j)  $p(1 + q^2) = q(z - a)$

Ans.  $4a(z - a) = 4 + (x + ay + b)^2$

(k)  $z^2(p^2 + q^2 + 1) = c^2$

Ans.  $(1+a^2)(c-z^2) = (x + ay + b)^2$

3. Solve the following partial differential equations.

(a)  $\sqrt{p} + \sqrt{q} = 2x$

Ans.  $z = \frac{1}{6}(a + 2x)^3 + a^2y + b$

(b)  $yp = 2yx + \log q$

Ans.  $az = ax^2 + a^2x + e^{ay} + ab$

(c)  $z^2(p^2 + q^2) = x^2 + y^2$

Ans.  $z^2 = x\sqrt{(a+x^2)} + a \log \left\{ x + \sqrt{(a+x^2)} \right\} + y\sqrt{(y^2 - a)} - a \log \left\{ y + \sqrt{(y^2 - a)} \right\} + C$

(d)  $p^2 - 2x^2 = q^2 - y$

Ans.  $z = \frac{2}{3}x^3 + ax \pm \frac{2}{3}(y + a)^{3/2} + b$

(e)  $p^2 + q^2 = z^2(x+y)$

Ans.  $\log z = \frac{2}{3}(a+x)^{3/2} + \frac{2}{3}(y-a)^{3/2} + C$

(f)  $z = px + qy + 2pq$

Ans.  $z = ax + by + 2ab$

(g)  $z = px + qy + p^2 + q^2$

Ans.  $z = ax + by + a^2 + b^2$

(h)  $z = px + qy + \log pq$

Ans.  $z = ax + by + \log ab$

---

**UNIT - V**  
**APPLICATIONS OF PARTIAL**  
**DIFFERENTIAL EQUATIONS**

---

# Chapter 9

## Applications of Partial Differential Equations

---

### INTRODUCTION

The problems related to fluid mechanics, solid state physics, heat transfer, electromagnetic theory, Wave equation and other areas of physics and engineering are governed by partial differential equations subject to certain given conditions, called boundary conditions. The process to find all solutions of a partial differential equation under given conditions is known as a boundary value problem. The method of solution of such equations differ from that used in the case of ordinary differential equations. Method of separation of variables is a powerful tool to solve such boundary value problem when partial differential equation is linear with homogenous boundary conditions. Most of the problems involving linear partial differential equations Can be solved by the method of separation of variables discussed below.

### METHOD OF SEPARATION OF VARIABLES

It involves a solution which breaks up into a product of functions each of which contains only one of the variables. The following example explain this method.

**Example 1** Apply the method of separation of variables to solve

$$\frac{\partial^2 z}{\partial x^2} - 2 \frac{\partial^2 z}{\partial x} + \frac{\partial z}{\partial x} = 0 \quad (\text{U.P.T.U. 2005, 09})$$

**Solution** Assume the trial solution  $z = X(x) Y(y)$  (i)

where  $X$  is a function of  $x$  alone and  $Y$  that of  $y$  alone, substituting this value of  $z$  in the given equation we have

$$X'' \cdot Y - 2X' \cdot Y + XY' = 0 \quad \text{where } X' = \frac{dX}{dx}, \quad Y' = \frac{dY}{dy} \text{ . etc}$$

separating the variables, we get

$$\frac{X'' - 2X'}{X} = -\frac{Y'}{Y} \quad (\text{ii})$$

since  $x$  and  $y$  are independent variables, therefore, (ii) can only be true if each side is equal to the same constant,  $K$ (say), so we have

$$\frac{X'' - 2X'}{X} = -\frac{Y'}{Y} = K$$

Therefore,  $\frac{X'' - 2X'}{X} = K$  i.e.  $X'' - 2X' - kX = 0$  (iii)

and  $-\frac{Y'}{Y} = k$  i.e.  $Y' + kY = 0$  (iv)

To solve the ordinary linear equation (iii) the auxiliary equation is  $m^2 - 2m - k = 0$

$$\Rightarrow m = 1 \pm \sqrt{(1+k)}$$

$\therefore$  The solution of (iii) is  $X = C_1 e^{\{1+\sqrt{(1+k)}\}x} + C_2 e^{\{1-\sqrt{(1+k)}\}x}$  and the solution of (iv) is  $Y = C_3 e^{-ky}$

Substituting these values of X and Y in (i), we get

$$z = \left\{ C_1 e^{\{1+\sqrt{(1+k)}\}x} + C_2 e^{\{1-\sqrt{(1+k)}\}x} \right\} \cdot C_3 e^{-ky}$$

$$\text{i.e. } z = \left\{ a e^{\{1+\sqrt{(1+k)}\}x} + b e^{\{1-\sqrt{(1+k)}\}x} \right\} e^{-ky}$$

where  $a = C_1 C_3$  and  $b = C_2 C_3$

which is the required complete solution.

**Example 2** Using the method of separation of variables, solve  $\frac{\partial u}{\partial x} = 2 \frac{\partial u}{\partial t} + u$ ,  
where  $u(x, 0) = 6 e^{-3x}$

(U.P.T.U. 2006)

**Solution** Assume the solution  $u(x, t) = X(x) T(t)$  (i)

Substituting in the given equation, we have

$$X'T = 2XT' + XT$$

$$\text{or } (X' - X) T = 2 XT'$$

$$\text{or } \frac{X' - X}{2X} = \frac{T'}{T} = k(\text{say})$$

$$\therefore X' - X - 2kX = 0$$

$$\text{or } \frac{X'}{X} = 1 + 2k \tag{ii}$$

$$\text{and } \frac{T'}{T} = k \tag{iii}$$

solving (ii),  $\log X = (1+2k)x + \log c$

$$\text{or } X = ce^{(1+2k)x}$$

From (iii),  $\log T = kt + \log c'$

$$\text{or } T = c'e^{kt}$$



Applications of Partial Differential Equations

Thus  $u(x,t) = XT$

$$= cc'e^{(1+2k)x} e^{kt}$$

$$\text{Now } 6e^{-3x} = u(x, 0) = cc'e^{(1+2k)x} \tag{iv}$$

$$\therefore cc' = 6 \text{ and } 1+2k = -3 \text{ or } k = -2$$

Substituting these values in (iv) we get

$$u = 6e^{-3x} e^{-2t} \text{ i.e. } u = 6e^{-(3x+2t)} \text{ which is the required solution}$$

**Example 3** Solve by the method of separation of variables,  $\frac{\partial u}{\partial x} = 2\frac{\partial u}{\partial y} + u$ , where

$$u(x, 0) = 3e^{-5x} - 2e^{-3x}$$

**Solution** Assume the trial solution  $u = X(x) Y(y)$  (i)

where  $X$  is the function of  $x$  alone and  $Y$  that of  $y$  alone.

substituting this value of  $u$  in the given equation we have

$$X' Y = 2XY' + XY$$

$$\frac{X'}{X} = \frac{2Y'}{Y} + 1 = k(\text{say}) \tag{ii}$$

$$\text{Now } \frac{X'}{X} = k \Rightarrow \frac{1}{X} \frac{dX}{dx} = k$$

$$\Rightarrow \frac{dX}{X} = k dx$$

on integrating, we get

$$\log_e X = kx + \log_e C_1$$

$$\Rightarrow X = c_1 e^{kx}$$

And taking last two terms of equation (ii), we have

$$\frac{2}{Y} \frac{dY}{dy} + 1 = k$$

$$\Rightarrow \frac{2}{Y} \frac{dY}{dy} = k - 1$$

$$\Rightarrow \frac{dY}{Y} = \frac{(k-1)}{2} dy$$

$$\text{on integrating, } \log_e Y = \frac{(k-1)}{2} y + \log c_2$$

$$\Rightarrow Y = c_2 e^{\frac{(k-1)y}{2}}$$

From (i), we get

$$u = c_1 c_2 e^{kx} e^{(k-1)y/2}$$

From (i), we get

$$u = c_1 c_2 e^{kx} e^{(k-1)y/2}$$

$$\Rightarrow u = \sum_{n=1}^{\infty} b_n e^{k_n x} e^{(k_n - 1)y/2} \quad \text{(iii)}$$

$$b_n = c_1 c_2 \text{ and } k = K_n$$

which is the most general solution of given equation, putting  $y=0$  and  $u = 3e^{-5x} - 2e^{-3x}$  in equation (iii) we get

$$3e^{-5x} - 2e^{-3x} = \sum_{n=1}^{\infty} b_n e^{k_n x} = b_1 e^{k_1 x} + b_2 e^{k_2 x}$$

comparing the terms on both sides, we get

$$b_1 = 3, k_1 = -5, b_2 = -2, k_2 = -3$$

Hence the required solution of given equation is from (iii), we have

$$u = 3e^{-5x} - e^{-3y} + (-2)e^{-3x} e^{-2y}$$

$$\Rightarrow u = 3e^{-(5x+3y)} - 2e^{-(3x+2y)}$$

**Example 4.**

Use the method of separation of variables to solve the equation.

$$\frac{\partial^2 v}{\partial x^2} = \frac{\partial v}{\partial t} \text{ given that } v=0 \text{ when } t \rightarrow \infty, \text{ as well as } v=0 \text{ at } x=0 \text{ and } x=l.$$

**Solution.** Assume the trial solution  $v = XT$  (i)

where  $X$  is a function of  $x$  alone and  $Y$  that of  $y$  given

$$\Rightarrow \frac{\partial v}{\partial t} = X \frac{dT}{dt} \quad \text{and} \quad \frac{\partial^2 v}{\partial x^2} = T \frac{d^2 X}{dx^2}$$

Substituting these values in the given differential equation, we get

$$\begin{aligned} T \frac{d^2 X}{dx^2} &= X \frac{dT}{dt} \text{ or } \frac{1}{T} \frac{dT}{dt} = \frac{1}{X} \frac{d^2 X}{dx^2} = -p^2 \text{ (say)} \\ \Rightarrow \frac{1}{T} \frac{dT}{dt} &= -p^2 \text{ or } \frac{dT}{dt} + p^2 T = 0 \end{aligned} \quad \text{(ii)}$$

$$\text{and } \frac{1}{X} \frac{d^2 X}{dx^2} = -p^2 \text{ or } \frac{d^2 X}{dx^2} + p^2 X = 0 \quad \text{(iii)}$$

Solving (ii) and (iii) we get

$$T = c_1 e^{-p^2 t} \text{ and } X = c_2 \cos px + c_3 \sin px$$

Substituting these values of  $X$  and  $T$  in (i) we get

$$v = c_1 e^{-p^2 t} (c_2 \cos pt + c_3 \sin pt) \quad \text{(iv)}$$

### Applications of Partial Differential Equations

putting  $x=0, v=0$  in (ii), we get

$$0 = c_1 e^{-p^2 t} c_2 \Rightarrow c_2 = 0 \text{ since } c_1 \neq 0$$

$$\Rightarrow v = c_1 e^{-p^2 t} c_3 \sin px \quad (v)$$

putting  $x=l, v=0$  in (v), we get

$$c_1 c_3 e^{-p^2 t} \sin pl = 0$$

$$\Rightarrow \sin pl = 0 = \sin n\pi$$

$$\Rightarrow p = n\pi/l, n \text{ is any integer}$$

$$\therefore v = c_1 c_3 e^{-(n^2 \pi^2 t)/l^2} \sin\left(\frac{n\pi x}{l}\right)$$

$$= b_n e^{-n^2 \pi^2 t/l^2} \sin\left(\frac{n\pi x}{l}\right)$$

where  $b_n = c_1 c_3$

The most general solution is  $v = \sum_{n=1}^{\infty} b_n e^{-n^2 \pi^2 t/l^2} \sin\left(\frac{n\pi x}{l}\right)$

### **PARTIAL DIFFERENTIAL EQUATIONS OF ENGINEERING**

A number of problems in engineering give rise to the following well known partial differential equations.

(i) Wave equation  $\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$

(ii) One dimensional heat flow equation  $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$

(iii) Two dimensional heat flow equation which in steady state becomes the two dimensional Laplace's equation  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$

(iv) Transmission line equations

(v) Vibrating membrane. Two dimensional wave equation.

(vi) Laplace's equation in three dimensions.


Besides these, the partial differential equations frequently occur in the theory of Elasticity and Hydraulics.

Starting with the method of separation of variables, we find their solutions subject to specific boundary conditions and the combination of such solution gives the desired solution. Quite often a certain condition is not applicable. In such cases, the most general solution is written as the sum of the particular solutions already found and the constants are determined using Fourier series so as to satisfy the remaining conditions.

**Vibrations of a stretched string (one dimensional wave Equation)**

Consider a tightly stretched elastic string of length  $l$  and fixed ends A and B and subjected to constant tension  $T$  as shown in figure. The tension  $T$  will be considered to be large as compared to the weight of the string so that the effects of gravity are negligible.

Let the string be released from rest and allowed to vibrate. We shall study the subsequent motion of the string, with no external forces acting on it, assuming that each point of the string makes small vibrations at right angles to the equilibrium position AB, of the string entirely in one plane.



Taking the end A as the origin, AB as the x-axis and AY perpendicular to it as the y-axis, so that the motion takes place entirely in the xy -plane. Above figure shown the string in the position APB at times  $t$ . Consider the motion of the element PQ of the string between its points  $P(x, y)$  and  $Q(x + \delta x, y + \delta y)$ , where the tangents make angles  $\psi$  and  $\psi + \delta\psi$  with the x axis. Clearly the element is moving upwards with the acceleration  $\frac{\partial^2 y}{\partial t^2}$ . Also the vertical component of the

force acting on this element

$$= T \sin (\psi + \delta\psi) - T \sin \psi$$

$$\approx T (\psi + \delta\psi - \psi) \because \sin \psi = \psi, \text{ as } \psi \text{ is very small}$$

$$= T \delta\psi \text{ (approximately)}$$

The acceleration of the elements in the QY direction is  $\frac{\partial^2 y}{\partial t^2}$ . If the length of PQ is

$\delta s$ , then the mass of PQ is  $m \cdot \delta s$ .

Hence, by Newton's second law, the equation of motion becomes

$$m \delta s \frac{\partial^2 y}{\partial t^2} = T \delta\psi$$

$$\text{or } \frac{\partial^2 y}{\partial t^2} = \frac{T}{m} \frac{\delta \psi}{\delta s}$$

As  $Q \rightarrow P$ ,  $\delta s \rightarrow 0$ . Therefore, taking limit as  $\delta s \rightarrow 0$ , the above equation becomes

$$\frac{\partial^2 y}{\partial t^2} = \frac{T}{m} \cdot \frac{\delta \psi}{\delta s}$$

Where  $\frac{\delta \psi}{\delta s}$  = Curvature at P of the deflection curve

$$\frac{\delta \psi}{\delta s} = \frac{\frac{\partial^2 y}{\partial x^2}}{\left[1 + \left(\frac{\partial y}{\partial x}\right)^2\right]^{3/2}} \text{ using formula for the radius of curvature}$$

$$= \frac{\partial^2 y}{\partial x^2}, \text{ approximately, since } \left(\frac{\partial y}{\partial x}\right)^2 \text{ is negligible because } \frac{\partial y}{\partial x} \text{ is small}$$

$$\therefore \frac{\partial^2 y}{\partial t^2} = \frac{T}{m} \frac{\partial^2 y}{\partial x^2}$$

Putting  $\frac{T}{m} = C^2$  (positive), the displacement  $y(x, t)$  is given by the equation

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2}$$

This partial differential equation is known as one dimensional wave equation.

### **Solution of the one dimensional wave Equation**

The one dimensional wave equation is

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2} \tag{i}$$

Assume that a solution of (i) is of the form  $y = X(x) T(t)$ . where  $X$  is a function of  $x$  alone and  $T$  is a function of  $t$  only.

$$\text{Then } \frac{\partial^2 y}{\partial t^2} = X \frac{d^2 T}{dt^2} \quad \text{and} \quad \frac{\partial^2 y}{\partial x^2} = T \frac{d^2 X}{dx^2}$$

putting these values in (i), we get

$$\frac{1}{X} \frac{d^2 X}{dx^2} = \frac{1}{C^2 T} \frac{d^2 T}{dt^2} = k(\text{say}) \tag{ii}$$

The (ii) leads to the ordinary differential equations.

$$\frac{d^2 X}{dx^2} - kX = 0 \quad \text{and} \quad \frac{d^2 T}{dt^2} - kC^2 T = 0 \tag{iii}$$

solving (iii) we get

(1) when  $K$  is negative say  $-p^2$ , then

$$X = c_1 \cos px + c_2 \sin px$$

$$T = c_3 \cos cpt + c_4 \sin cpt \tag{iv}$$

(2) when  $k = 0$ , then

$$X = C_5 x + C_6$$

$$T = C_7 t + C_8 \tag{v}$$

(3) when  $k$  is positive say  $p^2$ , then

$$\left. \begin{aligned} X &= C_9 e^{px} + C_{10} e^{-px} \\ T &= C_{11} e^{cpt} + C_{12} e^{-cpt} \end{aligned} \right\}$$

(vi)

of these three solutions we have to choose that solution which is consistent with the physical nature of the problem. As we are dealing with problems on vibration  $y$  must be a periodic function of  $x$  and  $t$ . Hence, the solution must involve trigonometric terms.

Accordingly the solution given by (iv) i.e. of the form

$$y = (C_1 \cos px + C_2 \sin px) (C_3 \cos cpt + C_4 \sin cpt)$$


is the only suitable solution of the wave equation.

**Example 5.** A string of length  $L$  is stretched and fastened to two fixed points. Find the solution of the wave equation  $y_{tt} = a^2 y_{xx}$ , when initial displacement is

$$y(x, 0) = f(x) = b \sin \left( \frac{\pi k}{L} \right) \text{ where symbols have usual meaning.}$$

(U.P.T.U. 2009)

**Solution.** Consider an elastic string tightly stretched between two points  $O$  and  $A$ . Let  $O$  be the origin and  $OA$  as  $x$ -axis on giving a small transverse displacement i.e. the displacement perpendicular to its length.



Applications of Partial Differential Equations

let  $y$  be the displacement at the point P ( $x, y$ ) at any time, the wave equation

$$y_{tt} = a^2 y_{xx}$$

$$\text{or } \frac{\partial^2 y}{\partial t^2} = a^2 \frac{\partial^2 y}{\partial x^2}$$

As the end points of the string are fixed, for all time,

$$y(0, t) = 0 \tag{ii}$$

$$y(L, t) = 0 \tag{iii}$$

Since, the initial transverse velocity of any point of the string is zero, therefore

$$\left( \frac{\partial y}{\partial t} \right)_{t=0} = 0 \tag{iv}$$

$$\text{Also } y(x, 0) = b \sin \frac{\pi x}{L} \tag{v}$$

The general solution of (i) is

$$y = (C_1 \cos px + C_2 \sin px) (C_3 \cos apt + C_4 \sin apt) \tag{vi}$$

Applying the boundary condition

$$y = 0 \text{ at } x = 0$$

$$0 = C_1 (C_3 \cos apt + C_4 \sin apt)$$

$$\therefore C_1 = 0$$

Therefore,

$$y = C_2 \sin px (C_3 \cos apt + C_4 \sin apt) \tag{vii}$$

Again applying  $\frac{\partial y}{\partial t} = 0$ , at  $t = 0$  on (vii)

$$\frac{\partial y}{\partial t} = C_2 \sin px \text{ ap. } (-C_3 \sin apt + C_4 \cos apt)$$

$$0 = C_2 \sin px \text{ ap. } C_4 \Rightarrow C_4 = 0$$

Then (vii) becomes  $y = C_2 C_3 \sin px \cos atp$  (viii)

Applying  $y = 0$  at  $x = L$

$$0 = C_2 C_3 \sin pL \cos atp$$

$$\therefore \sin pL = 0 = \sin n\pi, n = 0, 1, 2, 3, \dots$$

$$\therefore pL = n\pi$$

$$\text{or } p = \frac{n\pi}{L}$$

putting  $p = \frac{n\pi}{L}$  in (viii), we have

$$y = C_2 C_3 \sin \frac{n\pi x}{L} \cos \frac{n\pi at}{L} \quad \text{(ix)}$$

$$\text{At } t = 0, y = b \sin \frac{\pi x}{L}$$

$$b \sin \frac{\pi x}{L} = C_2 C_3 \sin \frac{n\pi x}{L}$$

$$\therefore C_2 C_3 = b, n = 1$$

putting  $C_2 C_3 = b, n = 1$  in (ix) we get

$$y = b \sin \left( \frac{\pi x}{L} \right) \cos \left( \frac{\pi at}{L} \right)$$

which is our required solution.

**Example 6.** A string is stretched and fastened to two points  $l$  apart. Motion is started by displacing the string the form  $y = a \sin \frac{\pi x}{l}$  from which it is released at a time  $t = 0$ . Show that the displacement of any point at a distance  $x$  from one end at time  $t$  is given by

$$y(x, t) = a \sin \left( \frac{\pi x}{l} \right) \cos \left( \frac{\pi ct}{l} \right)$$

(U.P.T.U. 2004, S.V.T.U. 2007)

**Solution :** Solving exactly just like as example 5.

**Example 7 :** A tightly stretched string with fixed end points  $x = 0$  and  $x = l$  is initially in a position given by  $y = y_0 \sin^3 \frac{\pi x}{l}$ .

If it is released from the rest from this position find the displacement  $y(x, t)$ .

**Solution** The equation to the vibrating string be

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2} \quad \text{(i)}$$

Here the initial conditions are

$$y(0, t) = 0, y(l, t) = 0$$

$$\frac{\partial y}{\partial t} = 0 \text{ at } t = 0, y(x, 0) = y_0 \sin^3 \frac{\pi x}{l}$$

The general solution of (i) is of the form

$$y = (C_1 \cos px + C_2 \sin px) (C_3 \cos cpt + C_4 \sin cpt) \quad \text{(ii)}$$

Now  $y = 0$  at  $x = 0$  gives  $C_1 = 0$



Applications of Partial Differential Equations

$$\therefore y = C_2 \sin px (C_3 \cos cpt + C_4 \sin cpt) \quad \text{(iii)}$$

Again  $\frac{\partial y}{\partial t} = 0$  at  $t = 0$  gives  $C_4 = 0$

$$y = c_2 c_3 \sin px \cos cpt \quad \text{(iv)}$$

At  $x = l, y = 0$

$$0 = C_2 C_3 \sin pl \cos cpt$$

$$\Rightarrow \sin pl = 0 = \sin n\pi, n = 0, 1, 2, \dots$$

$$\therefore p = \frac{n\pi}{l}$$

$$\therefore y = C_2 C_3 \sin \frac{n\pi x}{l} \cos \frac{n\pi ct}{l} \quad \text{(v)}$$

Let  $C_2 C_3 = b_n$ , As  $b_n$  is arbitrary constants

Therefore general solution is

$$y(x, t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \cos \frac{n\pi ct}{l} \quad \text{(vi)}$$

At  $t = 0, y = y_0 \sin^3 \frac{\pi x}{l}$ , so from equation (vi) we have

$$y_0 \sin^3 \left( \frac{\pi x}{l} \right) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$$

$$\Rightarrow \frac{y_0}{4} \left( 3 \sin \frac{\pi x}{l} - \sin \frac{3\pi x}{l} \right) = b_1 \sin \frac{\pi x}{l} + b_2 \sin \frac{2\pi x}{l} + b_3 \sin \frac{3\pi x}{l} + \dots$$

$$\therefore b_1 = \frac{3y_0}{4}, b_2 = 0, b_3 = -\frac{y_0}{4}, b_4 = b_5 = b_6 = \dots = 0$$

Hence (vi) becomes

$$y(x, t) = \frac{y_0}{4} \left( 3 \sin \frac{\pi x}{l} \cos \frac{\pi ct}{l} - \sin \frac{3\pi x}{l} \cos \frac{3\pi ct}{l} \right)$$

**Example 8:** A string is stretched and fastened to two points  $l$  apart. Motion is started by displacing the string into the form  $y = k(lx - x^2)$  from which it is released at time  $t = 0$ . Find the displacement of any point on the string at a distance of  $x$  from one end at time  $t$ .

(U.P.T.U. 2002)

**Solution**

The vibration of the string is given by the equation

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2} \quad \text{(i)}$$

As the end points of the string are fixed for all time,

$$y(0, t) = 0 \tag{ii}$$

$$y(l, t) = 0 \tag{iii}$$

Since the initial transverse velocity of any point of the string is zero, therefore

$$\left(\frac{\partial y}{\partial t}\right)_{t=0} = 0 \tag{iv}$$

$$\text{and } y(x, 0) = k(lx - x^2) \tag{v}$$

solution of (i) is

$$y = (c_1 \cos px + c_2 \sin px)(c_3 \cos cpt + c_4 \sin cpt) \tag{vi}$$

$$\text{At } x = 0, y = 0 \text{ gives } c_1 = 0$$

$$y = c_2 \sin px (c_3 \cos cpt + c_4 \sin cpt) \tag{vii}$$

$$\text{At } t = 0, \frac{\partial y}{\partial t} = 0$$

$$0 = c_2 \sin px \cdot cp \cdot c_4$$

$$\therefore c_4 = 0$$

$$y = c_2 c_3 \sin px \cos cpt \tag{viii}$$

At  $x = l, y = 0$


$$0 = C_2 C_3 \sin pl \cos cpt$$

$$\therefore \sin pl = 0 = \sin n\pi; n = 0, 1, 2, 3, \dots$$

$$\therefore p = \frac{n\pi}{l}$$

$$y = C_2 C_3 \sin \frac{n\pi x}{l} \cos \frac{n\pi ct}{l}$$

$$\text{Let } C_2 C_3 = b_n$$



$$y = b_n \sin \frac{n\pi x}{l} \cos \frac{n\pi ct}{l}$$

As  $b_n$  is arbitrary constants and a differential equation satisfy solution for all constants. Then we can write

Applications of Partial Differential Equations

$$y = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi t}{l} \cos \frac{n\pi x}{l} \quad (ix)$$

At  $t = 0, y = k(lx - x^2)$

$$k(lx - x^2) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$$

Applying half range Fourier sine series

$$b_n = \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$$

$$b_n = \frac{2k}{l} \int_0^l (lx - x^2) \sin \frac{n\pi x}{l} dx$$

$$b_n = \frac{2k}{l} \left[ (-1)^{n+1} \frac{2l^3}{n^3 \pi^3} + \frac{2l^3}{n^3 \pi^3} \right]$$

$$b_n = \begin{cases} \frac{8kl^2}{n^3 \pi^3}, & \text{when } n \text{ is odd} \\ 0, & \text{when } n \text{ is even} \end{cases}$$

0, when n is even

$$y = \sum_{n=1}^{\infty} \frac{8kl^2}{n^3 \pi^3} \sin \frac{n\pi x}{l} \cos \frac{n\pi t}{l}, \text{ when } n \text{ is odd}$$

$$\text{or } y = \sum_{n=1}^{\infty} \frac{8kl^2}{(2n-1)^3 \pi^3} \sin \frac{(2n-1)\pi x}{l} \cos \frac{(2n-1)\pi t}{l}$$

Which is required solution.

**Example 9** Solve the boundary value problem

$$\frac{\partial^2 y}{\partial t^2} = 4 \frac{\partial^2 y}{\partial x^2}, \text{ given that } y(0, t) = 0$$

$$y(5, t) = 0, y(x, 0) = 0 \text{ and } \left( \frac{\partial y}{\partial t} \right)_{x=0} = 5 \sin \pi x$$

**Solution:** Applying the method of separation of variables to the wave equation

$$\frac{\partial^2 y}{\partial t^2} = 2^2 \frac{\partial^2 y}{\partial x^2}.$$

The suitable solution is

$$y = (C_1 \cos px + C_2 \sin px) (C_3 \cos 2pt + C_4 \sin 2pt)$$

Applying the initial condition

$$y(x, 0) = 0 \text{ we have}$$

$$0 = C_3 (C_1 \cos px + C_2 \sin px)$$

$$\Rightarrow C_3 = 0$$

$$\therefore y = C_4(C_1 \cos px + C_2 \sin px) \sin 2pt$$

Now using  $y(0, t) = 0$ , we get

$$0 = C_1 C_4 \sin 2pt \Rightarrow C_1 = 0$$

$$\therefore y = C_2 \sin 2pt \sin px$$

Further  $y(5, t) = 0$  we have  $C_2 \sin 2pt \sin 5p = 0$

$$\Rightarrow \sin 5p = 0 = \sin n\pi$$

$$\Rightarrow p = \frac{n\pi}{5}, n = 1, 2, 3, \dots$$

Therefore,

$$y = c_2 \sin\left(\frac{n\pi 2t}{5}\right) \sin\left(\frac{n\pi x}{5}\right)$$

Also the boundary condition  $\left(\frac{\partial y}{\partial t}\right)_{x=0} = 5 \sin \pi x$

$$\therefore c_2 \frac{n\pi^2}{5} \cos\left(\frac{n\pi 2t}{5}\right) \sin\left(\frac{n\pi x}{5}\right) = 5 \sin \pi x$$

$$\Rightarrow n = 5 \text{ and } 2\pi c_2 = 5$$

Therefore, we have

$$y = \frac{5}{2\pi} \sin \pi x \sin 2\pi t$$


**Example 10:** A string of length  $l$  is fastened at both ends A and C. At a distance 'a' from the end A, the string is transversely displaced to a distance 'd' and is released from rest when it is in this position. Find the equation of the subsequent motion.

OR

Find the half period sine series for  $f(x)$  given in the range  $(l, 0)$  by the graph ABC as shown in figure.

(U.P.T.U. 2009)

Applications of Partial Differential Equations



**Solution** let  $y(x, t)$  is the displacement of the string Now, by the one dimensional wave equation we have

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2} \tag{i}$$

The solution of equation (i) is given by

$$y(x, t) = (C_1 \cos px + C_2 \sin px) (C_3 \cos cpt + C_4 \sin cpt) \tag{ii}$$

Now using the boundary conditions as follows

The boundary conditions are

$$\text{At } x=0 \text{ (at A), } y=0 \Rightarrow y(0, t)=0$$

$$\text{and At } x=l \text{ (at C), } y=0 \Rightarrow y(l, t)=0$$

From (ii), we have

$$0 = C_1 (C_3 \cos cpt + C_4 \sin cpt) \Rightarrow C_1 = 0$$

using  $C_1 = 0$  in equation (ii), we get

$$y(x, t) = C_2 \sin px (C_3 \cos cpt + C_4 \sin cpt) \tag{iii}$$

using second boundary condition, from (iii), we have

$$0 = C_2 \sin pl (C_3 \cos cpt + C_4 \sin cpt)$$

$$\Rightarrow \sin pl = 0 \Rightarrow \sin pl = \sin n\pi$$

$$\Rightarrow p = \frac{n\pi}{l}$$

using the value of  $p$  in (iii) we have

$$y(x, t) = C_2 \sin \frac{n\pi x}{l} \left( C_3 \cos \frac{n\pi ct}{l} + C_4 \sin \frac{n\pi ct}{l} \right) \tag{iv}$$

Next, the initial conditions are as follows:

$$\text{velocity } \frac{\partial y}{\partial t} = 0 \text{ at } t=0$$

and displacement at  $t=0$  is

$$y(x,0) = \begin{cases} \frac{d \cdot x}{a}, & 0 \leq x \leq a \\ \frac{d(x-l)}{a-l}, & a \leq x \leq l \end{cases} \quad \begin{array}{l} \therefore \text{Equation of AB is } y = \frac{d \cdot x}{a} \text{ and} \\ \text{equation of BC is } y = \frac{d(x-l)}{a-l} \end{array}$$

From (iv)

$$\frac{\partial y}{\partial t} = c_2 \sin \frac{n\pi x}{l} \left( -\frac{n\pi c c_3}{l} \sin \frac{n\pi ct}{l} + \frac{n\pi c c_4}{l} \cos \frac{n\pi ct}{l} \right)$$

using initial condition we get

$$0 = c_2 c_4 \frac{n\pi c}{l} \cdot \sin \frac{n\pi x}{l} \Rightarrow c_4 = 0$$

using  $c_4 = 0$  in equation (iv), we get

$$y(x,t) = c_2 c_3 \sin \frac{n\pi x}{l} \cdot \cos \frac{n\pi ct}{l}$$

$\therefore$  The general solution of the given problem is

$$y(x,t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \cdot \cos \frac{n\pi ct}{l} \quad (v)$$

Using initial condition in equation (v), we get

$$y(x,0) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$$

which is half range Fourier sine series, so we have

$$\begin{aligned} b_n &= \frac{2}{l} \int_0^l y(x,0) \cdot \sin \frac{n\pi x}{l} dx \\ &= \frac{2}{l} \int_0^a \frac{d}{a} \cdot x \cdot \sin \left( \frac{n\pi x}{l} \right) dx + \frac{2}{l} \frac{d}{(a-l)} \int_a^l (x-l) \sin \frac{n\pi x}{l} dx \\ &\quad + \frac{2d}{l(a-l)} \left[ (x-l) \left( \frac{-l}{n\pi} \right) \cos \frac{n\pi x}{l} - \left( \frac{-l^2}{n^2 \pi^2} \right) \sin \frac{n\pi x}{l} \right]_a^l \\ \Rightarrow b_n &= -\frac{2d}{n\pi} \cos \frac{n\pi a}{l} + \frac{2dl^2}{aln^2 \pi^2} \sin \frac{n\pi a}{l} + \frac{2d}{n\pi} \cos \frac{n\pi a}{l} - \frac{2dl^2}{l(a-l)n^2 \pi^2} \sin \frac{n\pi a}{l} \\ \Rightarrow b_n &= \frac{2dl^2}{a(l-a)n^2 \pi^2} \sin \frac{n\pi a}{l} \end{aligned}$$

$\therefore$  From (v), we get

$$y(x,t) = \frac{2dl^2}{a(l-a)\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} \sin \frac{n\pi a}{l} \cdot \sin \frac{n\pi x}{l} \cdot \cos \frac{n\pi ct}{l}$$

Applications of Partial Differential Equations

**Example 11:** A lightly stretched string of length  $l$  with fixed ends is initially in equilibrium position. it is set vibrating by giving each point a velocity  $V_0 \sin^3 \frac{\pi x}{l}$ .

Find the displacement  $y(x, t)$ .

(I.A.S. 2004, U.P.T.U. 2003)

**Solution** The equation of the vibrating string is  $\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2}$  (i)

The boundary condition are  $y(0, t) = 0, y(l, t) = 0$  (ii)

Also the initial conditions are  $y(x, 0) = 0$  (iii)

and  $\left(\frac{\partial y}{\partial t}\right)_{t=0} = V_0 \sin^3 \frac{\pi x}{l}$  (iv)

Since the vibration of the string is periodic, therefore, the solution of (i) is of the form

$$y(x, t) = (C_1 \cos px + C_2 \sin px) (C_3 \cos cpt + C_4 \sin cpt)$$

by (ii)  $y(0, t) = C_1(C_3 \cos cpt + C_4 \sin cpt) = 0$

For this to be true for all time  $C_1 = 0$

$\therefore y(x, t) = C_2 \sin px (C_3 \cos cpt + C_4 \sin cpt)$

Also  $y(l, t) = C_2 \sin pl (C_3 \cos cpt + C_4 \sin cpt) = 0$  for all  $t$ .

This gives  $pl = n\pi$  or  $p = \frac{n\pi}{l}$ ,  $n$  being an integer

Thus  $y(x, t) = C_2 \frac{n\pi x}{l} \left( C_3 \cos \frac{cn\pi}{l} t + C_4 \sin \frac{cn\pi}{l} t \right)$

$\therefore y(x, t) = b_n \sin \frac{n\pi x}{l} \sin \frac{cn\pi t}{l}$  where  $b_n = C_2 C_4$

Adding all such solutions the general solution of (i) is

$$y(x, t) = \sum b_n \sin \frac{n\pi x}{l} \sin \frac{cn\pi t}{l} \tag{v}$$

Now  $\frac{\partial y}{\partial t} = \sum b_n \sin \frac{n\pi x}{l} \cdot \frac{cn\pi}{l} \cos \frac{cn\pi t}{l}$

By (iv),  $V_0 \sin^3 \frac{\pi x}{l} = \left(\frac{\partial y}{\partial t}\right)_{t=0} = \sum \frac{cn\pi}{l} b_n \sin \frac{n\pi x}{l}$

or  $\frac{V}{4} \left( 3 \sin \frac{\pi x}{l} - \sin \frac{3\pi x}{l} \right) = \sum \frac{cn\pi}{l} b_n \sin \frac{n\pi x}{l}$

$= \frac{c\pi}{l} b_1 \sin \frac{\pi x}{l} + \frac{2c\pi}{l} b_2 \sin \frac{2\pi x}{l} + \frac{3c\pi}{l} b_3 \sin \frac{3\pi x}{l} + \dots$

Equating Coefficients from both sides, we get

$$\frac{3V_0}{4} = \frac{c\pi}{l} b_1, \quad 0 = \frac{2c\pi}{l} b_2, \quad -\frac{V_0}{4} = \frac{3c\pi}{l} b_3, \dots$$

$$\therefore \frac{3IV_0}{4c\pi}, b_3 = -\frac{IV_0}{12c\pi}, b_2 = b_4 = b_5 = \dots = 0$$

Substituting in (v), the desired solution is

$$y = \frac{IV_0}{12c\pi} \left( 9 \sin \frac{\pi x}{l} \sin \frac{c\pi t}{l} - \sin \frac{3\pi x}{l} \sin \frac{3c\pi t}{l} \right)$$

**Example 12:** A lightly stretched string with fixed end points  $x = 0$  and  $x = l$  is initially at rest in its equilibrium position. If it is vibrating by giving to each of its points a velocity  $\lambda x(l - x)$ . find the displacement of the string at any distance  $x$  from one end at any time  $t$ .

(U.P.T.U. 2002)

**Solution.** The equation of the vibrating string is

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2} \tag{i}$$

The boundary condition are  $y(0, t) = 0, y(l, t) = 0$  (ii)

Also the initial conditions are  $y(x, 0) = 0$  (iii)

and  $\left( \frac{\partial y}{\partial t} \right)_{t=0} = \lambda x(l - x)$  (iv)

As in example 11, the general solution of (i) satisfying the conditions (ii) and (iii) is

$$y(x, t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \cdot \sin \frac{n\pi ct}{l} \tag{v}$$

$$\frac{\partial y}{\partial t} = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \cdot \cos \frac{n\pi ct}{l} \left( \frac{n\pi c}{l} \right)$$

$$\text{By (iv), } \lambda x(l - x) = \left( \frac{\partial y}{\partial t} \right)_{t=0} = \frac{\pi c}{l} \sum_{n=1}^{\infty} n b_n \sin \frac{n\pi x}{l}$$

$$\therefore \frac{\pi c \pi}{l} b_n = \frac{2}{l} \int_0^l \lambda x(l - x) \sin \frac{n\pi x}{l} dx$$

$$= \frac{2\lambda}{l} \left[ (lx - x^2) \left( -\frac{l}{n\pi} \cos \frac{n\pi x}{l} \right) - (l - 2x) \left( -\frac{l^2}{n^2 \pi^2} \sin \frac{n\pi x}{l} \right) + (-2) \left( \frac{l^3}{n^3 \pi^3} \cos \frac{n\pi x}{l} \right) \right]_0^l$$

$$= \frac{4\lambda l^2}{n^3 \pi^3} (1 - \cos n\pi) = \frac{4\lambda l^2}{n^3 \pi^3} [1 - (-1)^n]$$



$$\begin{aligned} \text{or } b_n &= \frac{4\lambda l^2}{c\pi^4 n^4} [1 - (-1)^n] \\ &= \frac{8\lambda l^3}{c\pi^4 (2m-1)^4} \text{ taking } n = 2m-1 \end{aligned}$$

Hence, from (v) the desired solution is

$$y = \frac{8\lambda l^2}{c\pi^4} \sum_{m=1}^{\infty} \frac{1}{(2m-1)^4} \sin \frac{(2m-1)\pi x}{l} \sin \frac{(2m-1)\pi ct}{l}$$

Example 13. Solve completely the equation  $\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2}$ , representing the vibrations of a string of length  $l$ , fixed at both ends, given that  $y(0, t) = 0$ ,  $y(l, t) = 0$   
 $y(x, 0) = f(x)$ , and  $\frac{\partial}{\partial t} y(x, 0) = 0$ ,  $0 < x < l$

(U.P.T.U. 2005)

**Solution** Here the given equation is

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2} \tag{i}$$

The solution of equation (i) is given by

$$y = (C_1 \cos px + C_2 \sin px) (C_3 \cos cpt + C_4 \sin cpt) \tag{ii}$$

Now, applying the boundary conditions  $y = 0$  when  $x = 0$ , we get

$$\begin{aligned} 0 &= C_1(C_3 \cos cpt + C_4 \sin cpt) \\ \Rightarrow C_1 &= 0 \end{aligned}$$

Therefore, equation (ii) becomes

$$y = C_2 \sin px (C_3 \cos cpt + C_4 \sin cpt) \tag{iii}$$

Now putting  $x = l$  and  $y = 0$  in equation (iii), we get

$$\begin{aligned} 0 &= C_2 \sin pl (C_3 \cos cpt + C_4 \sin cpt) \\ \Rightarrow \sin pl &= 0 = \sin n\pi \end{aligned}$$

$$\text{or } pl = n\pi \Rightarrow p = \frac{n\pi}{l}$$

Thus, equation (iii) becomes

$$y = C_2 \sin \frac{n\pi}{l} x \left( C_3 \cos \frac{n\pi ct}{l} + C_4 \sin \frac{n\pi ct}{l} \right) \tag{iv}$$

Differentiating equation (iv) with respect to  $t$ , we get

$$\frac{\partial y}{\partial t} = C_2 \sin \frac{n\pi x}{l} \left( -C_3 \frac{n\pi c}{l} \sin \frac{n\pi ct}{l} + C_4 \frac{n\pi c}{l} \cos \frac{n\pi ct}{l} \right)$$

Using given boundary condition  $\frac{\partial y}{\partial t} = 0$ ,  $t=0$  we get

$$0 = C_2 \sin \frac{n\pi c}{l} \left( 0 + C_4 \frac{n\pi c}{l} \right)$$

$$\Rightarrow C_4 = 0$$

Thus, equation (iv) becomes

$$y = C_2 C_3 \sin \frac{n\pi x}{l} \cos \frac{n\pi ct}{l}$$

Now applying the last boundary condition given, we get

$$f(x) = b_n \sin \frac{n\pi x}{l}, \quad b_n = C_2 C_3$$

Where

$$b_n = \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$$

Thus, the required solution is

$$y = b_n \sin \frac{n\pi x}{l} \cos \frac{n\pi ct}{l}$$

### Solution of wave Equation By D'Almbert's Method

Transform the equation  $\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2}$  to its normal form using the transformation  $u = x + ct$ ,  $v = x - ct$  and hence solve it. Show that the solution may be put in the form  $y = \frac{1}{2} [f(x + ct) + f(x - ct)]$ . Assume initial condition  $y = f(x)$  and  $\frac{\partial y}{\partial t} = 0$  at  $t=0$

(U.P.T.U. 2003)

**Proof.** Consider one dimensional wave equation

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2} \tag{i}$$

Let  $u = x + ct$  and  $v = x - ct$ , be a transformation of  $x$  and  $t$  into  $u$  and  $v$ . then

$$\begin{aligned} \frac{\partial y}{\partial x} &= \frac{\partial y}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial y}{\partial v} \frac{\partial v}{\partial x} = \frac{\partial y}{\partial u} + \frac{\partial y}{\partial v} \quad \therefore \frac{\partial u}{\partial x} = 1 \\ & \qquad \qquad \qquad \frac{\partial v}{\partial x} = 1 \end{aligned}$$

Applications of Partial Differential Equations

or 
$$\frac{\partial}{\partial x} = \frac{\partial}{\partial u} + \frac{\partial}{\partial v}$$

$$\Rightarrow \frac{\partial^2 y}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial y}{\partial x} \right) = \left( \frac{\partial}{\partial u} + \frac{\partial}{\partial v} \right) \left( \frac{\partial y}{\partial u} + \frac{\partial y}{\partial v} \right)$$

$$= \frac{\partial^2 y}{\partial u^2} + \frac{\partial^2 y}{\partial u \partial v} + \frac{\partial^2 y}{\partial v \partial u} + \frac{\partial^2 y}{\partial v^2}$$

$$\Rightarrow \frac{\partial^2 y}{\partial x^2} = \frac{\partial^2 y}{\partial u^2} + 2 \frac{\partial^2 y}{\partial u \partial v} + \frac{\partial^2 y}{\partial v^2} \quad \text{(ii)}$$

and 
$$\frac{\partial y}{\partial t} = \frac{\partial y}{\partial u} \frac{\partial u}{\partial t} + \frac{\partial y}{\partial v} \frac{\partial v}{\partial t} = c \frac{\partial y}{\partial u} - c \frac{\partial y}{\partial v} \quad \therefore \frac{\partial u}{\partial t} = c, \frac{\partial v}{\partial t} = -c$$

or 
$$\frac{\partial}{\partial t} = c \left( \frac{\partial}{\partial u} - \frac{\partial}{\partial v} \right)$$

$$\therefore \frac{\partial^2 y}{\partial t^2} = \frac{\partial}{\partial t} \left( \frac{\partial y}{\partial t} \right) = c \left( \frac{\partial}{\partial u} - \frac{\partial}{\partial v} \right) \left( c \frac{\partial y}{\partial u} - c \frac{\partial y}{\partial v} \right)$$

$$= c^2 \left( \frac{\partial^2 y}{\partial u^2} - 2 \frac{\partial^2 y}{\partial u \partial v} + \frac{\partial^2 y}{\partial v^2} \right)$$

$$\Rightarrow \frac{\partial^2 y}{\partial t^2} = c^2 \left( \frac{\partial^2 y}{\partial u^2} - 2 \frac{\partial^2 y}{\partial u \partial v} + \frac{\partial^2 y}{\partial v^2} \right) \quad \text{(iii)}$$

marking use of equation (ii) and (iii) in equation (i), we get

$$c^2 \left( \frac{\partial^2 y}{\partial u^2} - 2 \frac{\partial^2 y}{\partial u \partial v} + \frac{\partial^2 y}{\partial v^2} \right) = c^2 \left( \frac{\partial^2 y}{\partial u^2} + 2 \frac{\partial^2 y}{\partial u \partial v} + \frac{\partial^2 y}{\partial v^2} \right)$$

$$\Rightarrow 4c^2 \frac{\partial^2 y}{\partial u \partial v} = 0$$

$$\Rightarrow \frac{\partial^2 y}{\partial u \partial v} = 0 \quad \text{(iv)}$$

Integrating equation (iv) w.r.t. 'v' we get

$$\frac{\partial y}{\partial u} = \phi(u) \quad \text{(v)}$$

where  $\phi(u)$  is a constant in respect to v

Again integrate equation (v), we get

$$y = \int \phi(u) du + \phi_2(v)$$

$$\Rightarrow y = \phi_1(u) + \phi_2(v)$$

$$\Rightarrow y(x, t) = \phi_1(x + ct) + \phi_2(x - ct) \quad \text{(vi)}$$

The solution (vi) is D' Alembert's solution of wave equation.

Now, we applying initial conditions  $y = f(t)$  and  $\frac{\partial y}{\partial t} = 0$  at  $t = 0$

From (vi), we get at  $t = 0$

$$f(x) = \phi_1(x) + \phi_2(x) \quad (\text{vii})$$

$$\text{and } \frac{\partial y}{\partial t} = c\phi_1'(x+ct) - c\phi_2'(x-ct)$$

$$\Rightarrow \left(\frac{\partial y}{\partial t}\right)_{t=0} = 0 = c\phi_1'(x+0) - c\phi_2'(x-0)$$

$$\Rightarrow \phi_1'(x) - \phi_2'(x) = 0$$

$$\Rightarrow \phi_1'(x) = \phi_2'(x)$$

on integrating, we get

$$\phi_1(x) = \phi_2(x) + c_1 \quad (\text{viii})$$

using equation (viii) in equation (vii), we get

$$f(x) = \phi_2(x) + c_1 + \phi_2(x) = 2\phi_2(x) + c_1$$

$$\Rightarrow \phi_2(x) = \frac{1}{2}[f(x) - c_1] \Rightarrow \phi_2(x-ct) = \frac{1}{2}[f(x-ct) - c_1]$$

$$\text{and } \phi_1(x) = \frac{1}{2}[f(x) + c_1] \Rightarrow \phi_1(x+ct) = \frac{1}{2}[f(x+ct) + c_1]$$

putting the values of  $\phi_1(x+ct)$  and  $\phi_2(x-ct)$  in eq<sup>n</sup>(vi) we get


$$y(x, t) = \frac{1}{2}[f(x+ct) + f(x-ct)]$$

### **VIBRATING MEMBRANE - TWO DIMENSIONAL WVAE EQUATION**

Consider a lightly stretched uniform membrane (such as the membrane of a drum) with tension  $T$  per unit length is the same in all directions at every point.

Consider the forces on an element  $\delta x \delta y$  of the membranes. Due to its displacement  $u$  perpendicular to the  $xy$  plane, the forces  $T\delta x$  and  $T\delta y$  act on the edges along the tangent to the membrane

Applications of Partial Differential Equations



The forces  $T\delta y$  (tangential to the membrane) on its opposite edges of length  $\delta y$  act at angles  $\alpha$  and  $\beta$  to the horizontal. So, their vertical component

$$= (T\delta y) \sin \beta - (T\delta y) \sin \alpha$$

$$= T\delta y (\tan \beta - \tan \alpha), \quad \because \alpha \text{ and } \beta \text{ are very small i.e. } \sin \alpha \approx \tan \alpha \text{ etc.}$$

$$= T\delta y \left\{ \left( \frac{\partial u}{\partial x} \right)_{x+\delta x} - \left( \frac{\partial u}{\partial x} \right)_x \right\}$$

$$= T\delta y \delta x \frac{\left[ \left( \frac{\partial u}{\partial x} \right)_{x+\delta x} - \left( \frac{\partial u}{\partial x} \right)_x \right]}{\delta x}$$

$$= T\delta y \delta x \frac{\partial^2 u}{\partial x^2}, \text{ upto a first order of approximation}$$

similarly, the forces  $T\delta x$  (the vertical component of the force) acting on the edges of length  $\delta x$  have the vertical component  $= T\delta x \delta y \frac{\partial^2 u}{\partial y^2}$

If  $m$  be the mass per unit area of the membrane, then the equation of motion of the element ABCD is

$$m\delta x \delta y \frac{\partial^2 u}{\partial t^2} = T \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) \delta x \delta y$$

$$\text{or } \frac{\partial^2 u}{\partial t^2} = \frac{T}{m} \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$$

$$\text{or } \frac{\partial^2 u}{\partial t^2} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) \quad \text{where } c^2 = \frac{T}{m}$$

This is the wave equation in two dimensions.

**Solution of the Two-Dimensional wave Equation**

The two dimensional wave equation is given by

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) \tag{i}$$

$$\text{Let } u = XYT \tag{ii}$$

be the solution of (i), where X is a function of x only, Y is a function of y only and T is a function of t only.

$$\Rightarrow \frac{\partial^2 u}{\partial x^2} = X''YT, \quad \frac{\partial^2 u}{\partial y^2} = XY''T \text{ and } \frac{\partial^2 u}{\partial t^2} = XYT''$$

substituting these values in (i), we get

$$\frac{1}{c^2} XYT'' = X''YT + XY''T$$

Dividing by XYT throughout, we get

$$\frac{1}{c^2} \frac{T''}{T} = \frac{X''}{X} + \frac{Y''}{Y} \tag{iii}$$

since each variable is independent, hence this will be true only when each member is a constant. Suitably choosing the constants, we have

$$\frac{d^2 X}{dx^2} + k^2 X = 0, \quad \frac{d^2 Y}{dy^2} + l^2 Y = 0$$

$$\text{and } \frac{d^2 T}{dt^2} + (k^2 + l^2)c^2 T = 0$$

Hence, the solution of these equations are given by

$$X = c_1 \cos kx + c_2 \sin kx$$

$$Y = c_3 \cos ly + c_4 \sin ly$$

$$\text{and } T = c_5 \cos \sqrt{(k^2 + l^2)}ct + c_6 \sin \sqrt{(k^2 + l^2)}ct$$

Hence from (ii), the solution of (i) is given by

$$u(x, y, t) = (c_1 \cos kx + c_2 \sin kx) (c_3 \cos ly + c_4 \sin ly)$$


$$\left[ c_5 \cos \sqrt{(k^2 + l^2)}ct + c_6 \sin \sqrt{(k^2 + l^2)}ct \right] \tag{iv}$$

Let us consider that the membrane is rectangular and stretched between the lines  $x=0, x=a, y=0, y=b$

Applications of Partial Differential Equations

Now the boundary condition are

- (1)  $u = 0$ , when  $x = 0$ , for all  $t$
- (2)  $u = 0$ , when  $x = a$ , for all  $t$
- (3)  $u = 0$ , when  $y = 0$ , for all  $t$
- (4)  $u = 0$ , when  $y = b$ , for all  $t$



Now using condition (1) in (iv), we get

$$0 = c_1 (c_3 \cos ly + c_4 \sin ly) \left[ c_5 \cos \sqrt{(k^2 + l^2)ct} + c_6 \sin \sqrt{(k^2 + l^2)ct} \right]$$

$$\Rightarrow c_1 = 0$$

substituting  $c_1 = 0$  in (iv) and using condition (ii), we get

$$\sin ka = 0 \quad \text{or} \quad k = \frac{m\pi}{a}, \text{ where } m \text{ is an integer}$$

Hence solution of (iii) becomes

$$u(x, y, t) = c_2 c_4 \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} (c_5 \cos pt + c_6 \sin pt) \tag{iv}$$

$$\text{where } p = \pi c \sqrt{\frac{m^2}{a^2} + \frac{n^2}{b^2}}$$

Now replacing the arbitrary constants, we can write the general solution as

$$u(x, y, t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} (A_{mn} \cos pt + B_{mn} \sin pt) \tag{v}$$

Equation (v) is the solution of the wave equation (i) which is zero on the boundary of the rectangular membrane.

Suppose the membrane starts from rest from the initial position.

$$u = f(x, y) \text{ i.e. } u(x, y, 0) = f(x, y)$$

Then using the condition  $\frac{\partial u}{\partial t} = 0$ , when  $t = 0$ , we get  $B_{mn} = 0$

Further using the condition  $u = f(x,y)$  when  $t=0$ , we get

$$f(x,y) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} \quad (vi)$$

This is a double Fourier series. multiplying both sides by  $\sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b}$  and integrating from  $x=0$  to  $x=a$  and  $y=0$  to  $y=b$ , every term on the right except one become zero. Thus, we have

$$\int_0^a \int_0^b f(x,y) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} dy dx = \frac{ab}{4} A_{mn}$$

$$\text{or } A_{mn} = \frac{4}{ab} \int_0^a \int_0^b f(x,y) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} dy dx \quad (vii)$$

Therefore from (v) required solution is

$$u(x,y,t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} \cos pt$$

where  $A_{mn}$  is given by (vii) and  $p = \pi c \sqrt{\left(\frac{m^2}{a^2} + \frac{n^2}{b^2}\right)}$

**Example 14:** Find the deflection  $u(x, y, t)$  of a square membrane with  $a = b = 1$  and  $c = 1$ , If the initial velocity is zero and the initial deflection is  $f(x,y) = A \sin \pi x \sin 2\pi y$

**Solution :** The deflection of the square membrane is given by the two dimensional wave equation.

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$$

The boundary conditions are

$$u(x, 0, t) = 0 = u(x, 1, t) \text{ and } u(0, y, t) = 0 = u(1, y, t)$$

The initial conditions are  $u(x, y, 0) = f(x, y) = A \sin \pi x \sin 2\pi y$ ,  $\left(\frac{\partial u}{\partial t}\right)_{t=0} = 0$

$\therefore$  Deflection


$$u(x,y,t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} \cos k_{mn} t \sin m\pi x. \sin n\pi y \quad (i)$$

where  $A_{mn} = 4 \int_0^1 \int_0^1 f(x,y) \sin m\pi x. \sin n\pi y dx dy \because a = b = 1, c = 1, k_{mn}^2 = \pi^2 (m^2 + n^2)$

$$= 4 \int_0^1 \int_0^1 (\sin \pi x)(\sin 2\pi y)(\sin m\pi x) \sin n\pi y dx dy$$



Applications of Partial Differential Equations



on integration, we find that  $A_{m1} = A_{m3} = A_{m4} = \dots = 0$

But

$$\begin{aligned} A_{m2} &= 4A \int_0^1 \int_0^1 (\sin \pi x)(\sin m\pi x) \sin^2 2\pi y \, dx \, dy \\ &= 2A \int_0^1 \int_0^1 (\sin \pi x)(\sin m\pi x)(1 - \cos 4\pi y) \, dx \, dy \\ &= 2A \int_0^1 \sin \pi x \sin m\pi x \left( y - \frac{1}{4\pi} \sin 4\pi y \right)_0^1 \, dx \\ &= 2A \int_0^1 \sin \pi x \sin m\pi x \, dx \end{aligned}$$

on integration we find that  $A_{22} = A_{32} = \dots = 0$

Also we find  $A_{12} = 2A \int_0^1 \sin \pi x \sin \pi x \, dx$

$$\begin{aligned} &= A \int_0^1 2 \sin^2 \pi x \, dx \\ &= A \int_0^1 (1 - \cos 2\pi x) \, dx \\ &= A \left( x - \frac{1}{2\pi} \sin 2\pi x \right)_0^1 = A \end{aligned}$$


Thus, from (1), we have

$$\begin{aligned} u(x, y, t) &= A_{12} (\cos k_{12}t) (\sin \pi x) \sin 2\pi y \\ &= A \cos \sqrt{5} \pi t \sin \pi x \sin 2\pi y \\ \therefore k_{12}^2 &= \pi^2 (1^2 + 2^2) \text{ i.e. } k_{12} = \pi\sqrt{5} \\ \text{since } c &= 1, m = 1, n = 2 \end{aligned}$$

**ONE - DIMENSIONAL HEAT FLOW**

Consider a homogeneous bar of uniform cross section A. Here we assume that the sides of the bar are insulated and the loss of heat from the sides by

conduction or radiation is negligible. Take one end of the bar as the origin and the direction of flow as the positive direction of  $x$  axis. Take  $k$  be the thermal conductivity 's' the specific heat and  $\rho$  be the density of the bar. The temperature  $u$  at any point of the bar depends on the distance  $x$  of the point from one end and the time  $t$ . The amount of heat crossing any section of the bar per second depends on the area  $A$  of the cross section, the rate of change of temperature with respect to ' $x$ ' (distance) normal to the area.



One dimensional heat flow

Therefore  $Q_1$ , the quantity of heat flowing into section at a distance  $x = -kA \left( \frac{\partial u}{\partial x} \right)_x$  per second. (The negative sign indicates that as  $x$  increases,  $u$  decreases).  $Q_2$ , the quantity of heat flowing out of the section at a distance  $x + \delta x = -kA \left( \frac{\partial u}{\partial x} \right)_{x+\delta x}$  per second.

Hence, the rate of increase of heat in the slab with thickness  $\delta x$  is

$$Q_1 - Q_2 = kA \left[ \left( \frac{\partial u}{\partial x} \right)_{x+\delta x} - \left( \frac{\partial u}{\partial x} \right)_x \right] \text{ per second} \quad \text{(i)}$$

But the rate of increase of heat of the slab =  $\rho A \delta x \frac{\partial u}{\partial t}$  (ii)

From (i) and (ii), we get

$$\rho A \delta x \frac{\partial u}{\partial t} = kA \left[ \left( \frac{\partial u}{\partial x} \right)_{x+\delta x} - \left( \frac{\partial u}{\partial x} \right)_x \right]$$

$$\text{or } \rho \frac{\partial u}{\partial t} = k \left[ \frac{\left( \frac{\partial u}{\partial x} \right)_{x+\delta x} - \left( \frac{\partial u}{\partial x} \right)_x}{\delta x} \right]$$

Applications of Partial Differential Equations

Taking limit as  $\delta x \rightarrow 0$ , we have

$$\begin{aligned} \rho s \frac{\partial u}{\partial t} &= k \frac{\partial^2 u}{\partial x^2} \\ \text{or } \frac{\partial u}{\partial t} &= \frac{k}{\rho s} \frac{\partial^2 u}{\partial x^2} \\ \text{or } \frac{\partial u}{\partial t} &= c^2 \frac{\partial^2 u}{\partial x^2} \quad (\text{iii}) \end{aligned}$$

where  $c^2 = \frac{k}{\rho s}$  is called the diffusivity of the material.

Equation (iii) is called the one-dimensional heat flow equation.

**Solution of one-dimensional heat Equation**

The equation of one dimensional heat flow is  $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$  (i)

Assume that a solution of (i) is

$$u = X(x) \cdot T(t)$$

where X is a function of x alone and T is a function t alone.

Then  $\frac{\partial u}{\partial t} = XT'$ ,  $\frac{\partial u}{\partial x} = X'T$  and  $\frac{\partial^2 u}{\partial x^2} = X''T$

Putting these values in (i), we get

$$XT' = C^2 X''T \Rightarrow \frac{X''}{X} = \frac{T'}{c^2 T} \quad (\text{ii})$$

Now L.H.S expression is a function of x only while R.H.S is a function of t only, so the two can be equal only when these are equal to a constant, say k.

$$\Rightarrow \frac{X''}{X} = k \quad \text{i.e. } X'' = kX$$

$$\text{or } \frac{d^2 X}{dx^2} - kX = 0 \quad (\text{iii})$$

$$\text{and } \frac{dT}{dt} - c^2 kT = 0 \quad (\text{iv})$$

There are arise following cases:

**Case I.** when  $k > 0$ , let  $k = p^2$

$$\text{Then } X = c_1 e^{px} + c_2 e^{-px} \text{ and } T = c_3 e^{-c^2 p^2 t}$$

$$\text{Therefore } u(x, t) = (c_1 e^{px} + c_2 e^{-px}) c_3 e^{-c^2 p^2 t}$$

**Case II.** when  $k < 0$ , let  $k = -p^2$

Then  $X = c_4 \cos px + c_5 \sin px$  and  $T = c_6 e^{-c^2 p^2 t}$

Therefore,  $u(x, t) = (c_4 \cos px + c_5 \sin px) c_6 e^{-c^2 p^2 t}$

**Case III** when  $k=0$ , then

$X = c_7 x + c_8$  and  $T = c_9$

Therefore  $u(x, t) = (c_7 x + c_8) c_9$

Here we are dealing with the heat conduction problem so the temperature  $u$  decreases with the increase of time  $t$  and hence solution given by case II is appropriate.

i.e.  $u(x, t) = (c_1 \cos px + c_2 \sin px) c_3 e^{-c^2 p^2 t}$  is the only suitable solution of the heat equation.

**Fourier series solution of one-dimensional heat equation**

Applying the boundary conditions and the initial condition, we get

$$u(x, t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} e^{-\frac{n^2 \pi^2 c^2 t}{l}}$$

where  $b_n = \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$

**Remark:** In steady state  $\frac{\partial u}{\partial t} = 0$ , so  $\frac{\partial^2 u}{\partial x^2} = 0$

**Example 15** An insulated rod of length  $l$  has its end A and B maintained  $0^\circ\text{C}$  and  $100^\circ\text{C}$  respectively until steady state condition Prevail. If B is suddenly reduced to  $0^\circ\text{C}$  and maintained at  $0^\circ\text{C}$  find the temperature at a distance  $x$  from A at time  $t$ .

(U.P.T.U.2004, 2005)

**Solution:** The equation of one dimensional heat flow be

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} \tag{i}$$

The boundary conditions are

(a)  $u(0, t) = 0^\circ\text{C}$  and (b)  $u(l, t) = 100^\circ\text{C}$

In steady state condition  $\frac{\partial u}{\partial t} = 0$ , here from (i) we get

$$\frac{\partial^2 u}{\partial x^2} = 0$$

on integration, we get  $u(x) = c_1 x + c_2$  (ii)

where  $c_1$  and  $c_2$  are constants to be determined.

At  $x=0$ , from equation (ii), we have

Applications of Partial Differential Equations

$$0 = C_2$$

$$\text{and at } x = l, 100 = c_1 l + 0 \Rightarrow c_1 = \frac{100}{l}$$

∴ form (ii)

$$u(x) = \frac{100}{l} x \tag{iii}$$

Now the temperature at B is suddenly changed we have again transient state. if  $u(x, t)$  is the subsequent temperature function, the boundary conditions are (a)

$$u(0, t) = 0^\circ\text{C} \quad (b') \quad u(l, t) = 0^\circ\text{C} \quad \text{and the initial condition (c')} \quad u(x, 0) = \frac{100}{l} x$$

since the subsequent steady state function  $u_s(x)$  satisfies the equation

$$\frac{\partial^2 u_s}{\partial x^2} = 0$$

$$\text{or } \frac{d^2 u_s}{dx^2} = 0 \Rightarrow u_s(x) = c_3 x + c_4$$

$$\text{at } x = 0, \text{ we get } 0 = c_4$$

and at  $x = l$ , we get

$$0 = c_3 l + 0 \Rightarrow c_3 = 0$$

$$\text{Thus } u_s(x) = 0 \tag{iv}$$

If  $u_T(x, t)$  is the temperature in transient state then the temperature distribution in the rod  $u(x, t)$  can be expressed in the form  $u(x, t) = u_s(x) + u_T(x, t)$

$$\Rightarrow u(x, t) = u_T(x, t) \quad \text{since } u_s(x) = 0 \tag{v}$$

Again from heat equation we have

$$\frac{\partial u_T}{\partial t} = c^2 \frac{\partial^2 u_T}{\partial x^2} \tag{vi}$$

The solution of equation (6) is

$$u_T(x, t) = (c_4 \cos px + c_2 \sin px) c_3 e^{-c^2 p^2 t} \tag{vii}$$

$$\text{At } x = 0, u(0, t) = 0$$

$$\Rightarrow 0 = c_1 e^{-c^2 p^2 t} \Rightarrow c_1 = 0$$

From (vii), we get

$$u(x, t) = c_2 \sin px c_3 e^{-c^2 p^2 t} \tag{viii}$$

$$\text{Again at } x = l, u(l, t) = 0$$

$$\Rightarrow 0 = c_1 c_2 \sin pl . e^{-c^2 p^2 t}$$

$$\Rightarrow \sin pl = 0 = \sin n\pi$$

$$\Rightarrow p = \frac{n\pi}{l}$$

From (viii), we get

$$u(x, t) = c_2 c_3 \sin \frac{n\pi x}{l} \cdot e^{-\frac{n^2 c^2 \pi^2 t}{l^2}}$$

$$\Rightarrow u(x, t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \cdot e^{-\frac{n^2 c^2 \pi^2 t}{l^2}} \quad \text{(ix)}$$

using initial condition i.e. at  $t = 0$ ,  $u = \frac{100}{l}x$ , we get

$$u(x, 0) = \frac{100}{l}x = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$$

$$\therefore b_n = \frac{2}{l} \int_0^l \frac{100}{l} x \sin \frac{n\pi x}{l} dx$$

$$= \frac{200}{l^2} \int_0^l x \sin \frac{n\pi x}{l} dx = \frac{200}{l^2} \left[ -\frac{xl}{n\pi} \cos \frac{n\pi x}{l} + \frac{l^2}{n^2 \pi^2} \sin \frac{n\pi x}{l} \right]_0^l$$

$$\Rightarrow b_n = \frac{200}{l^2} \left[ -\frac{l^2}{n\pi} \cos n\pi \right] = \frac{200}{n\pi} (-1)^{n+1}$$

Hence from equation (ix) we get

$$u(x, t) = \frac{200}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin \frac{n\pi x}{l} \cdot e^{-\frac{n^2 c^2 \pi^2 t}{l^2}}$$

**Example 16:** Determine the solution of one dimensional heat equation  $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$  subjected to the boundary conditions  $u(0, t) = 0$ ,  $u(l, t) = 0$  ( $t > 0$ ) and the initial condition  $u(x, 0) = x$ ,  $l$  being the length of the bar.

(I.A.S. 2007, U.P.T.U. 2006)

**Solution:** We have

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} \quad \text{(i)}$$

we know that the solution of equation (i) is given by

$$u(x, t) = (c_1 \cos px + c_2 \sin px) c_3 e^{-p^2 c^2 t} \quad \text{(ii)}$$

$$\text{At } x=0, u=0$$

$$\Rightarrow 0 = c_1 c_3 e^{-p^2 c^2 t}$$

$$\Rightarrow c_1 = 0$$

Applications of Partial Differential Equations

From (ii) we get

$$u(x, t) = c_2 c_3 \sin px e^{-p^2 c^2 t} \quad \text{(iii)}$$

Again at  $x=l, u=0$

$$\Rightarrow 0 = c_2 c_3 \sin pl . e^{-p^2 c^2 t}$$

$$\Rightarrow \sin pl = 0 = \sin n\pi$$

$$\Rightarrow p = \frac{n\pi}{l}$$

From (iii) the general solution of equation (i) is

$$u(x, t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} . e^{-\frac{n^2 c^2 \pi^2 t}{l^2}} \quad \text{(iv)}$$

At  $t=0, u=x$

$$\Rightarrow x = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$$

$$\therefore b_n = \frac{2}{l} \int_0^l x \sin \frac{n\pi x}{l} dx = \frac{2}{l} \left[ x \cdot \frac{l}{n\pi} \left( -\cos \frac{n\pi x}{l} \right) - \left( \frac{-l^2}{n^2 \pi^2} \sin \frac{n\pi x}{l} \right) \right]_0^l$$

$$= \frac{2}{l} \left[ \left( l \cdot \frac{l}{n\pi} (-\cos n\pi) + \frac{l^2}{n^2 \pi^2} \sin n\pi \right) - 0 \right]$$

$$\Rightarrow b_n = \frac{2}{l} \left[ -\frac{l^2}{n\pi} (-1)^n \right] = (-1)^{n+1} \frac{2l}{n\pi}$$

putting the value of  $b_n$  in equation (iv), we get

$$u(x, t) = \frac{2l}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin \frac{n\pi x}{l} . e^{-\frac{n^2 c^2 \pi^2 t}{l^2}}$$

**Example 17:** The ends A and B a rod 20 cm long have the temperature at 30°C and 80°C until steady state prevails. The temperature of the ends are changed to 40°C and 60°C respectively. Find the temperature distribution in the rod at time  $t$ . (I.A.S. 2005)

**Solution :** The heat equation in one dimensional is

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} \quad \text{(i)}$$

The boundary conditions are

$$(a) u(0, t) = 30^\circ\text{C} \quad (b) u(20, t) = 80^\circ\text{C}$$

In steady condition  $\frac{\partial u}{\partial t} = 0$

∴ From (i) we get  $\frac{\partial^2 u}{\partial x^2} = 0$ , on integration, we get

$$u(x) = c_1 x + c_2 \tag{ii}$$

at  $x = 0, u = 30$  so  $30 = 0 + c_2 \Rightarrow c_2 = 30$

and at  $x = 20, u = 80$  so  $80 = c_1 \cdot 20 + 30 \Rightarrow c_1 = \frac{5}{2}$

From equation (ii) we get

$$u(x) = \frac{5x}{2} + 30 \tag{iii}$$

Now the temperature at A and B are suddenly changed we have again gain transient state.

If  $u_1(x, t)$  is subsequent temperature function then the boundary conditions are

$$u_1(0, t) = 40^\circ\text{C} \text{ and } u_1(20, t) = 60^\circ\text{C}$$

and the initial condition i.e. at  $t = 0$ , is given by (iii)

Since the subsequent steady state function  $u_s(x)$  satisfies the equation

$$\frac{\partial^2 u_s}{\partial x^2} = 0 \quad \text{or} \quad \frac{d^2 u_s}{dx^2} = 0$$

The solution of above equation is

$$u_s(x) = c_3 x + c_4$$

At  $x = 0, u_s = 40 \Rightarrow 40 = 0 + c_4 \Rightarrow c_4 = 40 \quad \because u_s(0) = 40^\circ\text{C}$   
 $u_s(20) = 60^\circ\text{C}$

and at  $x = 20, u_s = 60 \Rightarrow 60 = 20 c_3 + 40 \Rightarrow c_3 = 1$

∴ from (iv), we get

$$u_s(x) = x + 40 \tag{v}$$

Thus the temperature distribution in the rod at time  $t$  is given by

$$u(x, t) = u_s(x) + u_T(x, t)$$

$$\Rightarrow u(x, t) = (x + 40) + u_T(x, t) \tag{vi}$$

where  $u_T(x, t)$  is the transient state function which satisfying the conditions

$$u_T(0, t) = u_1(0, t) - u_s(0) = 40 - 40 = 0$$

$$u_T(20, t) = u_1(20, t) - u_s(20) = 60 - 60 = 0$$

$$\text{and } u_T(x, 0) = u_1(x, 0) - u_s(x)$$

$$= \frac{5x}{2} + 30 - x - 40 = \frac{3x}{2} - 10$$

The general solution for  $u_T(x, t)$  is given by



Applications of Partial Differential Equations

$$u_T(x, t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{20} \cdot e^{-\frac{n^2 c^2 \pi^2 t}{l^2}} \quad \text{(vii)}$$

At  $t=0$ , from (vii) we get

$$\begin{aligned} \frac{3x}{2} - 10 &= \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{20} \\ \therefore b_n &= \frac{2}{20} \int_0^{20} \left( \frac{3x}{2} - 10 \right) \sin \frac{n\pi x}{20} dx \\ &= \frac{1}{10} \left[ \left( \frac{3x}{2} - 10 \right) \left( -\frac{20}{n\pi} \cos \frac{n\pi x}{20} \right) - \frac{3}{2} \left( -\frac{400}{n^2 \pi^2} \cos \frac{n\pi x}{20} \right) \right]_0^{20} \\ &= \frac{1}{10} \left[ -20 \left( \frac{20}{n\pi} \right) (-1)^n - (-10) \left( \frac{20}{n\pi} \right) \right] \\ &= -\frac{20}{n\pi} [2(-1)^n + 1] \end{aligned}$$

putting the value of  $b_n$  in equation (vii), we get


$$u_T(x, t) = -\frac{20}{\pi} \sum_{n=1}^{\infty} \frac{2(-1)^n + 1}{n} \sin \frac{n\pi x}{20} \cdot e^{-\frac{n^2 \pi^2 c^2 t}{400}} \quad \text{(viii)}$$

From (vi) and (viii), we get

$$u_T(x, t) = (x + 40) - \frac{20}{\pi} \sum_{n=1}^{\infty} \frac{2(-1)^n + 1}{n} \sin \frac{n\pi x}{20} \cdot e^{-\frac{n^2 \pi^2 c^2 t}{400}}$$

**TWO - DIMENSIONAL HEAT FLOW**

Consider the flow of heat in a metal plate of inform thickness  $\alpha$  (cm), density  $\rho$  (gr/cm<sup>3</sup>), specific heat  $s$ (cal/gr deg) and thermal conductivity  $k$  (cal/cm sec-deg) Let XOY plane be taken in one face of the plates as shown in figure. If the temperature at any point is independent of the  $z$  coordinate and depends only on  $x$ ,  $y$  and time  $t$ , then the flow is said to be two-dimensional. In this case, the heat flow is in the XY plane only and is zero along the normal to the XY-plane.



Consider a rectangular element ABCD of the plane with sides  $\delta x$  and  $\delta y$ .

The amount of heat entering the element in 1 sec. from the side AB

$$= -k\alpha\delta x \left( \frac{\partial u}{\partial y} \right)_y \quad (\text{see one dimensional heat flow})$$

and the amount of heat entering the element in 1 second from the side AD

$$= -k\alpha\delta y \left( \frac{\partial u}{\partial x} \right)_x$$

The quantity of heat flowing out through the side CD per sec

$$= -k\alpha\delta x \left( \frac{\partial u}{\partial y} \right)_{y+\delta y}$$

and the quantity of heat flowing out through the side BC per second

$$= -k\alpha\delta y \left( \frac{\partial u}{\partial x} \right)_{x+\delta x}$$

Hence the total gain of heat by the rectangular element ABCD per second

$$\begin{aligned} &= -k\alpha\delta x \left( \frac{\partial u}{\partial y} \right)_y - k\alpha\delta y \left( \frac{\partial u}{\partial x} \right)_x + k\alpha\delta x \left( \frac{\partial u}{\partial y} \right)_{y+\delta y} + k\alpha\delta y \left( \frac{\partial u}{\partial x} \right)_{x+\delta x} \\ &= k\alpha\delta x \left[ \left( \frac{\partial u}{\partial y} \right)_{y+\delta y} - \left( \frac{\partial u}{\partial y} \right)_y \right] + k\alpha\delta y \left[ \left( \frac{\partial u}{\partial x} \right)_{x+\delta x} - \left( \frac{\partial u}{\partial x} \right)_x \right] \\ &= k\alpha\delta x\delta y \left[ \frac{\left( \frac{\partial u}{\partial x} \right)_{x+\delta x} - \left( \frac{\partial u}{\partial x} \right)_x}{\delta x} + \frac{\left( \frac{\partial u}{\partial y} \right)_{y+\delta y} - \left( \frac{\partial u}{\partial y} \right)_y}{\delta y} \right] \end{aligned} \quad (i)$$

Also the rate of gain of heat by the element

$$= \rho\delta x\delta y\alpha s \frac{\partial u}{\partial t} \quad (ii)$$

Thus from equation (i) and (ii)

$$k\alpha\delta x\delta y \left[ \frac{\left( \frac{\partial u}{\partial x} \right)_{x+\delta x} - \left( \frac{\partial u}{\partial x} \right)_x}{\delta x} + \frac{\left( \frac{\partial u}{\partial y} \right)_{y+\delta y} - \left( \frac{\partial u}{\partial y} \right)_y}{\delta y} \right] = \rho\delta x\delta y\alpha s \frac{\partial u}{\partial t}$$

Dividing both sides by  $\alpha \delta x \delta y$  and taking limits as  $\delta x \rightarrow 0$ ,  $\delta y \rightarrow 0$ , we get

$$k \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) = \rho s \frac{\partial u}{\partial t}$$

i.e.  $\frac{\partial u}{\partial t} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$  (iii)

where  $c^2 = \frac{k}{\rho s}$  is the diffusivity

Equation (iii) gives the temperature distribution of the plate in the transient state.

**Cor.** In the steady state,  $u$  is independent of  $t$ , so that  $\frac{\partial u}{\partial t} = 0$  and the above equation reduces to,

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

and this is called Laplace's equation in two dimensions.

#### **Solution of Laplace Equation in Two Dimensions**

Laplace equation in two dimensions is given by

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \tag{i}$$

Let  $u = XY$  be a solution of (i)

where  $X$  is a function of  $x$  alone and  $Y$  is the function of  $y$  alone.

$$\text{Then } \frac{\partial^2 u}{\partial x^2} = X''Y \quad \text{and} \quad \frac{\partial^2 u}{\partial y^2} = XY''$$

Substituting these values in (i), we get

$$X''Y + XY'' = 0$$

or  $\frac{X''}{X} = -\frac{Y''}{Y}$  (ii)

Now in equation (ii) variables are separable since  $x$  and  $y$  are independent variable, this equation can hold only when both sides reduce to a constant, say  $k$ .

$$\text{i.e. } \frac{X''}{X} = -\frac{Y''}{Y} = k$$
$$\Rightarrow \frac{d^2 X}{dx^2} - kX = 0 \quad \text{and} \quad \frac{d^2 Y}{dy^2} + KY = 0 \tag{iii}$$

Solving equation (iii), we get

(a) when  $k$  is positive say  $p^2$ , then we have

$$X = c_1 e^{px} + c_2 e^{-px}, \quad Y = c_3 \cos py + c_4 \sin py$$

(b) when  $k$  is negative say  $-p^2$ , then we have

$$X = c_5 \cos px + c_6 \sin px, \quad Y = c_7 e^{py} + c_8 e^{-py}$$

(c) when  $k = 0$ , then

$$X = c_9 x + c_{10}, \quad Y = c_{11} y + c_{12}$$

Thus the various possible solution of (i) are

$$u = (c_1 e^{px} + c_2 e^{-px}) (c_3 \cos py + c_4 \sin py) \tag{iv}$$

$$u = (c_5 \cos py + c_6 \sin py) (c_7 e^{py} + c_8 e^{-py}) \tag{v}$$

$$u = (c_9 x + c_{10}) (c_{11} y + c_{12}) \tag{vi}$$


of these we take that solution which is consistent with the given boundary conditions, i.e., physical nature of the problem.

**Example 18:** Solve the Laplace equation  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$  subjected to the conditions

$$u(0, y) = u(l, y) = u(x, 0) = 0 \text{ and } u(x, a) = \sin \frac{n\pi x}{l} \quad (\text{U.P.T.U 2004})$$

**Solution.** The three possible solution of

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \tag{i}$$



are

$$u = (c_1 e^{px} + c_2 e^{-px}) (c_3 \cos py + c_4 \sin py) \tag{ii}$$

$$u = (c_5 \cos px + c_6 \sin px) (c_7 e^{py} + c_8 e^{-py}) \tag{iii}$$

$$u = (c_9 x + c_{10}) (c_{11} y + c_{12}) \tag{iv}$$

we have to solve (i) satisfying the following boundary conditions

$$u(0, y) = 0 \tag{v}$$

$$u(l, y) = 0 \tag{vi}$$

Applications of Partial Differential Equations

$$u(x, 0) = 0 \tag{vii}$$

$$u(x, a) = \sin n\pi x/l \tag{viii}$$

using (v) and (vi) in (ii), we get

$$c_1 + c_2 = 0, \text{ and } c_1 e^{pl} + c_2 e^{-pl} = 0$$

Solving these equations, we get  $c_1 = c_2 = 0$ , which lead to trivial solution. similarly we get a trivial solution by using (v) and (vi) in (iv). Hence the suitable for the present problem is solution (iii), using (v) in (iii),

we have  $c_5 (c_7 e^{py} + c_8 e^{-py}) = 0$  i.e.  $c_5 = 0$

$$\therefore \text{(iii) becomes } u = c_6 \sin px (c_7 e^{py} + c_8 e^{-py}) \tag{ix}$$

using (vi), we have  $c_6 \sin pl (c_7 e^{py} + c_8 e^{-py}) = 0$

$$\therefore \text{either } c_6 = 0 \text{ or } \sin pl = 0$$

If we take  $c_6 = 0$ , we get trivial solution

Thus  $\sin pl = 0 \Rightarrow pl = n\pi$

$$\Rightarrow p = \frac{n\pi}{l}, \text{ where } n = 0, 1, 2, \dots$$

$$\therefore \text{(ix) becomes } u = c_6 \sin\left(\frac{n\pi x}{l}\right) (c_7 e^{n\pi y/l} + c_8 e^{-n\pi y/l}) \tag{x}$$

Using (vii), we have  $0 = c_6 \sin\left(\frac{n\pi x}{l}\right) (c_7 + c_8)$  i.e.  $c_8 = -c_7$

Thus the solution suitable for this problem is

$$u(x, y) = b_n \sin\frac{n\pi x}{l} (e^{n\pi y/l} - e^{-n\pi y/l}) \text{ where } b_n = c_6 c_7$$

Now using the condition (viii), we have

$$u(x, a) = \sin\frac{n\pi x}{l} = b_n \sin\frac{n\pi x}{l} (e^{n\pi a/l} - e^{-n\pi a/l})$$

$$\text{we get } b_n = \frac{1}{(e^{n\pi a/l} - e^{-n\pi a/l})}$$

Hence the required solution is

$$u(x, y) = \frac{e^{n\pi y/l} - e^{-n\pi y/l}}{e^{n\pi a/l} - e^{-n\pi a/l}} \sin\frac{n\pi x}{l}$$

$$\Rightarrow u(x, y) = \frac{\sinh(n\pi y/l)}{\sinh(n\pi a/l)} \sin\frac{n\pi x}{l}$$

**Example 19:** A thin rectangular plate whose surface is impervious to heat flow, has at  $t = 0$  an arbitrary distribute of temperature  $f(x, y)$ , if four edges  $x = 0, x = a, y$

$x=0, y=b$  are kept at zero temperature. determine the temperature at a point of the plate as  $t$  increases.

(U.P.T.U. 2002)

**Solution** The two dimensional heat equation is


$$\frac{\partial u}{\partial t} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) \quad (1)$$

$$\text{where } c^2 = \frac{k}{\sigma\rho}$$

The initial temperature of the plate is  $f(x, y)$  and the temperature of the four edges of the plate are kept at zero temperature.

Now the function  $u(x, y, t)$  is required to satisfy (i) and the boundary and initial conditions given below:

The boundary conditions are



$$u(0, y, t) = 0 \quad (i)$$

$$u(a, y, t) = 0 \quad (ii)$$

$$u(x, 0, t) = 0 \quad (iii)$$

$$u(x, b, t) = 0 \quad (iv)$$

and the initial condition is

$$u(x, y, 0) = f(x, y) \quad (2)$$

Let the solution of the heat equation (1) be of the form

$$u(x, y, t) = X(x) Y(y) T(t) = XYT(\text{say}) \quad (3)$$

where  $X$  is a function of  $x$  only,  $Y$  is that of  $y$  only and  $T$  is that of  $t$  only.

using (3) in (1) we get

$$\frac{1}{c^2 T} \frac{dT}{dt} = \frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2}$$

Hence in order that (3) may satisfy (1) we have of these three possibilities :

Applications of Partial Differential Equations

$$(A) \frac{1}{X} \frac{d^2X}{dx^2} = 0, \quad \frac{1}{Y} \frac{d^2Y}{dy^2} = 0, \quad \frac{1}{c^2T} \frac{dT}{dt} = 0$$

$$(B) \frac{1}{X} \frac{d^2X}{dx^2} = p_1^2, \quad \frac{1}{Y} \frac{d^2Y}{dy^2} = p_2^2, \quad \frac{1}{c^2T} \frac{dT}{dt} = p^2$$

$$(C) \frac{1}{X} \frac{d^2X}{dx^2} = -p_1^2, \quad \frac{1}{Y} \frac{d^2Y}{dy^2} = -p_2^2, \quad \frac{1}{c^2T} \frac{dT}{dt} = -p^2$$

where  $p^2 = p_1^2 + p_2^2$

It can be easily observed that differential equation (C) only gives the solution for this situation, and the general of the differential equation in this case is

$$X = A_1 \cos p_1 x + B_1 \sin p_1 x$$

$$Y = A_2 \cos p_2 y + B_2 \sin p_2 y \text{ and } T = A_3 e^{-c^2 p^2 t}$$

$$\Rightarrow u(x, y, t) = (A_1 \cos p_1 x + B_1 \sin p_1 x) (A_2' \cos p_2 y + B_2' \sin p_2 y) e^{-c^2 p^2 t} \quad (4)$$

where  $A_2' = A_2 A_3$ ,  $B_2' = B_2 B_3$

Under boundary condition (i) we get

$$u(0, y, t) = A_1 (A_2' \cos p_2 y + B_2' \sin p_2 y) e^{-c^2 p^2 t} = 0$$

$$\Rightarrow A_1 = 0$$

Again using (ii)

$$u(a, y, t) = B_1 \sin p_1 a (A_2' \cos p_2 y + B_2' \sin p_2 y) e^{-c^2 p^2 t} = 0$$

$$\sin p_1 a = 0 \Rightarrow p_1 a = m\pi \quad \Rightarrow \quad p_1 = \frac{m\pi}{a} \text{ where } m = 1, 2, 3, \dots$$

similarly, making use of (iii) and (iv), we obtain

$$A_2' = 0 \text{ and } p_2 = \frac{n\pi}{b} \text{ (} n = 1, 2, 3, \dots \text{)}$$

Thus we have

$$u_{mn}(x, y, t) = A_{mn} e^{-c^2 p_{mn}^2 t} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b}$$

$$\text{where } p^2 = p_{mn}^2 = \pi^2 \left( \frac{m^2}{a^2} + \frac{n^2}{b^2} \right)$$

$$\Rightarrow u(x, y, t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} e^{-c^2 p_{mn}^2 t} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} \quad (5)$$

This solution satisfies the boundary conditions.

Now to derive the solution which satisfies the initial condition also, we proceed as follows:

$$\Rightarrow u(x, y, 0) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} e^{-c^2 p_{mn}^2 t} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} = f(x, y)$$

Hence, the L.H.S is the double Fourier sine series of  $f(x, y)$

$$\Rightarrow A_{mn} = \frac{2}{a} \cdot \frac{2}{b} \int_{x=0}^a \int_{y=0}^b f(x, y) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} dx dy \quad (6)$$

Hence the required solution of heat equation (1) is given by (5) with coefficients given by (6)

**Example 20** Solve the P.D.E by separation of variables method  $u_{xx} = u_y + 2u$ ,  $u(0, y) = 0$

$$\frac{\partial}{\partial x} u(0, y) = 1 + e^{-3y}$$

(U.P.T.U. 2009)

**Solution Hint.**  $\frac{X''}{X} = \frac{Y' + 2Y}{Y} = K$  (say)

$$\Rightarrow X = c_1 e^{\sqrt{k}x} + c_2 e^{-\sqrt{k}x}, Y = c_3 e^{(k-2)y}$$

$$u = XY$$

$$\text{Therefore } u(x, y) = \frac{1}{\sqrt{2}} \cdot \sinh \sqrt{2}x + \sin e^{-3y}$$

### EXERCISE

Solve the following P.D.E. by the method of separation of variables.

1.  $3 \frac{\partial u}{\partial x} + 2 \frac{\partial u}{\partial y} = 0$  given  $u(x, 0) = 4e^{-x}$

Ans.  $u(x, y) = 4e^{\frac{1}{2}(3y-2x)}$

2.  $\frac{\partial^2 z}{\partial x^2} + 4 \frac{\partial^2 z}{\partial y^2} = 0$

Ans.  $z = (c_1 e^{\sqrt{k}x} + c_2 e^{-\sqrt{k}x}) \left( c_3 \sin \frac{1}{2} \sqrt{k}y + c_4 \cos \frac{1}{2} \sqrt{k}y \right)$

3.  $\frac{\partial u}{\partial x} + 2 \frac{\partial u}{\partial t} + u, u(x, 0) = 6e^{-3x}$

(U.P.T.U. 2006)

Ans.  $u(x, t) = 6e^{(-3x + 2t)}$



Applications of Partial Differential Equations

4.  $2 \frac{\partial u}{\partial x} - 3 \frac{\partial u}{\partial y} = 0, u(x, 0) = 5e^{3x}$

Ans.  $u(x, y) = 5e^{3x + 2y}$

5.  $3 \frac{\partial u}{\partial x} + 2 \frac{\partial u}{\partial y} = 0$

(U.P.T.U. 2009)

Ans.  $u(x, y) = ce^{\frac{1}{6}k(2x-3y)}$

6. The vibrations of an elastic string is governed by the P.D.E.  $\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2}$ . The length of the string is  $\pi$  and the ends are fixed. The initial velocity is zero and the initial deflection is  $u(x, 0) = 2(\sin x + \sin 3x)$ . Find the deflection  $u(x, t)$  of the vibrating string for  $t > 0$ .

Ans.  $u(x, t) = 4 \cos x \cos 2t \sin 2x$

7. find the displacement of a string stretched between the fixed points  $(0, 0)$  and  $(1, 0)$  and released from rest from position  $a \sin \pi x + \cos 2\pi x$ .

Ans.  $u(x, t) = a \sin \pi x \cos \pi ct + b \sin 2\pi x \cos 2\pi ct$

8. Solve the wave equation  $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$  subjected to the conditions

(i)  $u(0, t) = u(l, t) = 0 \quad t \geq 0$

(ii)  $u(x, 0) = \begin{cases} \frac{2x}{l}, & 0 \leq x \leq \frac{l}{2} \\ \frac{2}{l}(l-x), & \frac{l}{2} \leq x \leq l \end{cases}$  (I.A.S 2006)

(iii)  $\left(\frac{\partial u}{\partial t}\right)_{t=0} = 0, \quad 0 \leq x \leq l$

Ans.  $u(x, t) = \frac{8}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} \sin \frac{n\pi}{2} \sin \frac{n\pi x}{l} \cos \frac{n\pi ct}{l}$

9. A tant string of length  $2l$  is fastened at both ends. The mid point of the string is taken to a height  $b$  and then released from the rest in that position. Find the displacement of the string.

Ans.  $y(x, t) = \frac{8b}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} \sin(2n-1) \frac{\pi}{2} \sin \frac{(2n-1)\pi x}{2l} \cos \frac{(2n-1)\pi ct}{2l}$

10. find the temperature  $u(x, t)$  in a homogeneous bar of heat conduction material of length  $l$  cm with its ends kept at zero temperature and initial temperature is  $dx(l-x)/l^2$

$$\text{Ans. } u(x, t) = \frac{8d}{\pi^3} \sum_{n=1}^{\infty} \frac{1}{(2n-1)^3} \sin \frac{(2n-1)\pi x}{2l} e^{-(2n-1)^2 \pi^2 c^2 t / l^2}$$

11. Solve the following Laplace equation

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

in a rectangle with  $u(0, y) = 0$ ,  $u(a, y) = 0$ ,  $u(x, b) = 0$  and  $u(x, 0) = f(x)$  along  $x$  axis.

(U.P.T.U. 2008)

$$\text{Ans. } u(x, y) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{a} \left\{ e^{\frac{n\pi y}{a}} - e^{-\frac{n\pi}{a}(y-2b)} \right\}$$

$$\text{where } b_n = \frac{2}{a \left( 1 - e^{-\frac{2n\pi b}{a}} \right)} \int_0^a f(x) \sin \frac{n\pi x}{a} dx$$

12. Find the temperature in a bar of length 2 whose end are kept at zero and lateral surface is insulated if the initial temperature is  $\sin \frac{n\pi x}{2} + 3 \sin \frac{5\pi x}{2}$

(U.P.T.U. 2009)

### Solution

**Hint.** Heat equation  $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$

$$u(x, t) = (c_1 \cos px + c_2 \sin px) (c_3 e^{-c^2 p^2 t})$$

$$u(0, t) = 0 \Rightarrow c_1 = 0$$

$$u(l, t) = 0 \Rightarrow p = \frac{n\pi}{l}$$

$$u(x, t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} e^{-\frac{c^2 n^2 \pi^2 t}{l^2}}$$

$$\text{where } b_n = \int_0^2 \left( \sin \frac{n\pi x}{2} + 3 \sin \frac{5\pi x}{2} \right) \sin \frac{n\pi x}{2} dx$$

**Choose the correct answer from the following:**

1. One dimensional wave equation  $\frac{\partial^2 z}{\partial t^2} = c^2 \frac{\partial^2 z}{\partial x^2}$  is:

Applications of Partial Differential Equations

- (a) elliptic    (b) parabolic    (c) hyperbolic    (d) circular.  
Ans. (c)

2. One dimensional heat flow equation  $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$  is:

- (a) circular    (b) hyperbolic    (c) parabolic    (d) elliptic.  
Ans. (c)

3. The two dimensional heat flow equation in steady state  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$  is:

- (a) elliptic    (b) circular    (c) parabolic    (d) hyperbolic.  
Ans. (a)

4. The differential equation  $4 \frac{\partial^2 u}{\partial x^2} + 4 \frac{\partial^2 u}{\partial x \partial y} + \frac{\partial^2 u}{\partial y^2} = 0$  is:

- (a) parabolic    (b) elliptic    (c) hyperbolic    (d) circular  
Ans. (a)

5. The general solution of  $\frac{\partial^2 u}{\partial x \partial y} = 0$  is:

- (a)  $u = f_1(x+y) + f_2(y)$     (b)  $u = f_1(x) + f_2(y)$   
(c)  $u = f(xy)$     (d)  $u = f_1(xy) + f_2(y)$ .  
Ans. (a)

6. The partial differential equation  $\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial y^2}$  is:

- (a) hyperbolic    (b) parabolic    (c) elliptic    (d) circular.  
Ans. (a)

7. The partial differential equation  $9 \frac{\partial^2 u}{\partial x^2} - 6 \frac{\partial^2 u}{\partial x \partial y} + \frac{\partial^2 u}{\partial y^2} = 0$  is:

- (a) hyperbolic    (b) elliptic    (c) circular    (d) parabolic  
Ans. (d)

8. The two dimensional heat equation in the transient state is:

- (a)  $\frac{\partial u}{\partial t} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$     (b)  $\frac{\partial^2 u}{\partial t^2} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$

(c)  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = c$                       (d)  $\frac{\partial u}{\partial t} = c \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial u}{\partial y} \right)$

Ans. (a)

9. The two dimensional wave equation  $\frac{\partial^2 u}{\partial t^2} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial u}{\partial y^2} \right)$  is:

- (a) circular    (b) elliptic                      (c) parabolic                      (d) hyperbolic.

Ans. (d)

10. The Laplace's equation in polar coordinates is:

(a)  $\frac{\partial^2 u}{\partial r^2} + \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$                       (b)  $\frac{\partial^2 u}{\partial r^2} - \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$   
 (c)  $\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$                       (d)  $\frac{\partial^2 u}{\partial r^2} - \frac{\partial u}{\partial r} + \frac{1}{r} \frac{\partial^2 u}{\partial \theta^2} = 0.$

Ans. (c)

11. The radio equations are:

(a)  $\frac{\partial^2 v}{\partial x^2} = LC \frac{\partial^2 v}{\partial t^2}$  and  $\frac{\partial^2 i}{\partial x^2} = LC \frac{\partial i}{\partial t}$     (b)  $\frac{\partial^2 v}{\partial x^2} = LC \frac{\partial^2 v}{\partial t^2}$  and  $\frac{\partial i}{\partial x} = LC \frac{\partial v}{\partial t}$   
 (c)  $\frac{\partial^2 v}{\partial x^2} = LC \frac{\partial^2 v}{\partial t^2}$  and  $\frac{\partial^2 i}{\partial x^2} = LC \frac{\partial^2 i}{\partial t^2}$     (d) none of these.

Ans. (c)

12. The telegraph equations are:

(a)  $\frac{\partial^2 v}{\partial x^2} = RC \frac{\partial v}{\partial t}$  and  $\frac{\partial^2 i}{\partial x^2} = RC \frac{\partial i}{\partial t}$     (b)  $\frac{\partial v}{\partial x} = RC \frac{\partial v}{\partial t}$  and  $\frac{\partial i}{\partial x} = RC \frac{\partial i}{\partial t}$   
 (c)  $\frac{\partial^2 v}{\partial x^2} = RC \frac{\partial^2 v}{\partial t^2}$  and  $\frac{\partial^2 i}{\partial x^2} = RC \frac{\partial^2 i}{\partial t^2}$     (d) none of these.

Ans. (a)

13. The equations for submarine cable are:

(a)  $\frac{\partial v}{\partial x} = RGv$  and  $\frac{\partial i}{\partial x} = RGi$                       (b)  $\frac{\partial^2 v}{\partial x^2} = RGv$  and  $\frac{\partial^2 i}{\partial x^2} = RGi$   
 (c)  $\frac{\partial^2 v}{\partial x^2} = RG$  and  $\frac{\partial^2 i}{\partial x^2} = RG$                       (d)  $\frac{\partial^2 v}{\partial x^2} = Rv$  and  $\frac{\partial^2 i}{\partial x^2} = Ri.$

Ans. (a)

Applications of Partial Differential Equations

**Fill in the blanks in the following problems:**

1. The partial differential equation  $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$  is of.....type.

Ans. Parabolic

2. When a vibrating string has an initial velocity, its initial conditions are.....

Ans.  $\left(\frac{\partial y}{\partial t}\right)_{t=0} = v$

3. The solution of  $\frac{\partial^2 u}{\partial x \partial y} = 0$  is of the form.

Ans.  $u = f_1(y) + f_2(x)$

4. The equation  $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$  is classified as.....

Ans. Elliptic partial differential equation

5. In two dimensional heat flow, the temperature along the normal to the  $xy$ -plane is.....

Ans. Zero

6. D' Alembert's solution of the wave equation  $\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$  is.....

Ans.  $y(x, t) = \frac{1}{2} [f(x + ct) + f(x - ct)]$

**Select True 'or' False answers in the following:**

7. The solution of  $\frac{\partial^2 z}{\partial x^2} = \sin(xy)$  is  $-y^2 \sin(xy) + x f_1(x) + f_2(y)$ .

(False)

8. The general solution of the equation  $\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial t^2} = 0$  is  $(C_1 \cos px + C_2 \sin px)(C_3 \cos pt + C_4 \sin pt)$ .

(True)

9. The two dimensional heat flow in transient state is  $\frac{\partial u}{\partial t} = C^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$ .

(True)

10. The telegraph equations are  $\frac{\partial^2 v}{\partial x^2} = RC \frac{\partial v}{\partial t}$  and  $\frac{\partial^2 i}{\partial x^2} = RC \frac{\partial i}{\partial t}$   
(True)
11. The radio equations are  $\frac{\partial v}{\partial x^2} = LC \frac{\partial^2 v}{\partial t^2}$  and  $\frac{\partial i}{\partial x} = LC \frac{\partial^2 i}{\partial t^2}$   
(False)
12. The Laplace's equation in three dimensions is  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial u}{\partial z} = 0$   
(False)